

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	SNF1	SNF2	p	test
n		497	128		
Subtype (%)	SNF1	497 (100.0)	0 (0.0)	<0.001	
	SNF2	0 (0.0)	128 (100.0)		
vital status (%)	Alive	132 (93.6)	31 (93.9)	1.000	exact
	Dead	9 (6.4)	2 (6.1)		
days to birth (median [IQR])		-21534.50 [-24923.00, -17939.75]	-20151.00 [-23467.25, -17744.50]	0.077	nonnorm
days to death (median [IQR])		1590.50 [642.75, 2406.00]	848.50 [372.50, 2688.50]	0.483	nonnorm
days to last followup (median [IQR])		754.00 [476.50, 1611.00]	789.00 [463.50, 1592.00]	0.979	nonnorm
race list (%)	AMERICAN INDIAN OR ALASKA NATIVE	0 (0.0)	1 (0.8)	<0.001	exact
	ASIAN	26 (5.3)	6 (4.8)		
	BLACK OR AFRICAN AMERICAN	79 (16.3)	45 (35.7)		
	WHITE	381 (78.4)	74 (58.7)		
history of neoadjuvant treatment (%)	No	490 (98.8)	127 (100.0)	0.607	exact
	Yes	6 (1.2)	0 (0.0)		
age at initial pathologic diagnosis (median [IQR])		58.00 [48.00, 67.00]	55.00 [48.00, 63.50]	0.105	nonnorm
ethnicity (%)	HISPANIC OR LATINO	20 (4.5)	7 (6.0)	0.662	
	NOT HISPANIC OR LATINO	427 (95.5)	110 (94.0)		
primary lymph node presentation assessment (%)	NO	21 (5.3)	5 (4.9)	1.000	
	YES	376 (94.7)	98 (95.1)		
histological type (%)	Infiltrating Carcinoma NOS	0 (0.0)	1 (0.8)	<0.001	exact
	Infiltrating Ductal Carcinoma	331 (66.6)	105 (82.7)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	SNF1	SNF2	p	test
menopause status (%)	Infiltrating Lobular Carcinoma	116 (23.3)	2 (1.6)	0.039	exact
	Medullary Carcinoma	0 (0.0)	5 (3.9)		
	Metaplastic Carcinoma	1 (0.2)	6 (4.7)		
	Mixed Histology (please specify)	21 (4.2)	0 (0.0)		
	Mucinous Carcinoma	10 (2.0)	0 (0.0)		
	Other, specify	18 (3.6)	8 (6.3)		
	Indeterminate (neither Pre or Postmenopausal)	9 (2.0)	5 (4.2)		
	Peri (6-12 months since last menstrual period)	13 (2.9)	7 (5.9)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	318 (69.7)	87 (73.7)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	116 (25.4)	19 (16.1)		
breast carcinoma progesterone receptor status (%)	Indeterminate	1 (0.2)	1 (0.8)	<0.001	exact
breast carcinoma estrogen receptor status (%)	Negative	75 (16.1)	109 (90.1)	<0.001	
	Positive	389 (83.7)	11 (9.1)		
	Negative	28 (6.0)	102 (83.6)		
lab proc her2 neu immuno-histochemistry receptor status (%)	Positive	437 (94.0)	20 (16.4)	0.190	exact
	Equivocal	89 (22.1)	21 (20.0)		
	Indeterminate	7 (1.7)	3 (2.9)		
	Negative	244 (60.7)	72 (68.6)		
	Positive	62 (15.4)	9 (8.6)		
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.020	nonnorm

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	SNF1	SNF2	p	test
number of lymphnodes positive by he (median [IQR])		1.00 [0.00, 3.00]	0.00 [0.00, 2.00]	<0.001	nonnorm
er level cell percentage category (%)	<10%	10 (3.9)	25 (73.5)	<0.001	exact
	10-19%	19 (7.4)	1 (2.9)		
	20-29%	2 (0.8)	2 (5.9)		
	30-39%	4 (1.6)	1 (2.9)		
	40-49%	3 (1.2)	0 (0.0)		
	50-59%	8 (3.1)	0 (0.0)		
	60-69%	7 (2.7)	0 (0.0)		
	70-79%	39 (15.1)	1 (2.9)		
	80-89%	18 (7.0)	0 (0.0)		
	90-99%	148 (57.4)	4 (11.8)		
progesterone receptor level cell percent category (%)	<10%	51 (21.9)	28 (93.3)	<0.001	exact
	10-19%	18 (7.7)	0 (0.0)		
	20-29%	8 (3.4)	0 (0.0)		
	30-39%	8 (3.4)	0 (0.0)		
	40-49%	11 (4.7)	0 (0.0)		
	50-59%	10 (4.3)	0 (0.0)		
	60-69%	11 (4.7)	1 (3.3)		
	70-79%	29 (12.4)	0 (0.0)		
	80-89%	23 (9.9)	0 (0.0)		
	90-99%	64 (27.5)	1 (3.3)		
distant metastasis present ind2 (%)	NO	210 (97.7)	43 (97.7)	1.000	exact
	YES	5 (2.3)	1 (2.3)		
stage event pathologic stage (%)	Stage I	41 (8.3)	9 (7.1)	0.032	exact
	Stage IA	40 (8.1)	9 (7.1)		
	Stage IB	3 (0.6)	0 (0.0)		
	Stage II	4 (0.8)	1 (0.8)		
	Stage IIA	141 (28.6)	61 (48.0)		
	Stage IIB	123 (24.9)	25 (19.7)		
	Stage III	2 (0.4)	0 (0.0)		
	Stage IIIA	87 (17.6)	13 (10.2)		
	Stage IIIB	12 (2.4)	2 (1.6)		
	Stage IIIC	32 (6.5)	4 (3.1)		
	Stage IV	5 (1.0)	2 (1.6)		
	Stage X	3 (0.6)	1 (0.8)		