

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	MOFA1	MOFA2	MOFA3	p	test
n		342	96	187		
Subtype (%)	MOFA1	342 (100.0)	0 (0.0)	0 (0.0)	<0.001	
	MOFA2	0 (0.0)	96 (100.0)	0 (0.0)		
	MOFA3	0 (0.0)	0 (0.0)	187 (100.0)		
vital status (%)	Alive	94 (94.9)	26 (92.9)	43 (91.5)	0.616	exact
	Dead	5 (5.1)	2 (7.1)	4 (8.5)		
days to birth (median [IQR])		-21310.50 [- 24324.00, -17687.75]	-19621.50 [- 24058.50, -17708.50]	-22189.00 [- 25123.25, -18528.50]	0.118	nonnorm
days to death (median [IQR])		1365.00 [574.00, 2235.00]	912.00 [524.00, 1759.00]	1910.50 [685.50, 2612.75]	0.566	nonnorm
days to last followup (median [IQR])		856.00 [492.00, 1783.00]	730.00 [456.75, 1565.50]	664.00 [446.50, 1347.50]	0.117	nonnorm
race list (%)	AMERICAN INDIAN OR ALASKA NATIVE	0 (0.0)	0 (0.0)	1 (0.5)	<0.001	exact
	ASIAN	15 (4.5)	4 (4.3)	13 (7.1)		
	BLACK OR AFRICAN AMERICAN	54 (16.1)	39 (41.5)	31 (16.9)		
	WHITE	266 (79.4)	51 (54.3)	138 (75.4)		
history of neoadjuvant treatment (%)	No	341 (100.0)	95 (100.0)	181 (96.8)	0.001	exact
	Yes	0 (0.0)	0 (0.0)	6 (3.2)		
age at initial pathologic diagnosis (median [IQR])		58.00 [48.00, 66.00]	53.00 [48.00, 65.00]	60.00 [50.00, 68.00]	0.165	nonnorm
ethnicity (%)	HISPANIC OR LATINO	17 (5.5)	4 (4.5)	6 (3.6)	0.699	exact
	NOT HISPANIC OR LATINO	292 (94.5)	84 (95.5)	161 (96.4)		
primary lymph node presentation assessment (%)	NO	11 (4.0)	3 (3.8)	12 (8.1)	0.201	exact

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	MOFA1	MOFA2	MOFA3	p	test
	YES	262 (96.0)	75 (96.2)	137 (91.9)		
histological type (%)	Infiltrating Carcinoma NOS	0 (0.0)	1 (1.1)	0 (0.0)	<0.001	exact
	Infiltrating Ductal Carcinoma	214 (62.6)	80 (84.2)	142 (75.9)		
	Infiltrating Lobular Carcinoma	89 (26.0)	2 (2.1)	27 (14.4)		
	Medullary Carcinoma	0 (0.0)	4 (4.2)	1 (0.5)		
	Metaplastic Carcinoma	3 (0.9)	2 (2.1)	2 (1.1)		
	Mixed Histology (please specify)	11 (3.2)	0 (0.0)	10 (5.3)		
	Mucinous Carcinoma	9 (2.6)	0 (0.0)	1 (0.5)		
	Other, specify	16 (4.7)	6 (6.3)	4 (2.1)		
menopause status (%)	Indeterminate (neither Pre or Postmenopausal)	10 (3.2)	2 (2.3)	2 (1.2)	0.204	exact
	Peri (6-12 months since last menstrual period)	10 (3.2)	6 (6.9)	4 (2.3)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	213 (67.4)	63 (72.4)	129 (75.4)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	83 (26.3)	16 (18.4)	36 (21.1)		
breast carcinoma progesterone receptor status (%)	Indeterminate	1 (0.3)	1 (1.1)	0 (0.0)	<0.001	exact
	Negative	60 (18.5)	84 (93.3)	40 (23.4)		
	Positive	264 (81.2)	5 (5.6)	131 (76.6)		
breast carcinoma estrogen receptor status (%)	Negative	28 (8.6)	78 (85.7)	24 (14.0)	<0.001	
	Positive	297 (91.4)	13 (14.3)	147 (86.0)		
lab proc her2 neu immuno-histochemistry receptor status (%)	Equivocal	59 (21.5)	17 (21.2)	34 (22.4)	<0.001	exact
	Indeterminate	4 (1.5)	2 (2.5)	4 (2.6)		
	Negative	183 (66.5)	58 (72.5)	75 (49.3)		
	Positive	29 (10.5)	3 (3.8)	39 (25.7)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	MOFA1	MOFA2	MOFA3	p	test
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.016	nonnorm
number of lymphnodes positive by her (median [IQR])		1.00 [0.00, 3.00]	0.00 [0.00, 1.00]	1.00 [0.00, 4.00]	0.001	nonnorm
er level cell percentage category (%)	<10%	10 (5.5)	16 (69.6)	9 (10.5)	<0.001	exact
	10-19%	10 (5.5)	1 (4.3)	9 (10.5)		
	20-29%	0 (0.0)	2 (8.7)	2 (2.3)		
	30-39%	2 (1.1)	1 (4.3)	2 (2.3)		
	40-49%	2 (1.1)	0 (0.0)	1 (1.2)		
	50-59%	7 (3.8)	0 (0.0)	1 (1.2)		
	60-69%	5 (2.7)	0 (0.0)	2 (2.3)		
	70-79%	27 (14.8)	1 (4.3)	12 (14.0)		
	80-89%	10 (5.5)	0 (0.0)	8 (9.3)		
	90-99%	110 (60.1)	2 (8.7)	40 (46.5)		
progesterone receptor level cell percent category (%)	<10%	37 (21.3)	19 (100.0)	23 (32.9)	<0.001	exact
	10-19%	11 (6.3)	0 (0.0)	7 (10.0)		
	20-29%	4 (2.3)	0 (0.0)	4 (5.7)		
	30-39%	6 (3.4)	0 (0.0)	2 (2.9)		
	40-49%	8 (4.6)	0 (0.0)	3 (4.3)		
	50-59%	9 (5.2)	0 (0.0)	1 (1.4)		
	60-69%	9 (5.2)	0 (0.0)	3 (4.3)		
	70-79%	23 (13.2)	0 (0.0)	6 (8.6)		
	80-89%	15 (8.6)	0 (0.0)	8 (11.4)		
	90-99%	52 (29.9)	0 (0.0)	13 (18.6)		
distant metastasis present ind2 (%)	NO	141 (96.6)	32 (97.0)	80 (100.0)	0.234	exact
	YES	5 (3.4)	1 (3.0)	0 (0.0)		
stage event pathologic stage (%)	Stage I	31 (9.1)	7 (7.4)	12 (6.5)	0.016	exact
	Stage IA	33 (9.7)	7 (7.4)	9 (4.8)		
	Stage IB	3 (0.9)	0 (0.0)	0 (0.0)		
	Stage II	3 (0.9)	1 (1.1)	1 (0.5)		
	Stage IIA	92 (27.1)	47 (49.5)	63 (33.9)		
	Stage IIB	89 (26.3)	20 (21.1)	39 (21.0)		
	Stage III	1 (0.3)	0 (0.0)	1 (0.5)		
	Stage IIIA	57 (16.8)	6 (6.3)	37 (19.9)		
	Stage IIIB	6 (1.8)	2 (2.1)	6 (3.2)		
	Stage IIIC	17 (5.0)	3 (3.2)	16 (8.6)		
	Stage IV	4 (1.2)	1 (1.1)	2 (1.1)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

level	MOFA1	MOFA2	MOFA3	p	test
Stage X	3 (0.9)	1 (1.1)	0 (0.0)		