

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	SNF1	SNF2	p	test
n		488	137		
Subtype (%)	SNF1	488 (100.0)	0 (0.0)	<0.001	
	SNF2	0 (0.0)	137 (100.0)		
vital status (%)	Alive	129 (93.5)	34 (94.4)	1.000	exact
	Dead	9 (6.5)	2 (5.6)		
days to birth (median [IQR])		-21514.00 [-24813.00, -17930.00]	-20227.00 [-23948.00, -17763.00]	0.163	nonnorm
days to death (median [IQR])		1590.50 [606.25, 2430.00]	939.50 [473.50, 2357.50]	0.527	nonnorm
days to last followup (median [IQR])		759.00 [477.00, 1611.00]	722.00 [450.00, 1572.00]	0.912	nonnorm
race list (%)	AMERICAN INDIAN OR ALASKA NATIVE	0 (0.0)	1 (0.7)	<0.001	exact
	ASIAN	24 (5.0)	8 (5.9)		
	BLACK OR AFRICAN AMERICAN	77 (16.1)	47 (34.8)		
	WHITE	376 (78.8)	79 (58.5)		
history of neoadjuvant treatment (%)	No	481 (98.8)	136 (100.0)	0.348	exact
	Yes	6 (1.2)	0 (0.0)		
age at initial pathologic diagnosis (median [IQR])		58.00 [48.00, 67.00]	55.00 [48.00, 65.00]	0.211	nonnorm
ethnicity (%)	HISPANIC OR LATINO	21 (4.8)	6 (4.8)	1.000	
	NOT HISPANIC OR LATINO	418 (95.2)	119 (95.2)		
primary lymph node presentation assessment (%)	NO	19 (4.9)	7 (6.3)	0.724	
	YES	370 (95.1)	104 (93.7)		
histological type (%)	Infiltrating Carcinoma NOS	0 (0.0)	1 (0.7)	<0.001	exact
	Infiltrating Ductal Carcinoma	323 (66.2)	113 (83.1)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	SNF1	SNF2	p	test
menopause status (%)	Infiltrating Lobular Carcinoma	116 (23.8)	2 (1.5)	0.064	exact
	Medullary Carcinoma	0 (0.0)	5 (3.7)		
	Metaplastic Carcinoma	1 (0.2)	6 (4.4)		
	Mixed Histology (please specify)	21 (4.3)	0 (0.0)		
	Mucinous Carcinoma	10 (2.0)	0 (0.0)		
	Other, specify	17 (3.5)	9 (6.6)		
	Indeterminate (neither Pre or Postmenopausal)	9 (2.0)	5 (4.0)		
	Peri (6-12 months since last menstrual period)	13 (2.9)	7 (5.6)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	313 (69.7)	92 (73.6)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	114 (25.4)	21 (16.8)		
breast carcinoma progesterone receptor status (%)	Indeterminate	1 (0.2)	1 (0.8)	<0.001	exact
breast carcinoma estrogen receptor status (%)	Negative	72 (15.7)	112 (88.2)	<0.001	
	Positive	386 (84.1)	14 (11.0)		
	Negative	27 (5.9)	103 (80.5)		
lab proc her2 neu immuno-histochemistry receptor status (%)	Positive	432 (94.1)	25 (19.5)	0.526	exact
	Equivocal	89 (22.4)	21 (19.3)		
	Indeterminate	7 (1.8)	3 (2.8)		
	Negative	243 (61.1)	73 (67.0)	0.005	nonnorm
	Positive	59 (14.8)	12 (11.0)		
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	SNF1	SNF2	p	test
number of lymphnodes positive by he (median [IQR])		1.00 [0.00, 3.00]	0.00 [0.00, 2.00]	0.004	nonnorm
er level cell percentage category (%)	<10%	11 (4.3)	24 (68.6)	<0.001	exact
	10-19%	19 (7.4)	1 (2.9)		
	20-29%	2 (0.8)	2 (5.7)		
	30-39%	3 (1.2)	2 (5.7)		
	40-49%	3 (1.2)	0 (0.0)		
	50-59%	8 (3.1)	0 (0.0)		
	60-69%	7 (2.7)	0 (0.0)		
	70-79%	39 (15.2)	1 (2.9)		
	80-89%	18 (7.0)	0 (0.0)		
	90-99%	147 (57.2)	5 (14.3)		
progesterone receptor level cell percent category (%)	<10%	51 (22.0)	28 (90.3)	<0.001	exact
	10-19%	18 (7.8)	0 (0.0)		
	20-29%	8 (3.4)	0 (0.0)		
	30-39%	8 (3.4)	0 (0.0)		
	40-49%	11 (4.7)	0 (0.0)		
	50-59%	10 (4.3)	0 (0.0)		
	60-69%	10 (4.3)	2 (6.5)		
	70-79%	29 (12.5)	0 (0.0)		
	80-89%	23 (9.9)	0 (0.0)		
	90-99%	64 (27.6)	1 (3.2)		
distant metastasis present ind2 (%)	NO	209 (97.7)	44 (97.8)	1.000	exact
	YES	5 (2.3)	1 (2.2)		
stage event pathologic stage (%)	Stage I	41 (8.5)	9 (6.6)	0.066	exact
	Stage IA	39 (8.1)	10 (7.4)		
	Stage IB	3 (0.6)	0 (0.0)		
	Stage II	4 (0.8)	1 (0.7)		
	Stage IIA	138 (28.5)	64 (47.1)		
	Stage IIB	122 (25.2)	26 (19.1)		
	Stage III	2 (0.4)	0 (0.0)		
	Stage IIIA	84 (17.4)	16 (11.8)		
	Stage IIIB	12 (2.5)	2 (1.5)		
	Stage IIIC	31 (6.4)	5 (3.7)		
	Stage IV	5 (1.0)	2 (1.5)		
	Stage X	3 (0.6)	1 (0.7)		