

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	wMKL1	wMKL2	wMKL3	wMKL4	wMKL5	wMKL6	wMKL7	wMKL8	p	test
n		45	115	18	96	77	143	45	86		
Subtype (%)	wMKL1	45 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	<0.001	exact
	wMKL2	0 (0.0)	115 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	wMKL3	0 (0.0)	0 (0.0)	18 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	wMKL4	0 (0.0)	0 (0.0)	0 (0.0)	96 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	wMKL5	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	77 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	wMKL6	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	143 (100.0)	0 (0.0)	0 (0.0)		
	wMKL7	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	45 (100.0)	0 (0.0)		
	wMKL8	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	86 (100.0)		
vital status (%)	Alive	11 (91.7)	35 (97.2)	6 (100.0)	22 (88.0)	20 (95.2)	35 (92.1)	9 (90.0)	25 (96.2)	0.817	exact
	Dead	1 (8.3)	1 (2.8)	0 (0.0)	3 (12.0)	1 (4.8)	3 (7.9)	1 (10.0)	1 (3.8)		
days to birth (median [IQR])		-20211.50 [-23870.25, -17610.00]	-20329.00 [-23144.00, -17341.25]	-23318.00 [-25833.00, -20288.00]	-21767.50 [-25387.25, -17656.75]	-19778.00 [-24075.00, -17763.00]	-22179.00 [-25119.00, -18985.00]	-23036.00 [-26004.50, -18754.25]	-21327.00 [-23397.00, -18177.00]	0.061	nonnorm
days to death (median [IQR])		NA [NA, NA]	571.00 [524.00, 860.00]	616.00 [616.00, 616.00]	2421.00 [567.25, 2523.50]	912.00 [573.00, 1759.00]	1910.50 [1221.75, 2517.50]	1286.00 [825.50, 2206.00]	1742.50 [1328.50, 1840.75]	0.534	nonnorm
days to last followup (median [IQR])		975.00 [416.00, 1779.00]	856.00 [476.00, 1808.00]	915.00 [618.50, 1705.50]	947.50 [535.25, 1928.50]	771.00 [480.00, 1604.00]	641.00 [409.00, 1298.00]	746.50 [461.25, 1413.75]	735.50 [495.00, 1215.75]	0.433	nonnorm
race list (%)	AMERICAN INDIAN OR ALASKA NATIVE	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.7)	0 (0.0)	0 (0.0)	<0.001	exact
	ASIAN	2 (4.5)	6 (5.4)	1 (5.6)	6 (6.5)	2 (2.6)	13 (9.3)	1 (2.3)	1 (1.2)		
	BLACK OR AFRICAN AMERICAN	5 (11.4)	24 (21.4)	3 (16.7)	21 (22.8)	31 (40.3)	29 (20.7)	3 (6.8)	8 (9.4)		
	WHITE	37 (84.1)	82 (73.2)	14 (77.8)	65 (70.7)	44 (57.1)	97 (69.3)	40 (90.9)	76 (89.4)		

Table 1: Summarization of clinical variables stratified by subtype (continued)

	level	wMKL1	wMKL2	wMKL3	wMKL4	wMKL5	wMKL6	wMKL7	wMKL8	p	test
history of neoadjuvant treatment (%)	No	45 (100.0)	113 (100.0)	18 (100.0)	96 (100.0)	77 (100.0)	138 (96.5)	44 (97.8)	86 (100.0)	0.065	exact
	Yes	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	5 (3.5)	1 (2.2)	0 (0.0)		
age at initial pathologic diagnosis (median [IQR])		55.00 [48.00, 65.00]	55.00 [47.00, 63.00]	62.00 [52.75, 70.00]	59.00 [47.75, 69.00]	54.00 [48.00, 65.00]	59.00 [50.00, 68.00]	63.00 [51.00, 71.00]	58.00 [49.25, 64.00]	0.068	nonnorm
ethnicity (%)	HISPANIC OR LATINO	3 (7.9)	5 (4.8)	3 (17.6)	4 (4.5)	4 (5.6)	6 (4.6)	0 (0.0)	2 (2.7)	0.248	exact
	NOT HISPANIC OR LATINO	35 (92.1)	100 (95.2)	14 (82.4)	85 (95.5)	67 (94.4)	125 (95.4)	38 (100.0)	73 (97.3)		
primary lymph node presentation assessment (%)	NO	1 (3.4)	5 (5.6)	1 (5.9)	1 (1.4)	3 (4.8)	10 (9.1)	3 (7.9)	2 (2.5)	0.335	exact
	YES	28 (96.6)	85 (94.4)	16 (94.1)	73 (98.6)	60 (95.2)	100 (90.9)	35 (92.1)	77 (97.5)		
histological type (%)	Infiltrating Carcinoma NOS	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.3)	0 (0.0)	0 (0.0)	0 (0.0)	<0.001	exact
	Infiltrating Ductal Carcinoma	25 (55.6)	93 (80.9)	8 (44.4)	62 (64.6)	62 (81.6)	120 (83.9)	25 (55.6)	41 (47.7)		
	Infiltrating Lobular Carcinoma	19 (42.2)	7 (6.1)	4 (22.2)	24 (25.0)	1 (1.3)	11 (7.7)	16 (35.6)	36 (41.9)		
	Medullary Carcinoma	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	4 (5.3)	1 (0.7)	0 (0.0)	0 (0.0)		
	Metaplastic Carcinoma	0 (0.0)	2 (1.7)	1 (5.6)	0 (0.0)	4 (5.3)	0 (0.0)	0 (0.0)	0 (0.0)		
	Mixed Histology (please specify)	0 (0.0)	1 (0.9)	1 (5.6)	6 (6.2)	0 (0.0)	6 (4.2)	4 (8.9)	3 (3.5)		
	Mucinous Carcinoma	0 (0.0)	3 (2.6)	2 (11.1)	1 (1.0)	0 (0.0)	1 (0.7)	0 (0.0)	3 (3.5)		
	Other, specify	1 (2.2)	9 (7.8)	2 (11.1)	3 (3.1)	4 (5.3)	4 (2.8)	0 (0.0)	3 (3.5)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	wMKL1	wMKL2	wMKL3	wMKL4	wMKL5	wMKL6	wMKL7	wMKL8	p	test
menopausal status (%)	Indeterminate (neither Pre or Post-menopausal)	2 (4.5)	4 (3.6)	0 (0.0)	3 (3.6)	2 (2.9)	1 (0.8)	1 (2.4)	1 (1.3)	0.194	exact
	Peri (6-12 months since last menstrual period)	5 (11.4)	3 (2.7)	0 (0.0)	2 (2.4)	6 (8.7)	3 (2.3)	0 (0.0)	1 (1.3)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	25 (56.8)	74 (66.1)	14 (82.4)	62 (73.8)	46 (66.7)	99 (76.7)	33 (78.6)	52 (67.5)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	12 (27.3)	31 (27.7)	3 (17.6)	17 (20.2)	15 (21.7)	26 (20.2)	8 (19.0)	23 (29.9)		
breast carcinoma progesterone receptor status (%)	Indeterminate	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.1)	1 (1.4)	0 (0.0)	0 (0.0)	0 (0.0)	<0.001	exact
	Negative	4 (8.9)	46 (41.8)	6 (40.0)	15 (16.3)	66 (93.0)	36 (26.7)	4 (10.5)	7 (8.8)		
	Positive	41 (91.1)	64 (58.2)	9 (60.0)	76 (82.6)	4 (5.6)	99 (73.3)	34 (89.5)	73 (91.2)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	wMKL1	wMKL2	wMKL3	wMKL4	wMKL5	wMKL6	wMKL7	wMKL8	p	test
breast carcinoma estrogen receptor status (%)	Negative	1 (2.2)	36 (32.4)	4 (26.7)	3 (3.3)	61 (85.9)	21 (15.6)	2 (5.3)	2 (2.5)	<0.001	exact
	Positive	44 (97.8)	75 (67.6)	11 (73.3)	89 (96.7)	10 (14.1)	114 (84.4)	36 (94.7)	78 (97.5)		
lab proc her2 neu immuno-histo-chemistry receptor status (%)	Equivocal	5 (13.2)	21 (21.4)	3 (27.3)	16 (20.3)	15 (23.8)	25 (21.0)	9 (28.1)	16 (23.9)	0.005	exact
	Indeterminate	0 (0.0)	2 (2.0)	0 (0.0)	2 (2.5)	2 (3.2)	4 (3.4)	0 (0.0)	0 (0.0)		
number of lymph nodes positive by ihc (median [IQR])	Negative	30 (78.9)	64 (65.3)	7 (63.6)	52 (65.8)	43 (68.3)	58 (48.7)	15 (46.9)	47 (70.1)		
	Positive	3 (7.9)	11 (11.2)	1 (9.1)	9 (11.4)	3 (4.8)	32 (26.9)	8 (25.0)	4 (6.0)	0.019	nonnorm
number of lymph nodes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 1.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.25]	0.00 [0.00, 0.50]		
number of lymph nodes positive by he (median [IQR])		1.00 [0.00, 2.75]	1.00 [0.00, 2.00]	1.00 [0.00, 5.00]	1.00 [0.00, 3.00]	0.00 [0.00, 1.25]	1.00 [0.00, 4.00]	1.00 [0.00, 3.00]	1.00 [0.00, 2.00]	0.044	nonnorm
er level cell percentage category (%)	<10%	1 (4.0)	7 (13.7)	2 (22.2)	3 (5.8)	13 (72.2)	8 (11.4)	1 (5.9)	0 (0.0)	<0.001	exact
	10-19%	0 (0.0)	5 (9.8)	1 (11.1)	2 (3.8)	1 (5.6)	8 (11.4)	1 (5.9)	2 (4.0)		
	20-29%	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	2 (11.1)	1 (1.4)	1 (5.9)	0 (0.0)		
	30-39%	0 (0.0)	0 (0.0)	1 (11.1)	1 (1.9)	1 (5.6)	1 (1.4)	1 (5.9)	0 (0.0)		
	40-49%	0 (0.0)	0 (0.0)	1 (11.1)	0 (0.0)	0 (0.0)	2 (2.9)	0 (0.0)	0 (0.0)		
	50-59%	1 (4.0)	0 (0.0)	0 (0.0)	3 (5.8)	0 (0.0)	1 (1.4)	0 (0.0)	3 (6.0)		
	60-69%	0 (0.0)	1 (2.0)	0 (0.0)	3 (5.8)	0 (0.0)	1 (1.4)	1 (5.9)	1 (2.0)		

Table 1: Summarization of clinical variables stratified by subtype (continued)

	level	wMKL1	wMKL2	wMKL3	wMKL4	wMKL5	wMKL6	wMKL7	wMKL8	p	test
	70-79%	4 (16.0)	8 (15.7)	1 (11.1)	6 (11.5)	1 (5.6)	10 (14.3)	1 (5.9)	9 (18.0)		
	80-89%	1 (4.0)	1 (2.0)	0 (0.0)	3 (5.8)	0 (0.0)	4 (5.7)	4 (23.5)	5 (10.0)		
	90-99%	18 (72.0)	29 (56.9)	3 (33.3)	31 (59.6)	0 (0.0)	34 (48.6)	7 (41.2)	30 (60.0)		
progesterone receptor level cell percent category (%)	<10%	2 (7.7)	12 (26.1)	3 (37.5)	17 (34.7)	15 (100.0)	23 (41.1)	1 (6.2)	6 (12.8)	<0.001	exact
	10-19%	1 (3.8)	4 (8.7)	0 (0.0)	3 (6.1)	0 (0.0)	4 (7.1)	3 (18.8)	3 (6.4)		
	20-29%	1 (3.8)	1 (2.2)	1 (12.5)	0 (0.0)	0 (0.0)	1 (1.8)	3 (18.8)	1 (2.1)		
	30-39%	1 (3.8)	2 (4.3)	1 (12.5)	1 (2.0)	0 (0.0)	3 (5.4)	0 (0.0)	0 (0.0)		
	40-49%	2 (7.7)	1 (2.2)	0 (0.0)	4 (8.2)	0 (0.0)	2 (3.6)	1 (6.2)	1 (2.1)		
	50-59%	2 (7.7)	1 (2.2)	0 (0.0)	2 (4.1)	0 (0.0)	2 (3.6)	0 (0.0)	3 (6.4)		
	60-69%	1 (3.8)	5 (10.9)	0 (0.0)	3 (6.1)	0 (0.0)	2 (3.6)	0 (0.0)	1 (2.1)		
	70-79%	3 (11.5)	8 (17.4)	1 (12.5)	4 (8.2)	0 (0.0)	5 (8.9)	1 (6.2)	7 (14.9)		
	80-89%	3 (11.5)	1 (2.2)	1 (12.5)	5 (10.2)	0 (0.0)	3 (5.4)	5 (31.2)	5 (10.6)		
	90-99%	10 (38.5)	11 (23.9)	1 (12.5)	10 (20.4)	0 (0.0)	11 (19.6)	2 (12.5)	20 (42.6)		
distant metastasis present ind2 (%)	NO	20 (100.0)	48 (96.0)	8 (80.0)	32 (97.0)	26 (100.0)	56 (100.0)	22 (100.0)	41 (97.6)	0.048	exact
	YES	0 (0.0)	2 (4.0)	2 (20.0)	1 (3.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (2.4)		
stage event pathologic stage (%)	Stage I	1 (2.2)	6 (5.3)	2 (11.1)	17 (17.9)	6 (7.9)	9 (6.3)	2 (4.5)	7 (8.2)	0.053	exact
	Stage IA	4 (8.9)	15 (13.2)	2 (11.1)	8 (8.4)	4 (5.3)	6 (4.2)	3 (6.8)	7 (8.2)		
	Stage IB	0 (0.0)	0 (0.0)	1 (5.6)	1 (1.1)	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.2)		
	Stage II	1 (2.2)	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.3)	1 (0.7)	0 (0.0)	2 (2.4)		
	Stage IIA	15 (33.3)	35 (30.7)	7 (38.9)	23 (24.2)	35 (46.1)	46 (32.2)	18 (40.9)	23 (27.1)		
	Stage IIB	11 (24.4)	28 (24.6)	2 (11.1)	22 (23.2)	17 (22.4)	32 (22.4)	7 (15.9)	29 (34.1)		
	Stage III	0 (0.0)	0 (0.0)	1 (5.6)	0 (0.0)	0 (0.0)	1 (0.7)	0 (0.0)	0 (0.0)		
	Stage IIIA	8 (17.8)	20 (17.5)	1 (5.6)	18 (18.9)	6 (7.9)	29 (20.3)	8 (18.2)	10 (11.8)		
	Stage IIIB	1 (2.2)	3 (2.6)	0 (0.0)	1 (1.1)	2 (2.6)	6 (4.2)	0 (0.0)	1 (1.2)		
	Stage IIIC	4 (8.9)	4 (3.5)	2 (11.1)	3 (3.2)	2 (2.6)	10 (7.0)	6 (13.6)	5 (5.9)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

level	wMKL1	wMKL2	wMKL3	wMKL4	wMKL5	wMKL6	wMKL7	wMKL8	p	test
Stage IV	0 (0.0)	2 (1.8)	0 (0.0)	1 (1.1)	2 (2.6)	2 (1.4)	0 (0.0)	0 (0.0)		
Stage X	0 (0.0)	1 (0.9)	0 (0.0)	1 (1.1)	1 (1.3)	1 (0.7)	0 (0.0)	0 (0.0)		