

Summary of ordinal clinical variables

Contents

Table 1: Summary of ordinal clinical variables

Variable	Levels	NEMO1	NEMO2	p	test
number of lymphnodes positive by ihc (%)	0	114 (78.6)	36 (97.3)	0.0046	JT
	1	25 (17.2)	1 (2.7)		
	2	3 (2.1)	0 (0.0)		
	3	2 (1.4)	0 (0.0)		
	4	1 (0.7)	0 (0.0)		
number of lymphnodes positive by he (%)	0	184 (41.0)	68 (58.1)	0.001	JT
	1	90 (20.0)	15 (12.8)		
	2	43 (9.6)	14 (12.0)		
	3	30 (6.7)	4 (3.4)		
	4	22 (4.9)	1 (0.9)		
	5	17 (3.8)	3 (2.6)		
	6	10 (2.2)	5 (4.3)		
	7	5 (1.1)	1 (0.9)		
	8	7 (1.6)	0 (0.0)		
	9	5 (1.1)	1 (0.9)		
	10	3 (0.7)	3 (2.6)		
	11	6 (1.3)	1 (0.9)		
	12	6 (1.3)	0 (0.0)		
	13	2 (0.4)	0 (0.0)		
	14	4 (0.9)	1 (0.9)		
	17	1 (0.2)	0 (0.0)		
	19	2 (0.4)	0 (0.0)		
	20	1 (0.2)	0 (0.0)		
	21	2 (0.4)	0 (0.0)		
	22	1 (0.2)	0 (0.0)		
	23	1 (0.2)	0 (0.0)		
	24	1 (0.2)	0 (0.0)		
	27	2 (0.4)	0 (0.0)		
	28	1 (0.2)	0 (0.0)		
	29	2 (0.4)	0 (0.0)		
	35	1 (0.2)	0 (0.0)		