

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	NEMO1	NEMO2	p	test
n		497	128		
Subtype (%)	NEMO1	497 (100.0)	0 (0.0)	<0.001	
	NEMO2	0 (0.0)	128 (100.0)		
vital status (%)		10 (6.7)	1 (2.9)	0.908	exact
	Alive	131 (87.3)	32 (91.4)		
	Dead	9 (6.0)	2 (5.7)		
days to birth (median [IQR])		-21557.00 [-24895.50, -18006.75]	-20066.00 [-23932.25, -17679.75]	0.059	nonnorm
days to death (median [IQR])		1590.50 [642.75, 2456.00]	848.50 [372.50, 2083.75]	0.277	nonnorm
days to last followup (median [IQR])		755.00 [477.00, 1611.00]	738.00 [445.50, 1621.50]	0.977	nonnorm
race list (%)		11 (2.2)	2 (1.6)	<0.001	exact
	AMERICAN INDIAN OR ALASKA NATIVE	0 (0.0)	1 (0.8)		
	ASIAN	26 (5.2)	6 (4.7)		
	BLACK OR AFRICAN AMERICAN	80 (16.1)	44 (34.4)		
	WHITE	380 (76.5)	75 (58.6)		
history of neoadjuvant treatment (%)		1 (0.2)	1 (0.8)	0.291	exact
	No	490 (98.6)	127 (99.2)		
	Yes	6 (1.2)	0 (0.0)	0.082	nonnorm
age at initial pathologic diagnosis (median [IQR])		58.00 [49.00, 67.00]	54.00 [48.00, 65.00]		
ethnicity (%)		50 (10.1)	11 (8.6)	0.866	
	HISPANIC OR LATINO	21 (4.2)	6 (4.7)		
	NOT HISPANIC OR LATINO	426 (85.7)	111 (86.7)		
primary lymph node presentation assessment (%)		100 (20.1)	25 (19.5)	0.974	
	NO	21 (4.2)	5 (3.9)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	NEMO1	NEMO2	p	test
histological type (%)	YES	376 (75.7)	98 (76.6)	<0.001	exact
		0 (0.0)	1 (0.8)		
	Infiltrating Carcinoma NOS	0 (0.0)	1 (0.8)		
	Infiltrating Ductal Carcinoma	331 (66.6)	105 (82.0)		
	Infiltrating Lobular Carcinoma	116 (23.3)	2 (1.6)		
	Medullary Carcinoma	1 (0.2)	4 (3.1)		
	Metaplastic Carcinoma	1 (0.2)	6 (4.7)		
	Mixed Histology (please specify)	21 (4.2)	0 (0.0)		
	Mucinous Carcinoma	10 (2.0)	0 (0.0)		
menopause status (%)	Other, specify	17 (3.4)	9 (7.0)	0.090	exact
		39 (7.8)	12 (9.4)		
	Indeterminate (neither Pre or Postmenopausal)	9 (1.8)	5 (3.9)		
	Peri (6-12 months since last menstrual period)	13 (2.6)	7 (5.5)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	321 (64.6)	84 (65.6)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	115 (23.1)	20 (15.6)		
		32 (6.4)	7 (5.5)		
breast carcinoma progesterone receptor status (%)	Indeterminate	1 (0.2)	1 (0.8)	<0.001	exact
	Negative	74 (14.9)	110 (85.9)		
	Positive	390 (78.5)	10 (7.8)		
breast carcinoma estrogen receptor status (%)		32 (6.4)	6 (4.7)	<0.001	
	Negative	28 (5.6)	102 (79.7)		
	Positive	437 (87.9)	20 (15.6)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	NEMO1	NEMO2	p	test
lab proc her2 neu immuno- histochemistry receptor status (%)		95 (19.1)	23 (18.0)	0.476	exact
	Equivocal	89 (17.9)	21 (16.4)		
	Indeterminate	7 (1.4)	3 (2.3)		
	Negative	245 (49.3)	71 (55.5)		
	Positive	61 (12.3)	10 (7.8)		
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.008	nonnorm
number of lymphnodes positive by he (median [IQR])		1.00 [0.00, 3.00]	0.00 [0.00, 2.00]	0.001	nonnorm
er level cell percentage category (%)		237 (47.7)	96 (75.0)	<0.001	exact
	<10%	11 (2.2)	24 (18.8)		
	10-19%	19 (3.8)	1 (0.8)		
	20-29%	2 (0.4)	2 (1.6)		
	30-39%	4 (0.8)	1 (0.8)		
	40-49%	3 (0.6)	0 (0.0)		
	50-59%	8 (1.6)	0 (0.0)		
	60-69%	7 (1.4)	0 (0.0)		
	70-79%	39 (7.8)	1 (0.8)		
	80-89%	18 (3.6)	0 (0.0)		
	90-99%	149 (30.0)	3 (2.3)		
progesterone receptor level cell percent category (%)		262 (52.7)	100 (78.1)	<0.001	exact
	<10%	52 (10.5)	27 (21.1)		
	10-19%	18 (3.6)	0 (0.0)		
	20-29%	8 (1.6)	0 (0.0)		
	30-39%	8 (1.6)	0 (0.0)		
	40-49%	11 (2.2)	0 (0.0)		
	50-59%	10 (2.0)	0 (0.0)		
	60-69%	11 (2.2)	1 (0.8)		
	70-79%	29 (5.8)	0 (0.0)		
	80-89%	23 (4.6)	0 (0.0)		
	90-99%	65 (13.1)	0 (0.0)		
distant metastasis present ind2 (%)		281 (56.5)	85 (66.4)	0.117	exact
	NO	211 (42.5)	42 (32.8)		
	YES	5 (1.0)	1 (0.8)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	NEMO1	NEMO2	p	test
stage event pathologic stage (%)		4 (0.8)	1 (0.8)	0.033	exact
	Stage I	42 (8.5)	8 (6.2)		
	Stage IA	40 (8.0)	9 (7.0)		
	Stage IB	3 (0.6)	0 (0.0)		
	Stage II	4 (0.8)	1 (0.8)		
	Stage IIA	140 (28.2)	62 (48.4)		
	Stage IIB	124 (24.9)	24 (18.8)		
	Stage III	2 (0.4)	0 (0.0)		
	Stage IIIA	86 (17.3)	14 (10.9)		
	Stage IIIB	12 (2.4)	2 (1.6)		
	Stage IIIC	32 (6.4)	4 (3.1)		
	Stage IV	5 (1.0)	2 (1.6)		
	Stage X	3 (0.6)	1 (0.8)		