

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	ab-SNF1	ab-SNF2	p	test
n		506	119		
Subtype (%)	ab-SNF1	506 (100.0)	0 (0.0)	<0.001	
	ab-SNF2	0 (0.0)	119 (100.0)		
vital status (%)	Alive	134 (93.7)	29 (93.5)	1.000	exact
	Dead	9 (6.3)	2 (6.5)		
days to birth (median [IQR])		-21514.00 [-24808.00, -17949.50]	-20052.00 [-24000.50, -17726.00]	0.114	nonnorm
days to death (median [IQR])		1673.00 [669.50, 2395.00]	785.00 [322.00, 2854.00]	0.368	nonnorm
days to last followup (median [IQR])		760.50 [480.50, 1611.00]	718.00 [436.75, 1582.00]	0.651	nonnorm
race list (%)	AMERICAN INDIAN OR ALASKA NATIVE	0 (0.0)	1 (0.9)	<0.001	exact
	ASIAN	26 (5.3)	6 (5.1)		
	BLACK OR AFRICAN AMERICAN	79 (16.0)	45 (38.5)		
	WHITE	390 (78.8)	65 (55.6)		
history of neoadjuvant treatment (%)	No	499 (98.8)	118 (100.0)	0.601	exact
	Yes	6 (1.2)	0 (0.0)		
age at initial pathologic diagnosis (median [IQR])		58.00 [48.25, 67.00]	54.00 [48.00, 65.00]	0.151	nonnorm
ethnicity (%)	HISPANIC OR LATINO	22 (4.8)	5 (4.5)	1.000	
	NOT HISPANIC OR LATINO	432 (95.2)	105 (95.5)		
primary lymph node presentation assessment (%)	NO	21 (5.2)	5 (5.2)	1.000	exact
	YES	383 (94.8)	91 (94.8)		
histological type (%)	Infiltrating Carcinoma NOS	0 (0.0)	1 (0.8)	<0.001	exact
	Infiltrating Ductal Carcinoma	339 (67.0)	97 (82.2)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	ab-SNF1	ab-SNF2	p	test
menopause status (%)	Infiltrating Lobular Carcinoma	116 (22.9)	2 (1.7)	0.031	exact
	Medullary Carcinoma	0 (0.0)	5 (4.2)		
	Metaplastic Carcinoma	1 (0.2)	6 (5.1)		
	Mixed Histology (please specify)	21 (4.2)	0 (0.0)		
	Mucinous Carcinoma	10 (2.0)	0 (0.0)		
	Other, specify	19 (3.8)	7 (5.9)		
	Indeterminate (neither Pre or Postmenopausal)	9 (1.9)	5 (4.6)		
	Peri (6-12 months since last menstrual period)	13 (2.8)	7 (6.4)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	326 (70.1)	79 (72.5)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	117 (25.2)	18 (16.5)		
breast carcinoma progesterone receptor status (%)	Indeterminate	1 (0.2)	1 (0.9)	<0.001	exact
breast carcinoma estrogen receptor status (%)	Negative	80 (16.9)	104 (92.0)	<0.001	
	Positive	392 (82.9)	8 (7.1)		
	Negative	31 (6.6)	99 (86.8)		
lab proc her2 neu immuno-histochemistry receptor status (%)	Positive	442 (93.4)	15 (13.2)	0.156	exact
	Equivocal	91 (22.2)	19 (19.6)		
	Indeterminate	7 (1.7)	3 (3.1)		
	Negative	249 (60.7)	67 (69.1)		
	Positive	63 (15.4)	8 (8.2)		
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.003	nonnorm

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	ab-SNF1	ab-SNF2	p	test
number of lymphnodes positive by he (median [IQR])		1.00 [0.00, 3.00]	0.00 [0.00, 2.00]	0.001	nonnorm
er level cell percentage category (%)	<10%	13 (4.9)	22 (78.6)	<0.001	exact
	10-19%	19 (7.2)	1 (3.6)		
	20-29%	2 (0.8)	2 (7.1)		
	30-39%	4 (1.5)	1 (3.6)		
	40-49%	3 (1.1)	0 (0.0)		
	50-59%	8 (3.0)	0 (0.0)		
	60-69%	7 (2.7)	0 (0.0)		
	70-79%	40 (15.2)	0 (0.0)		
	80-89%	18 (6.8)	0 (0.0)		
	90-99%	150 (56.8)	2 (7.1)		
progesterone receptor level cell percent category (%)	<10%	54 (22.7)	25 (100.0)	<0.001	exact
	10-19%	18 (7.6)	0 (0.0)		
	20-29%	8 (3.4)	0 (0.0)		
	30-39%	8 (3.4)	0 (0.0)		
	40-49%	11 (4.6)	0 (0.0)		
	50-59%	10 (4.2)	0 (0.0)		
	60-69%	12 (5.0)	0 (0.0)		
	70-79%	29 (12.2)	0 (0.0)		
	80-89%	23 (9.7)	0 (0.0)		
	90-99%	65 (27.3)	0 (0.0)		
distant metastasis present ind2 (%)	NO	216 (97.7)	37 (97.4)	1.000	exact
	YES	5 (2.3)	1 (2.6)		
stage event pathologic stage (%)	Stage I	43 (8.6)	7 (5.9)	0.044	exact
	Stage IA	40 (8.0)	9 (7.6)		
	Stage IB	3 (0.6)	0 (0.0)		
	Stage II	4 (0.8)	1 (0.8)		
	Stage IIA	145 (28.9)	57 (48.3)		
	Stage IIB	125 (24.9)	23 (19.5)		
	Stage III	2 (0.4)	0 (0.0)		
	Stage IIIA	88 (17.5)	12 (10.2)		
	Stage IIIB	12 (2.4)	2 (1.7)		
	Stage IIIC	32 (6.4)	4 (3.4)		
	Stage IV	5 (1.0)	2 (1.7)		
	Stage X	3 (0.6)	1 (0.8)		