

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	CIMLR1	CIMLR2	p	test
n		326	299		
Subtype (%)	CIMLR1	326 (100.0)	0 (0.0)	<0.001	
	CIMLR2	0 (0.0)	299 (100.0)		
vital status (%)	Alive	83 (94.3)	80 (93.0)	0.969	
	Dead	5 (5.7)	6 (7.0)		
days to birth (median [IQR])		-21056.00 [-24294.50, -17563.50]	-21391.00 [-24861.00, -18156.50]	0.218	nonnorm
days to death (median [IQR])		1673.00 [720.50, 2446.50]	959.00 [538.00, 2417.00]	0.346	nonnorm
days to last followup (median [IQR])		776.00 [499.00, 1561.50]	726.00 [448.50, 1681.50]	0.727	nonnorm
race list (%)	AMERICAN INDIAN OR ALASKA NATIVE	0 (0.0)	1 (0.3)	0.200	exact
	ASIAN	16 (5.0)	16 (5.4)		
	BLACK OR AFRICAN AMERICAN	56 (17.7)	68 (23.1)		
	WHITE	245 (77.3)	210 (71.2)		
history of neoadjuvant treatment (%)	No	321 (98.8)	296 (99.3)	0.688	exact
	Yes	4 (1.2)	2 (0.7)		
age at initial pathologic diagnosis (median [IQR])		57.00 [48.00, 66.00]	58.00 [49.00, 67.00]	0.297	nonnorm
ethnicity (%)	HISPANIC OR LATINO	10 (3.4)	17 (6.2)	0.170	
	NOT HISPANIC OR LATINO	282 (96.6)	255 (93.8)		
primary lymph node presentation assessment (%)	NO	12 (4.6)	14 (5.9)	0.635	
	YES	251 (95.4)	223 (94.1)		
histological type (%)	Infiltrating Carcinoma NOS	0 (0.0)	1 (0.3)	<0.001	exact
	Infiltrating Ductal Carcinoma	211 (64.9)	225 (75.3)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	CIMLR1	CIMLR2	p	test
menopause status (%)	Infiltrating Lobular Carcinoma	77 (23.7)	41 (13.7)	0.244	
	Medullary Carcinoma	2 (0.6)	3 (1.0)		
	Metaplastic Carcinoma	2 (0.6)	5 (1.7)		
	Mixed Histology (please specify)	16 (4.9)	5 (1.7)		
	Mucinous Carcinoma	1 (0.3)	9 (3.0)		
	Other, specify	16 (4.9)	10 (3.3)		
	Indeterminate (neither Pre or Postmenopausal)	8 (2.7)	6 (2.2)		
	Peri (6-12 months since last menstrual period)	8 (2.7)	12 (4.3)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	203 (68.1)	202 (73.2)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	79 (26.5)	56 (20.3)		
breast carcinoma progesterone receptor status (%)	Indeterminate	1 (0.3)	1 (0.4)	<0.001	exact
breast carcinoma estrogen receptor status (%)	Negative	61 (19.9)	123 (43.9)	<0.001	
	Positive	244 (79.7)	156 (55.7)		
	Negative	42 (13.7)	88 (31.4)		
lab proc her2 neu immuno-histochemistry receptor status (%)	Positive	265 (86.3)	192 (68.6)	0.164	exact
	Equivocal	56 (21.4)	54 (22.0)		
	Indeterminate	4 (1.5)	6 (2.4)		
	Negative	157 (59.9)	159 (64.9)		
	Positive	45 (17.2)	26 (10.6)		
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.951	nonnorm

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	CIMLR1	CIMLR2	p	test
number of lymphnodes positive by he (median [IQR])		1.00 [0.00, 3.00]	1.00 [0.00, 2.00]	0.233	nonnorm
er level cell percentage category (%)	<10%	14 (9.2)	21 (15.1)	0.472	exact
	10-19%	8 (5.2)	12 (8.6)		
	20-29%	1 (0.7)	3 (2.2)		
	30-39%	2 (1.3)	3 (2.2)		
	40-49%	2 (1.3)	1 (0.7)		
	50-59%	5 (3.3)	3 (2.2)		
	60-69%	2 (1.3)	5 (3.6)		
	70-79%	23 (15.0)	17 (12.2)		
	80-89%	11 (7.2)	7 (5.0)		
	90-99%	85 (55.6)	67 (48.2)		
progesterone receptor level cell percent category (%)	<10%	31 (21.7)	48 (40.0)	0.031	exact
	10-19%	12 (8.4)	6 (5.0)		
	20-29%	7 (4.9)	1 (0.8)		
	30-39%	4 (2.8)	4 (3.3)		
	40-49%	6 (4.2)	5 (4.2)		
	50-59%	4 (2.8)	6 (5.0)		
	60-69%	6 (4.2)	6 (5.0)		
	70-79%	16 (11.2)	13 (10.8)		
	80-89%	17 (11.9)	6 (5.0)		
	90-99%	40 (28.0)	25 (20.8)		
distant metastasis present ind2 (%)	NO	140 (97.9)	113 (97.4)	1.000	exact
	YES	3 (2.1)	3 (2.6)		
stage event pathologic stage (%)	Stage I	27 (8.4)	23 (7.7)	0.146	exact
	Stage IA	23 (7.1)	26 (8.7)		
	Stage IB	2 (0.6)	1 (0.3)		
	Stage II	4 (1.2)	1 (0.3)		
	Stage IIA	99 (30.7)	103 (34.6)		
	Stage IIB	81 (25.2)	67 (22.5)		
	Stage III	2 (0.6)	0 (0.0)		
	Stage IIIA	55 (17.1)	45 (15.1)		
	Stage IIIB	2 (0.6)	12 (4.0)		
	Stage IIIC	19 (5.9)	17 (5.7)		
	Stage IV	5 (1.6)	2 (0.7)		
	Stage X	3 (0.9)	1 (0.3)		