

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	LRAcluster1	LRAcluster2	LRAcluster3	LRAcluster4	p	test
n		63	310	73	179		
Subtype (%)	LRAcluster1	63 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	<0.001	
	LRAcluster2	0 (0.0)	310 (100.0)	0 (0.0)	0 (0.0)		
	LRAcluster3	0 (0.0)	0 (0.0)	73 (100.0)	0 (0.0)		
	LRAcluster4	0 (0.0)	0 (0.0)	0 (0.0)	179 (100.0)		
vital status (%)	Alive	17 (94.4)	87 (94.6)	17 (94.4)	42 (91.3)	0.919	exact
	Dead	1 (5.6)	5 (5.4)	1 (5.6)	4 (8.7)		
days to birth (median [IQR])		-20210.00 [-23681.00, -17713.00]	-21291.00 [-24324.00, -17648.50]	-20080.00 [-24075.00, -18214.00]	-22016.50 [-25200.00, -18345.75]	0.433	nonnorm
days to death (median [IQR])		570.00 [372.50, 742.75]	1508.00 [577.00, 2373.00]	1759.00 [967.00, 2965.00]	1541.00 [521.00, 2450.50]	0.085	nonnorm
days to last followup (median [IQR])		738.00 [437.50, 1676.00]	862.50 [505.25, 1691.00]	882.50 [456.75, 2080.25]	647.00 [427.00, 1226.50]	0.084	nonnorm
race list (%)	AMERICAN INDIAN OR ALASKA NATIVE	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.6)	0.001	exact
	ASIAN	4 (6.6)	12 (4.0)	3 (4.1)	13 (7.4)		
	BLACK OR AFRICAN AMERICAN	18 (29.5)	47 (15.5)	27 (37.0)	32 (18.3)		
	WHITE	39 (63.9)	244 (80.5)	43 (58.9)	129 (73.7)		
history of neoadjuvant treatment (%)	No	62 (100.0)	309 (100.0)	73 (100.0)	173 (96.6)	0.002	exact
	Yes	0 (0.0)	0 (0.0)	0 (0.0)	6 (3.4)		
age at initial pathologic diagnosis (median [IQR])		55.00 [48.00, 64.50]	58.00 [48.00, 66.00]	54.00 [49.00, 65.00]	59.00 [49.50, 68.00]	0.538	nonnorm
ethnicity (%)	HISPANIC OR LATINO	2 (3.4)	15 (5.4)	4 (6.1)	6 (3.8)	0.820	exact
	NOT HISPANIC OR LATINO	57 (96.6)	264 (94.6)	62 (93.9)	154 (96.2)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	LRACluster1	LRACluster2	LRACluster3	LRACluster4	p	test
primary lymph node presentation assessment (%)	NO	3 (5.7)	8 (3.3)	3 (5.0)	12 (8.5)	0.165	exact
	YES	50 (94.3)	237 (96.7)	57 (95.0)	130 (91.5)		
histological type (%)	Infiltrating Carcinoma NOS	0 (0.0)	0 (0.0)	1 (1.4)	0 (0.0)	<0.001	exact
	Infiltrating Ductal Carcinoma	45 (71.4)	196 (63.2)	58 (80.6)	137 (76.5)		
	Infiltrating Lobular Carcinoma	9 (14.3)	81 (26.1)	2 (2.8)	26 (14.5)		
	Medullary Carcinoma	0 (0.0)	0 (0.0)	5 (6.9)	0 (0.0)		
	Metaplastic Carcinoma	3 (4.8)	0 (0.0)	3 (4.2)	1 (0.6)		
	Mixed Histology (please specify)	0 (0.0)	11 (3.5)	0 (0.0)	10 (5.6)		
	Mucinous Carcinoma	0 (0.0)	9 (2.9)	0 (0.0)	1 (0.6)		
	Other, specify	6 (9.5)	13 (4.2)	3 (4.2)	4 (2.2)		
menopause status (%)	Indeterminate (neither Pre or Postmenopausal)	2 (3.3)	8 (2.8)	2 (3.1)	2 (1.2)	0.098	exact
	Peri (6-12 months since last menstrual period)	2 (3.3)	9 (3.2)	6 (9.2)	3 (1.8)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	47 (77.0)	190 (66.7)	47 (72.3)	121 (74.2)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	10 (16.4)	78 (27.4)	10 (15.4)	37 (22.7)		
breast carcinoma progesterone receptor status (%)	Indeterminate	1 (1.7)	1 (0.3)	0 (0.0)	0 (0.0)	<0.001	exact
	Negative	45 (77.6)	42 (14.1)	60 (90.9)	37 (22.4)		
	Positive	12 (20.7)	254 (85.5)	6 (9.1)	128 (77.6)		
breast carcinoma estrogen receptor status (%)	Negative	43 (72.9)	11 (3.7)	54 (81.8)	22 (13.3)	<0.001	
	Positive	16 (27.1)	286 (96.3)	12 (18.2)	143 (86.7)		
lab proc her2 neu immuno-histochemistry receptor status (%)	Equivocal	9 (18.0)	53 (20.9)	15 (26.3)	33 (22.4)	<0.001	exact

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	LRAcluster1	LRAcluster2	LRAcluster3	LRAcluster4	p	test
number of lymphnodes positive by ihc (median [IQR])	Indeterminate	0 (0.0)	4 (1.6)	2 (3.5)	4 (2.7)	0.117	nonnorm
	Negative	35 (70.0)	172 (68.0)	38 (66.7)	71 (48.3)		
	Positive	6 (12.0)	24 (9.5)	2 (3.5)	39 (26.5)		
		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	<0.001	nonnorm
	number of lymphnodes positive by he (median [IQR])	0.00 [0.00, 2.00]	1.00 [0.00, 3.00]	0.00 [0.00, 1.00]	1.00 [0.00, 4.00]		
	er level cell percentage category (%)	<10%	10 (45.5)	6 (3.5)	14 (73.7)	<0.001	exact
		10-19%	2 (9.1)	8 (4.7)	1 (5.3)	<0.001	exact
		20-29%	0 (0.0)	0 (0.0)	2 (10.5)		
		30-39%	0 (0.0)	2 (1.2)	1 (5.3)		
		40-49%	0 (0.0)	2 (1.2)	0 (0.0)		
		50-59%	0 (0.0)	7 (4.1)	0 (0.0)		
		60-69%	1 (4.5)	4 (2.4)	0 (0.0)		
		70-79%	4 (18.2)	24 (14.1)	0 (0.0)		
		80-89%	0 (0.0)	10 (5.9)	0 (0.0)		
		90-99%	5 (22.7)	107 (62.9)	1 (5.3)		
	progesterone receptor level cell percent category (%)	<10%	12 (63.2)	31 (19.3)	16 (94.1)		
distant metastasis present ind2 (%)		10-19%	0 (0.0)	11 (6.8)	0 (0.0)	0.018	exact
		20-29%	1 (5.3)	3 (1.9)	0 (0.0)		
		30-39%	0 (0.0)	6 (3.7)	0 (0.0)		
		40-49%	0 (0.0)	8 (5.0)	0 (0.0)		
		50-59%	0 (0.0)	9 (5.6)	0 (0.0)		
		60-69%	0 (0.0)	9 (5.6)	1 (5.9)		
		70-79%	3 (15.8)	20 (12.4)	0 (0.0)		
		80-89%	1 (5.3)	14 (8.7)	0 (0.0)		
		90-99%	2 (10.5)	50 (31.1)	0 (0.0)		
	NO	18 (85.7)	133 (97.8)	26 (100.0)	76 (100.0)		
stage event pathologic stage (%)	YES	3 (14.3)	3 (2.2)	0 (0.0)	0 (0.0)	0.015	exact
	Stage I	5 (7.9)	29 (9.4)	7 (9.7)	9 (5.1)		
		Stage IA	5 (7.9)	31 (10.1)	4 (5.6)		
		Stage IB	0 (0.0)	3 (1.0)	0 (0.0)		
		Stage II	0 (0.0)	3 (1.0)	1 (1.4)		
		Stage IIA	28 (44.4)	81 (26.4)	35 (48.6)		
		Stage IIB	14 (22.2)	81 (26.4)	14 (19.4)		
		Stage III	1 (1.6)	0 (0.0)	0 (0.0)		
		Stage IIIA	6 (9.5)	52 (16.9)	6 (8.3)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

level	LRAcluster1	LRAcluster2	LRAcluster3	LRAcluster4	p	test
Stage IIIB	0 (0.0)	6 (2.0)	2 (2.8)	6 (3.4)		
Stage IIIC	3 (4.8)	15 (4.9)	1 (1.4)	17 (9.6)		
Stage IV	1 (1.6)	3 (1.0)	1 (1.4)	2 (1.1)		
Stage X	0 (0.0)	3 (1.0)	1 (1.4)	0 (0.0)		