

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	MFAMFA1	MFAMFA2	MFAMFA3	p
n		269	228	128	
Subtype (%)	MFA1	269 (100.0)	0 (0.0)	0 (0.0)	<0.001
	MFA2	0 (0.0)	228 (100.0)	0 (0.0)	
	MFA3	0 (0.0)	0 (0.0)	128 (100.0)	
vital status (%)	Alive	68 (93.2)	66 (94.3)	29 (93.5)	1.000
	Dead	5 (6.8)	4 (5.7)	2 (6.5)	
days to birth (median [IQR])		-20694.00 [-23928.00, -17529.00]	-22277.50 [-25301.00, -18948.25]	-20080.00 [-24000.50, -17768.50]	0.011
days to death (median [IQR])		1726.50 [604.75, 2334.75]	1590.50 [748.50, 2406.00]	848.50 [372.50, 2562.50]	0.648
days to last followup (median [IQR])		847.00 [482.50, 1737.50]	703.00 [477.00, 1467.00]	730.00 [443.25, 1565.50]	0.469
race list (%)	AMERICAN INDIAN OR ALASKA NATIVE	0 (0.0)	0 (0.0)	1 (0.8)	<0.001
	ASIAN	10 (3.8)	14 (6.3)	8 (6.3)	
	BLACK OR AFRICAN AMERICAN	40 (15.2)	38 (17.0)	46 (36.5)	
	WHITE	213 (81.0)	171 (76.7)	71 (56.3)	
history of neoadjuvant treatment (%)	No	266 (98.9)	224 (98.7)	127 (100.0)	0.670
	Yes	3 (1.1)	3 (1.3)	0 (0.0)	
age at initial pathologic diagnosis (median [IQR])		56.00 [47.00, 65.00]	60.00 [50.75, 69.00]	54.00 [48.00, 65.00]	0.024
ethnicity (%)	HISPANIC OR LATINO	10 (4.2)	10 (4.8)	7 (5.9)	0.776
	NOT HISPANIC OR LATINO	227 (95.8)	199 (95.2)	111 (94.1)	
primary lymph node presentation assessment (%)	NO	6 (2.7)	13 (7.6)	7 (6.7)	0.065
	YES	219 (97.3)	157 (92.4)	98 (93.3)	
histological type (%)	Infiltrating Carcinoma NOS	0 (0.0)	0 (0.0)	1 (0.8)	<0.001

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	MFAMFA1	MFAMFA2	MFAMFA3	p
	Infiltrating Ductal Carcinoma	148 (55.0)	182 (79.8)	106 (83.5)	
	Infiltrating Lobular Carcinoma	96 (35.7)	20 (8.8)	2 (1.6)	
	Medullary Carcinoma	0 (0.0)	0 (0.0)	5 (3.9)	
	Metaplastic Carcinoma	2 (0.7)	0 (0.0)	5 (3.9)	
	Mixed Histology (please specify)	11 (4.1)	10 (4.4)	0 (0.0)	
	Mucinous Carcinoma	4 (1.5)	6 (2.6)	0 (0.0)	
	Other, specify	8 (3.0)	10 (4.4)	8 (6.3)	
menopause status (%)	Indeterminate (neither Pre or Postmenopausal)	3 (1.2)	6 (2.8)	5 (4.2)	0.042
	Peri (6-12 months since last menstrual period)	8 (3.3)	5 (2.4)	7 (5.9)	
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	163 (66.5)	158 (74.9)	84 (71.2)	
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	71 (29.0)	42 (19.9)	22 (18.6)	
breast carcinoma progesterone receptor status (%)	Indeterminate	1 (0.4)	0 (0.0)	1 (0.8)	<0.001
	Negative	42 (16.3)	35 (16.7)	107 (89.2)	
	Positive	214 (83.3)	174 (83.3)	12 (10.0)	
breast carcinoma estrogen receptor status (%)	Negative	16 (6.2)	12 (5.7)	102 (84.3)	<0.001
	Positive	241 (93.8)	197 (94.3)	19 (15.7)	
lab proc her2 neu immuno-histochemistry receptor status (%)	Equivocal	43 (19.1)	45 (25.3)	22 (21.2)	0.004
	Indeterminate	2 (0.9)	6 (3.4)	2 (1.9)	
	Negative	154 (68.4)	91 (51.1)	71 (68.3)	
	Positive	26 (11.6)	36 (20.2)	9 (8.7)	
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 0.50]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.017

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	MFAMFA1	MFAMFA2	MFAMFA3	p
number of lymphnodes positive by he (median [IQR])		1.00 [0.00, 3.00]	1.00 [0.00, 4.00]	0.00 [0.00, 2.00]	0.001
er level cell percentage category (%)	<10%	8 (5.2)	3 (2.9)	24 (72.7)	<0.001
	10-19%	8 (5.2)	11 (10.6)	1 (3.0)	
	20-29%	2 (1.3)	0 (0.0)	2 (6.1)	
	30-39%	2 (1.3)	2 (1.9)	1 (3.0)	
	40-49%	1 (0.6)	2 (1.9)	0 (0.0)	
	50-59%	5 (3.2)	3 (2.9)	0 (0.0)	
	60-69%	4 (2.6)	3 (2.9)	0 (0.0)	
	70-79%	26 (16.8)	13 (12.5)	1 (3.0)	
	80-89%	12 (7.7)	6 (5.8)	0 (0.0)	
	90-99%	87 (56.1)	61 (58.7)	4 (12.1)	
progesterone receptor level cell percent category (%)	<10%	30 (20.7)	22 (24.7)	27 (93.1)	<0.001
	10-19%	9 (6.2)	9 (10.1)	0 (0.0)	
	20-29%	4 (2.8)	4 (4.5)	0 (0.0)	
	30-39%	4 (2.8)	4 (4.5)	0 (0.0)	
	40-49%	6 (4.1)	5 (5.6)	0 (0.0)	
	50-59%	7 (4.8)	3 (3.4)	0 (0.0)	
	60-69%	5 (3.4)	6 (6.7)	1 (3.4)	
	70-79%	22 (15.2)	7 (7.9)	0 (0.0)	
	80-89%	16 (11.0)	7 (7.9)	0 (0.0)	
	90-99%	42 (29.0)	22 (24.7)	1 (3.4)	
distant metastasis present ind2 (%)	NO	111 (96.5)	99 (99.0)	43 (97.7)	0.479
	YES	4 (3.5)	1 (1.0)	1 (2.3)	
stage event pathologic stage (%)	Stage I	28 (10.4)	13 (5.8)	9 (7.1)	0.020
	Stage IA	24 (9.0)	16 (7.1)	9 (7.1)	
	Stage IB	3 (1.1)	0 (0.0)	0 (0.0)	
	Stage II	2 (0.7)	2 (0.9)	1 (0.8)	
	Stage IIA	80 (29.9)	63 (28.0)	59 (46.5)	
	Stage IIB	64 (23.9)	57 (25.3)	27 (21.3)	
	Stage III	1 (0.4)	1 (0.4)	0 (0.0)	
	Stage IIIA	41 (15.3)	46 (20.4)	13 (10.2)	
	Stage IIIB	3 (1.1)	9 (4.0)	2 (1.6)	
	Stage IIIC	20 (7.5)	12 (5.3)	4 (3.1)	
	Stage IV	2 (0.7)	3 (1.3)	2 (1.6)	
	Stage X	0 (0.0)	3 (1.3)	1 (0.8)	