

# Summary\_of\_clinical\_variables

## Contents

Table 1: Summarization of clinical variables stratified by subtype

|                                                             | level                             | MDICC1                              | MDICC2                              | p      | test    |
|-------------------------------------------------------------|-----------------------------------|-------------------------------------|-------------------------------------|--------|---------|
| n                                                           |                                   | 248                                 | 216                                 |        |         |
| Subtype (%)                                                 | MDICC1                            | 248 (100.0)                         | 0 ( 0.0)                            | <0.001 |         |
|                                                             | MDICC2                            | 0 ( 0.0)                            | 216 (100.0)                         |        |         |
| vital status (%)                                            | Alive                             | 4 (100.0)                           | 7 ( 87.5)                           | 1.000  | exact   |
|                                                             | Dead                              | 0 ( 0.0)                            | 1 ( 12.5)                           |        |         |
| days to birth<br>(median [IQR])                             |                                   | -22997.00 [-26134.00,<br>-19188.00] | -20705.00 [-23667.00,<br>-17474.00] | <0.001 | nonnorm |
| days to death<br>(median [IQR])                             |                                   | 1563.00 [883.00,<br>2483.00]        | 1032.00 [683.00,<br>1742.50]        | 0.238  | nonnorm |
| days to last<br>followup<br>(median [IQR])                  |                                   | 752.00 [418.00,<br>1277.00]         | 1015.50 [445.50,<br>2112.75]        | 0.012  | nonnorm |
| race list (%)                                               | ASIAN                             | 10 ( 5.3)                           | 19 ( 9.7)                           | 0.001  | exact   |
|                                                             | BLACK OR AFRICAN<br>AMERICAN      | 16 ( 8.5)                           | 41 ( 21.0)                          |        |         |
|                                                             | WHITE                             | 162 ( 86.2)                         | 135 ( 69.2)                         |        |         |
| history of<br>neoadjuvant<br>treatment (%)                  | No                                | 242 ( 97.6)                         | 215 ( 99.5)                         | 0.128  | exact   |
|                                                             | Yes                               | 6 ( 2.4)                            | 1 ( 0.5)                            |        |         |
| age at initial<br>pathologic<br>diagnosis<br>(median [IQR]) |                                   | 62.00 [52.00, 71.00]                | 56.00 [47.00, 64.00]                | <0.001 | nonnorm |
| ethnicity (%)                                               | HISPANIC OR<br>LATINO             | 6 ( 3.5)                            | 4 ( 2.2)                            | 0.535  | exact   |
|                                                             | NOT HISPANIC OR<br>LATINO         | 167 ( 96.5)                         | 177 ( 97.8)                         |        |         |
| primary lymph<br>node<br>presentation<br>assessment<br>(%)  | NO                                | 2 ( 1.9)                            | 0 ( 0.0)                            | 0.498  | exact   |
|                                                             | YES                               | 106 ( 98.1)                         | 106 (100.0)                         |        |         |
| histological<br>type (%)                                    | Infiltrating Ductal<br>Carcinoma  | 175 ( 70.6)                         | 165 ( 76.4)                         | 0.264  | exact   |
|                                                             | Infiltrating Lobular<br>Carcinoma | 49 ( 19.8)                          | 37 ( 17.1)                          |        |         |
|                                                             | Medullary Carcinoma               | 0 ( 0.0)                            | 1 ( 0.5)                            |        |         |

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

|                                                             | level                                                                                        | MDICC1            | MDICC2            | p      | test    |
|-------------------------------------------------------------|----------------------------------------------------------------------------------------------|-------------------|-------------------|--------|---------|
|                                                             | Metaplastic Carcinoma                                                                        | 1 ( 0.4)          | 2 ( 0.9)          |        |         |
|                                                             | Mixed Histology (please specify)                                                             | 4 ( 1.6)          | 4 ( 1.9)          |        |         |
|                                                             | Mucinous Carcinoma                                                                           | 6 ( 2.4)          | 1 ( 0.5)          |        |         |
|                                                             | Other, specify                                                                               | 13 ( 5.2)         | 6 ( 2.8)          |        |         |
| menopause status (%)                                        | Indeterminate (neither Pre or Postmenopausal)                                                | 8 ( 3.4)          | 12 ( 5.9)         | 0.073  | exact   |
|                                                             | Peri (6-12 months since last menstrual period)                                               | 8 ( 3.4)          | 10 ( 5.0)         |        |         |
|                                                             | Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)            | 178 ( 75.4)       | 129 ( 63.9)       |        |         |
|                                                             | Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement) | 42 ( 17.8)        | 51 ( 25.2)        |        |         |
| breast carcinoma progesterone receptor status (%)           | Indeterminate                                                                                | 2 ( 0.8)          | 0 ( 0.0)          | <0.001 | exact   |
|                                                             | Negative                                                                                     | 38 ( 15.7)        | 122 ( 57.5)       |        |         |
|                                                             | Positive                                                                                     | 202 ( 83.5)       | 90 ( 42.5)        |        |         |
| breast carcinoma estrogen receptor status (%)               | Indeterminate                                                                                | 2 ( 0.8)          | 0 ( 0.0)          | <0.001 | exact   |
|                                                             | Negative                                                                                     | 11 ( 4.5)         | 100 ( 47.2)       |        |         |
|                                                             | Positive                                                                                     | 229 ( 94.6)       | 112 ( 52.8)       |        |         |
| lab proc her2 neu immuno-histochemistry receptor status (%) | Equivocal                                                                                    | 47 ( 21.3)        | 24 ( 12.9)        | 0.050  | exact   |
|                                                             | Indeterminate                                                                                | 1 ( 0.5)          | 1 ( 0.5)          |        |         |
|                                                             | Negative                                                                                     | 132 ( 59.7)       | 112 ( 60.2)       |        |         |
|                                                             | Positive                                                                                     | 41 ( 18.6)        | 49 ( 26.3)        |        |         |
| number of lymphnodes positive by ihc (median [IQR])         |                                                                                              | 0.00 [0.00, 0.00] | 0.00 [0.00, 0.00] | 0.412  | nonnorm |
| number of lymphnodes positive by he (median [IQR])          |                                                                                              | 0.00 [0.00, 2.00] | 0.00 [0.00, 2.00] | 0.513  | nonnorm |

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

|                                                                              | level      | MDICC1             | MDICC2            | p     | test    |
|------------------------------------------------------------------------------|------------|--------------------|-------------------|-------|---------|
| er level cell<br>percentage<br>category<br>(median [IQR])                    |            | 1.00 [1.00, 11.00] | 1.00 [1.00, 2.00] | 0.585 | nonnorm |
| progesterone<br>receptor level<br>cell percent<br>category<br>(median [IQR]) |            | 1.00 [1.00, 3.00]  | 1.00 [1.00, 2.00] | 0.723 | nonnorm |
| distant<br>metastasis<br>present ind2<br>(%)                                 | NO         | 72 ( 97.3)         | 57 ( 95.0)        | 0.656 | exact   |
|                                                                              | YES        | 2 ( 2.7)           | 3 ( 5.0)          |       |         |
| stage event<br>pathologic<br>stage (%)                                       | Stage I    | 20 ( 8.1)          | 20 ( 9.4)         | 0.192 | exact   |
|                                                                              | Stage IA   | 24 ( 9.8)          | 14 ( 6.6)         |       |         |
|                                                                              | Stage IB   | 3 ( 1.2)           | 0 ( 0.0)          |       |         |
|                                                                              | Stage IIA  | 87 ( 35.4)         | 69 ( 32.5)        |       |         |
|                                                                              | Stage IIB  | 53 ( 21.5)         | 53 ( 25.0)        |       |         |
|                                                                              | Stage IIIA | 25 ( 10.2)         | 29 ( 13.7)        |       |         |
|                                                                              | Stage IIIB | 9 ( 3.7)           | 3 ( 1.4)          |       |         |
|                                                                              | Stage IIIC | 11 ( 4.5)          | 17 ( 8.0)         |       |         |
|                                                                              | Stage IV   | 7 ( 2.8)           | 5 ( 2.4)          |       |         |
|                                                                              | Stage X    | 7 ( 2.8)           | 2 ( 0.9)          |       |         |