

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	MSNE1	MSNE2	MSNE3	MSNE4	MSNE5	p	test
n		72	112	96	66	118		
Subtype (%)	MSNE1	72 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	<0.001	
	MSNE2	0 (0.0)	112 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	MSNE3	0 (0.0)	0 (0.0)	96 (100.0)	0 (0.0)	0 (0.0)		
	MSNE4	0 (0.0)	0 (0.0)	0 (0.0)	66 (100.0)	0 (0.0)		
	MSNE5	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	118 (100.0)		
vital status (%)	Alive	0 (0.0)	1 (100.0)	3 (100.0)	4 (100.0)	3 (100.0)	0.163	exact
	Dead	1 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
days to birth (median [IQR])		-20589.50 [-17667.50, 22730.50]	-23900.00 [-21345.50, 27065.50]	-21479.00 [-18506.00, 25658.00]	-20471.00 [-17070.00, 23543.00]	-21710.50 [-17311.25, 24785.75]	<0.001	nonnorm
days to death (median [IQR])		1009.00 [548.00, 1556.00]	921.00 [468.00, 1920.00]	1286.00 [865.00, 2115.50]	1013.00 [672.00, 1695.25]	2000.00 [1220.75, 2867.00]	0.178	nonnorm
days to last followup (median [IQR])		997.00 [397.00, 1831.00]	734.00 [486.50, 1255.50]	797.00 [381.50, 1382.50]	941.50 [506.50, 2301.25]	987.00 [515.00, 2173.00]	0.114	nonnorm
race list (%)	ASIAN	4 (6.2)	4 (4.9)	11 (15.1)	4 (7.1)	6 (5.6)	<0.001	exact
	BLACK OR AFRICAN AMERICAN	13 (20.0)	9 (11.0)	9 (12.3)	18 (32.1)	8 (7.5)		
	WHITE	48 (73.8)	69 (84.1)	53 (72.6)	34 (60.7)	93 (86.9)		
history of neoadjuvant treatment (%)	No	72 (100.0)	111 (99.1)	92 (95.8)	66 (100.0)	116 (98.3)	0.203	exact

Table 1: Summarization of clinical variables stratified by subtype
(continued)

	level	MSNE1	MSNE2	MSNE3	MSNE4	MSNE5	p	test
age at initial pathologic diagnosis (median [IQR])	Yes	0 (0.0) 56.00 [48.00, 62.00]	1 (0.9) 65.00 [58.00, 74.00]	4 (4.2) 57.50 [50.00, 70.00]	0 (0.0) 55.50 [46.00, 64.00]	2 (1.7) 59.00 [47.00, 67.00]	<0.001	nonnorm
ethnicity (%)	HISPANIC OR LATINO	0 (0.0)	2 (2.7)	1 (1.4)	2 (3.8)	5 (5.1)	0.403	exact
	NOT HISPANIC OR LATINO	59 (100.0)	73 (97.3)	69 (98.6)	50 (96.2)	93 (94.9)		
primary lymph node pre-sentation assessment (%)	NO	0 (0.0)	1 (2.3)	0 (0.0)	1 (2.9)	0 (0.0)	0.439	exact
	YES	29 (100.0)	43 (97.7)	32 (100.0)	34 (97.1)	74 (100.0)		
histological type (%)	Infiltrating Ductal Carcinoma	71 (98.6)	78 (69.6)	83 (86.5)	51 (77.3)	57 (48.3)	<0.001	exact
	Infiltrating Lobular Carcinoma	0 (0.0)	16 (14.3)	9 (9.4)	8 (12.1)	53 (44.9)		
	Medullary Carcinoma	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.5)	0 (0.0)		
	Metaplastic Carcinoma	1 (1.4)	0 (0.0)	0 (0.0)	1 (1.5)	1 (0.8)		
	Mixed Histology (please specify)	0 (0.0)	2 (1.8)	1 (1.0)	3 (4.5)	2 (1.7)		
	Mucinous Carcinoma	0 (0.0)	5 (4.5)	1 (1.0)	0 (0.0)	1 (0.8)		
	Other, specify	0 (0.0)	11 (9.8)	2 (2.1)	2 (3.0)	4 (3.4)		
menopause status (%)	Indeterminat (neither Pre or Post-menopausal)	5 (7.6)	3 (2.8)	7 (7.6)	4 (6.3)	1 (0.9)	0.075	exact

Table 1: Summarization of clinical variables stratified by subtype
(continued)

	level	MSNE1	MSNE2	MSNE3	MSNE4	MSNE5	p	test
	Peri (6-12 months since last menstrual period)	4 (6.1)	3 (2.8)	5 (5.4)	1 (1.6)	5 (4.6)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	45 (68.2)	88 (80.7)	59 (64.1)	42 (66.7)	73 (67.6)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	12 (18.2)	15 (13.8)	21 (22.8)	16 (25.4)	29 (26.9)		
breast carcinoma progesterone receptor status (%)	Indeterminat	0 (0.0)	1 (0.9)	1 (1.1)	0 (0.0)	0 (0.0)	<0.001	exact
	Negative	64 (88.9)	12 (11.0)	28 (30.4)	38 (60.3)	18 (15.3)		
	Positive	8 (11.1)	96 (88.1)	63 (68.5)	25 (39.7)	100 (84.7)		
breast carcinoma estrogen receptor status (%)	Indeterminate	0 (0.0)	0 (0.0)	2 (2.2)	0 (0.0)	0 (0.0)	<0.001	exact
	Negative	59 (81.9)	0 (0.0)	8 (8.7)	36 (57.1)	8 (6.8)		
	Positive	13 (18.1)	109 (100.0)	82 (89.1)	27 (42.9)	110 (93.2)		
lab proc her2 neu immunohistochemistry receptor status (%)	Equivocal	11 (16.9)	26 (26.5)	11 (12.9)	9 (16.4)	14 (13.5)	<0.001	exact

Table 1: Summarization of clinical variables stratified by subtype
(continued)

	level	MSNE1	MSNE2	MSNE3	MSNE4	MSNE5	p	test
	Indeterminate	0 (0.0)	1 (1.0)	0 (0.0)	0 (0.0)	1 (1.0)		
	Negative	45 (69.2)	56 (57.1)	38 (44.7)	33 (60.0)	72 (69.2)		
	Positive	9 (13.8)	15 (15.3)	36 (42.4)	13 (23.6)	17 (16.3)		
number of lymphn-odes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 1.00]	0.193	nonnorm
number of lymphn-odes positive by he (median [IQR])		0.00 [0.00, 2.00]	0.00 [0.00, 1.00]	1.00 [0.00, 3.00]	0.00 [0.00, 1.00]	0.00 [0.00, 3.00]	0.012	nonnorm
er level cell percent-age category (median [IQR])		1.00 [1.00, 2.00]	1.00 [1.00, 11.00]	1.00 [1.00, 2.00]	1.00 [1.00, 2.00]	1.50 [1.00, 11.00]	<0.001	nonnorm
progesterone receptor level cell percent category (median [IQR])		1.00 [1.00, 2.00]	1.00 [1.00, 7.00]	1.00 [1.00, 1.00]	1.00 [1.00, 2.00]	1.00 [1.00, 5.75]	<0.001	nonnorm
distant metastasis present ind2 (%)	NO	15 (88.2)	29 (96.7)	20 (95.2)	18 (100.0)	47 (97.9)	0.374	exact
	YES	2 (11.8)	1 (3.3)	1 (4.8)	0 (0.0)	1 (2.1)		
stage event pathologic stage (%)	Stage I	4 (5.8)	14 (12.6)	4 (4.2)	7 (10.8)	11 (9.4)	0.006	exact
	Stage IA	6 (8.7)	13 (11.7)	2 (2.1)	7 (10.8)	10 (8.5)		
	Stage IB	0 (0.0)	3 (2.7)	0 (0.0)	0 (0.0)	0 (0.0)		
	Stage IIA	30 (43.5)	41 (36.9)	31 (32.3)	23 (35.4)	31 (26.5)		
	Stage IIB	17 (24.6)	23 (20.7)	25 (26.0)	13 (20.0)	28 (23.9)		
	Stage IIIA	9 (13.0)	8 (7.2)	17 (17.7)	4 (6.2)	16 (13.7)		
	Stage IIIB	0 (0.0)	4 (3.6)	4 (4.2)	3 (4.6)	1 (0.9)		
	Stage IIIC	2 (2.9)	2 (1.8)	5 (5.2)	5 (7.7)	14 (12.0)		
	Stage IV	1 (1.4)	1 (0.9)	6 (6.2)	2 (3.1)	2 (1.7)		
	Stage X	0 (0.0)	2 (1.8)	2 (2.1)	1 (1.5)	4 (3.4)		