

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	RWR-F1	RWR-F2	p	test
n		269	195		
Subtype (%)	RWR-F1	269 (100.0)	0 (0.0)	<0.001	
	RWR-F2	0 (0.0)	195 (100.0)		
vital status (%)	Alive	6 (100.0)	5 (83.3)	1.000	exact
	Dead	0 (0.0)	1 (16.7)		
days to birth (median [IQR])		-22833.00 [-26432.50, -18799.00]	-20675.00 [-24029.00, -17660.00]	<0.001	nonnorm
days to death (median [IQR])		1787.00 [888.75, 2813.50]	1020.50 [796.75, 1595.25]	0.078	nonnorm
days to last followup (median [IQR])		812.00 [456.00, 1547.00]	973.50 [392.75, 1848.00]	0.603	nonnorm
race list (%)	ASIAN	13 (5.9)	16 (9.9)	<0.001	exact
	BLACK OR AFRICAN AMERICAN	16 (7.2)	41 (25.3)		
	WHITE	192 (86.9)	105 (64.8)		
history of neoadjuvant treatment (%)	No	264 (98.1)	193 (99.0)	0.704	exact
	Yes	5 (1.9)	2 (1.0)		
age at initial pathologic diagnosis (median [IQR])		62.00 [51.00, 72.00]	56.00 [48.00, 65.00]	<0.001	nonnorm
ethnicity (%)	HISPANIC OR LATINO	9 (4.4)	1 (0.7)	0.049	exact
	NOT HISPANIC OR LATINO	196 (95.6)	148 (99.3)		
primary lymph node presentation assessment (%)	NO	2 (1.5)	0 (0.0)	0.535	exact
	YES	134 (98.5)	78 (100.0)		
histological type (%)	Infiltrating Ductal Carcinoma	162 (60.2)	178 (91.3)	<0.001	exact
	Infiltrating Lobular Carcinoma	79 (29.4)	7 (3.6)		
	Medullary Carcinoma	0 (0.0)	1 (0.5)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	RWR-F1	RWR-F2	p	test
	Metaplastic Carcinoma	0 (0.0)	3 (1.5)		
	Mixed Histology (please specify)	5 (1.9)	3 (1.5)		
	Mucinous Carcinoma	7 (2.6)	0 (0.0)		
	Other, specify	16 (5.9)	3 (1.5)		
menopause status (%)	Indeterminate (neither Pre or Postmenopausal)	10 (4.0)	10 (5.4)	0.238	exact
	Peri (6-12 months since last menstrual period)	10 (4.0)	8 (4.3)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	187 (73.9)	120 (64.9)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	46 (18.2)	47 (25.4)		
breast carcinoma progesterone receptor status (%)	Indeterminate	2 (0.8)	0 (0.0)	<0.001	exact
	Negative	33 (12.6)	127 (66.1)		
	Positive	227 (86.6)	65 (33.9)		
breast carcinoma estrogen receptor status (%)	Indeterminate	0 (0.0)	2 (1.0)	<0.001	exact
	Negative	4 (1.5)	107 (55.7)		
	Positive	258 (98.5)	83 (43.2)		
lab proc her2 neu immuno-histochemistry receptor status (%)	Equivocal	46 (19.8)	25 (14.3)	0.002	exact
	Indeterminate	2 (0.9)	0 (0.0)		
	Negative	147 (63.4)	97 (55.4)		
	Positive	37 (15.9)	53 (30.3)		
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.043	nonnorm
number of lymphnodes positive by he (median [IQR])		0.00 [0.00, 2.00]	0.00 [0.00, 2.00]	0.518	nonnorm

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	RWR-F1	RWR-F2	p	test
er level cell percentage category (median [IQR])		1.00 [1.00, 11.00]	1.00 [1.00, 2.00]	<0.001	nonnorm
progesterone receptor level cell percent category (median [IQR])		1.00 [1.00, 6.00]	1.00 [1.00, 2.00]	<0.001	nonnorm
distant metastasis present ind2 (%)	NO	82 (96.5)	47 (95.9)	1.000	exact
	YES	3 (3.5)	2 (4.1)		
stage event pathologic stage (%)	Stage I	30 (11.3)	10 (5.2)	0.054	exact
	Stage IA	24 (9.0)	14 (7.3)		
	Stage IB	2 (0.8)	1 (0.5)		
	Stage IIA	83 (31.2)	73 (38.0)		
	Stage IIB	55 (20.7)	51 (26.6)		
	Stage IIIA	32 (12.0)	22 (11.5)		
	Stage IIIB	6 (2.3)	6 (3.1)		
	Stage IIIC	18 (6.8)	10 (5.2)		
	Stage IV	7 (2.6)	5 (2.6)		
	Stage X	9 (3.4)	0 (0.0)		