

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	SNF1	SNF2	p	test
n		313	151		
Subtype (%)	SNF1	313 (100.0)	0 (0.0)	<0.001	
	SNF2	0 (0.0)	151 (100.0)		
vital status (%)	Alive	7 (100.0)	4 (80.0)	0.417	exact
	Dead	0 (0.0)	1 (20.0)		
days to birth (median [IQR])		-22710.00 [-26024.50, -18674.50]	-20332.00 [-23530.00, -17474.00]	<0.001	nonnorm
days to death (median [IQR])		1563.00 [883.00, 2573.00]	991.00 [651.00, 1624.00]	0.068	nonnorm
days to last followup (median [IQR])		857.00 [448.00, 1518.25]	972.00 [410.50, 1961.50]	0.479	nonnorm
race list (%)	ASIAN	17 (6.9)	12 (8.8)	<0.001	exact
	BLACK OR AFRICAN AMERICAN	20 (8.1)	37 (27.2)		
	WHITE	210 (85.0)	87 (64.0)		
history of neoadjuvant treatment (%)	No	306 (97.8)	151 (100.0)	0.102	exact
	Yes	7 (2.2)	0 (0.0)		
age at initial pathologic diagnosis (median [IQR])		62.00 [50.00, 71.00]	55.00 [47.00, 64.00]	<0.001	nonnorm
ethnicity (%)	HISPANIC OR LATINO	9 (3.9)	1 (0.8)	0.104	exact
	NOT HISPANIC OR LATINO	219 (96.1)	125 (99.2)		
primary lymph node presentation assessment (%)	NO	2 (1.4)	0 (0.0)	1.000	exact
	YES	143 (98.6)	69 (100.0)		
histological type (%)	Infiltrating Ductal Carcinoma	207 (66.1)	133 (88.1)	<0.001	exact
	Infiltrating Lobular Carcinoma	78 (24.9)	8 (5.3)		
	Medullary Carcinoma	0 (0.0)	1 (0.7)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	SNF1	SNF2	p	test
	Metaplastic Carcinoma	0 (0.0)	3 (2.0)		
	Mixed Histology (please specify)	6 (1.9)	2 (1.3)		
	Mucinous Carcinoma	7 (2.2)	0 (0.0)		
	Other, specify	15 (4.8)	4 (2.6)		
menopause status (%)	Indeterminate (neither Pre or Postmenopausal)	12 (4.1)	8 (5.6)	0.204	exact
	Peri (6-12 months since last menstrual period)	12 (4.1)	6 (4.2)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	216 (73.2)	91 (63.6)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	55 (18.6)	38 (26.6)		
breast carcinoma progesterone receptor status (%)	Indeterminate	2 (0.7)	0 (0.0)	<0.001	exact
	Negative	38 (12.4)	122 (82.4)		
	Positive	266 (86.9)	26 (17.6)		
breast carcinoma estrogen receptor status (%)	Indeterminate	2 (0.7)	0 (0.0)	<0.001	exact
	Negative	5 (1.6)	106 (71.6)		
	Positive	299 (97.7)	42 (28.4)		
lab proc her2 neu immuno-histochemistry receptor status (%)	Equivocal	51 (18.7)	20 (14.9)	0.225	exact
	Indeterminate	2 (0.7)	0 (0.0)		
	Negative	167 (61.2)	77 (57.5)		
	Positive	53 (19.4)	37 (27.6)		
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.063	nonnorm
number of lymphnodes positive by he (median [IQR])		0.00 [0.00, 2.00]	0.00 [0.00, 1.00]	0.091	nonnorm

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	SNF1	SNF2	p	test
er level cell percentage category (median [IQR])		1.00 [1.00, 11.00]	1.00 [1.00, 2.00]	0.003	nonnorm
progesterone receptor level cell percent category (median [IQR])		1.00 [1.00, 5.00]	1.00 [1.00, 2.00]	0.003	nonnorm
distant metastasis present ind2 (%)	NO	89 (96.7)	40 (95.2)	0.648	exact
	YES	3 (3.3)	2 (4.8)		
stage event pathologic stage (%)	Stage I	32 (10.3)	8 (5.4)	0.054	exact
	Stage IA	25 (8.1)	13 (8.8)		
	Stage IB	3 (1.0)	0 (0.0)		
	Stage IIA	95 (30.6)	61 (41.2)		
	Stage IIB	66 (21.3)	40 (27.0)		
	Stage IIIA	39 (12.6)	15 (10.1)		
	Stage IIIB	10 (3.2)	2 (1.4)		
	Stage IIIC	21 (6.8)	7 (4.7)		
	Stage IV	10 (3.2)	2 (1.4)		
	Stage X	9 (2.9)	0 (0.0)		