

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	Spectrum1	Spectrum2	p	test
n		150	314		
Subtype (%)	Spectrum1	150 (100.0)	0 (0.0)	<0.001	
	Spectrum2	0 (0.0)	314 (100.0)		
vital status (%)	Alive	4 (80.0)	7 (100.0)	0.417	exact
	Dead	1 (20.0)	0 (0.0)		
days to birth (median [IQR])		-20504.00 [-23628.00, -17569.00]	-22672.00 [-26024.50, -18611.00]	<0.001	nonnorm
days to death (median [IQR])		967.00 [599.50, 1485.00]	1602.50 [892.50, 2550.50]	0.045	nonnorm
days to last followup (median [IQR])		996.00 [414.50, 2033.00]	822.00 [446.50, 1504.00]	0.272	nonnorm
race list (%)	ASIAN	11 (8.2)	18 (7.2)	<0.001	exact
	BLACK OR AFRICAN AMERICAN	37 (27.6)	20 (8.0)		
	WHITE	86 (64.2)	211 (84.7)		
history of neoadjuvant treatment (%)	No	150 (100.0)	307 (97.8)	0.102	exact
	Yes	0 (0.0)	7 (2.2)		
age at initial pathologic diagnosis (median [IQR])		55.50 [48.00, 64.00]	61.50 [50.00, 71.00]	0.001	nonnorm
ethnicity (%)	HISPANIC OR LATINO	1 (0.8)	9 (3.9)	0.105	exact
	NOT HISPANIC OR LATINO	124 (99.2)	220 (96.1)		
primary lymph node presentation assessment (%)	NO	0 (0.0)	2 (1.4)	1.000	exact
	YES	70 (100.0)	142 (98.6)		
histological type (%)	Infiltrating Ductal Carcinoma	131 (87.3)	209 (66.6)	<0.001	exact
	Infiltrating Lobular Carcinoma	9 (6.0)	77 (24.5)		
	Medullary Carcinoma	1 (0.7)	0 (0.0)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	Spectrum1	Spectrum2	p	test
	Metaplastic Carcinoma	3 (2.0)	0 (0.0)		
	Mixed Histology (please specify)	2 (1.3)	6 (1.9)		
	Mucinous Carcinoma	0 (0.0)	7 (2.2)		
	Other, specify	4 (2.7)	15 (4.8)		
menopause status (%)	Indeterminate (neither Pre or Postmenopausal)	8 (5.6)	12 (4.1)	0.488	exact
	Peri (6-12 months since last menstrual period)	6 (4.2)	12 (4.1)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	93 (65.5)	214 (72.3)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	35 (24.6)	58 (19.6)		
breast carcinoma progesterone receptor status (%)	Indeterminate	0 (0.0)	2 (0.7)	<0.001	exact
	Negative	120 (81.6)	40 (13.0)		
	Positive	27 (18.4)	265 (86.3)		
breast carcinoma estrogen receptor status (%)	Indeterminate	0 (0.0)	2 (0.7)	<0.001	exact
	Negative	106 (72.1)	5 (1.6)		
	Positive	41 (27.9)	300 (97.7)		
lab proc her2 neu immuno-histochemistry receptor status (%)	Equivocal	20 (15.0)	51 (18.6)	0.383	exact
	Indeterminate	0 (0.0)	2 (0.7)		
	Negative	78 (58.6)	166 (60.6)		
	Positive	35 (26.3)	55 (20.1)		
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.073	nonnorm
number of lymphnodes positive by he (median [IQR])		0.00 [0.00, 1.00]	0.00 [0.00, 2.00]	0.095	nonnorm

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	Spectrum1	Spectrum2	p	test
er level cell percentage category (median [IQR])		1.00 [1.00, 2.00]	1.00 [1.00, 11.00]	0.001	nonnorm
progesterone receptor level cell percent category (median [IQR])		1.00 [1.00, 2.00]	1.00 [1.00, 5.00]	0.002	nonnorm
distant metastasis present ind2 (%)	NO	41 (95.3)	88 (96.7)	0.656	exact
	YES	2 (4.7)	3 (3.3)		
stage event pathologic stage (%)	Stage I	9 (6.1)	31 (10.0)	0.135	exact
	Stage IA	13 (8.8)	25 (8.0)		
	Stage IB	0 (0.0)	3 (1.0)		
	Stage IIA	59 (40.1)	97 (31.2)		
	Stage IIB	39 (26.5)	67 (21.5)		
	Stage IIIA	15 (10.2)	39 (12.5)		
	Stage IIIB	2 (1.4)	10 (3.2)		
	Stage IIIC	8 (5.4)	20 (6.4)		
	Stage IV	2 (1.4)	10 (3.2)		
	Stage X	0 (0.0)	9 (2.9)		