

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	ab-SNF1	ab-SNF2	p	test
n		318	146		
Subtype (%)	ab-SNF1	318 (100.0)	0 (0.0)	<0.001	
	ab-SNF2	0 (0.0)	146 (100.0)		
vital status (%)	Alive	7 (100.0)	4 (80.0)	0.417	exact
	Dead	0 (0.0)	1 (20.0)		
days to birth (median [IQR])		-22703.00 [-26024.50, -18674.50]	-20346.00 [-23530.00, -17474.00]	<0.001	nonnorm
days to death (median [IQR])		1602.50 [892.50, 2550.50]	967.00 [599.50, 1485.00]	0.045	nonnorm
days to last followup (median [IQR])		849.00 [447.00, 1516.00]	984.00 [410.75, 2003.00]	0.368	nonnorm
race list (%)	ASIAN	18 (7.2)	11 (8.3)	<0.001	exact
	BLACK OR AFRICAN AMERICAN	20 (8.0)	37 (28.0)		
	WHITE	213 (84.9)	84 (63.6)		
history of neoadjuvant treatment (%)	No	311 (97.8)	146 (100.0)	0.104	exact
	Yes	7 (2.2)	0 (0.0)		
age at initial pathologic diagnosis (median [IQR])		62.00 [50.00, 71.00]	55.00 [47.25, 64.00]	<0.001	nonnorm
ethnicity (%)	HISPANIC OR LATINO	9 (3.9)	1 (0.8)	0.174	exact
	NOT HISPANIC OR LATINO	222 (96.1)	122 (99.2)		
primary lymph node presentation assessment (%)	NO	2 (1.4)	0 (0.0)	1.000	exact
	YES	144 (98.6)	68 (100.0)		
histological type (%)	Infiltrating Ductal Carcinoma	212 (66.7)	128 (87.7)	<0.001	exact
	Infiltrating Lobular Carcinoma	78 (24.5)	8 (5.5)		
	Medullary Carcinoma	0 (0.0)	1 (0.7)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	ab-SNF1	ab-SNF2	p	test
	Metaplastic Carcinoma	0 (0.0)	3 (2.1)		
	Mixed Histology (please specify)	6 (1.9)	2 (1.4)		
	Mucinous Carcinoma	7 (2.2)	0 (0.0)		
	Other, specify	15 (4.7)	4 (2.7)		
menopause status (%)	Indeterminate (neither Pre or Postmenopausal)	12 (4.0)	8 (5.8)	0.349	exact
	Peri (6-12 months since last menstrual period)	12 (4.0)	6 (4.3)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	218 (72.7)	89 (64.5)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	58 (19.3)	35 (25.4)		
breast carcinoma progesterone receptor status (%)	Indeterminate	2 (0.6)	0 (0.0)	<0.001	exact
	Negative	41 (13.2)	119 (83.2)		
	Positive	268 (86.2)	24 (16.8)		
breast carcinoma estrogen receptor status (%)	Indeterminate	2 (0.6)	0 (0.0)	<0.001	exact
	Negative	5 (1.6)	106 (74.1)		
	Positive	304 (97.7)	37 (25.9)		
lab proc her2 neu immuno-histochemistry receptor status (%)	Equivocal	51 (18.3)	20 (15.5)	0.545	exact
	Indeterminate	2 (0.7)	0 (0.0)		
	Negative	168 (60.4)	76 (58.9)		
	Positive	57 (20.5)	33 (25.6)		
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.073	nonnorm
number of lymphnodes positive by he (median [IQR])		0.00 [0.00, 2.00]	0.00 [0.00, 1.00]	0.135	nonnorm

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	ab-SNF1	ab-SNF2	p	test
er level cell percentage category (median [IQR])		1.00 [1.00, 11.00]	1.00 [1.00, 2.00]	0.001	nonnorm
progesterone receptor level cell percent category (median [IQR])		1.00 [1.00, 4.75]	1.00 [1.00, 2.00]	0.004	nonnorm
distant metastasis present ind2 (%)	NO	90 (96.8)	39 (95.1)	0.641	exact
	YES	3 (3.2)	2 (4.9)		
stage event pathologic stage (%)	Stage I	32 (10.2)	8 (5.6)	0.109	exact
	Stage IA	25 (7.9)	13 (9.1)		
	Stage IB	3 (1.0)	0 (0.0)		
	Stage IIA	99 (31.4)	57 (39.9)		
	Stage IIB	67 (21.3)	39 (27.3)		
	Stage IIIA	39 (12.4)	15 (10.5)		
	Stage IIIB	10 (3.2)	2 (1.4)		
	Stage IIIC	21 (6.7)	7 (4.9)		
	Stage IV	10 (3.2)	2 (1.4)		
	Stage X	9 (2.9)	0 (0.0)		