

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	iClusterBayes1	iClusterBayes2	iClusterBayes4	iClusterBayes5
n		96	182	68	118
Subtype (%)	iClusterBayes1	96 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)
	iClusterBayes2	0 (0.0)	182 (100.0)	0 (0.0)	0 (0.0)
	iClusterBayes4	0 (0.0)	0 (0.0)	68 (100.0)	0 (0.0)
	iClusterBayes5	0 (0.0)	0 (0.0)	0 (0.0)	118 (100.0)
vital status (%)	Alive	5 (100.0)	5 (100.0)	0 (0.0)	1 (100.0)
	Dead	0 (0.0)	0 (0.0)	1 (100.0)	0 (0.0)
days to birth (median [IQR])		-20940.00 [-23536.50, -16980.50]	-22931.00 [-26686.25, -18868.75]	-20747.50 [-22799.75, -17613.00]	-21641.00 [-23536.50, -18519.50]
days to death (median [IQR])		1532.50 [993.75, 2671.00]	1642.00 [956.00, 2108.00]	991.00 [651.00, 1624.00]	1068.00 [719.00, 1919.50]
days to last followup (median [IQR])		1039.00 [621.50, 2249.50]	878.00 [539.50, 1540.00]	965.00 [411.00, 1842.00]	679.00 [354.00, 1359.00]
race list (%)	ASIAN	5 (5.7)	8 (5.4)	3 (4.8)	13 (11.0)
	BLACK OR AFRICAN AMERICAN	23 (26.1)	8 (5.4)	15 (24.2)	11 (12.7)
	WHITE	60 (68.2)	132 (89.2)	44 (71.0)	61 (71.7)
history of neoadjuvant treatment (%)	No	96 (100.0)	178 (97.8)	68 (100.0)	115 (97.4)
	Yes	0 (0.0)	4 (2.2)	0 (0.0)	3 (2.5)
age at initial pathologic diagnosis (median [IQR])		57.00 [46.00, 64.00]	62.00 [51.00, 72.75]	56.50 [48.00, 62.00]	59.00 [50.00, 64.00]
ethnicity (%)	HISPANIC OR LATINO	4 (4.7)	6 (4.4)	0 (0.0)	0 (0.0)
	NOT HISPANIC OR LATINO	81 (95.3)	130 (95.6)	55 (100.0)	78 (100.0)
primary lymph node presentation assessment (%)	NO	1 (1.6)	0 (0.0)	0 (0.0)	1 (2.5)
	YES	62 (98.4)	89 (100.0)	28 (100.0)	33 (97.5)
histological type (%)	Infiltrating Ductal Carcinoma	52 (54.2)	111 (61.0)	67 (98.5)	110 (93.2)
	Infiltrating Lobular Carcinoma	33 (34.4)	50 (27.5)	0 (0.0)	3 (2.5)

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	iClusterBayes1	iClusterBayes2	iClusterBayes4	iClusterBayes5
menopause status (%)	Medullary Carcinoma	1 (1.0)	0 (0.0)	0 (0.0)	0 (0.0)
	Metaplastic Carcinoma	2 (2.1)	0 (0.0)	1 (1.5)	0 (0.0)
	Mixed Histology (please specify)	4 (4.2)	3 (1.6)	0 (0.0)	1 (0.8)
	Mucinous Carcinoma	0 (0.0)	6 (3.3)	0 (0.0)	1 (0.8)
	Other, specify	4 (4.2)	12 (6.6)	0 (0.0)	3 (2.5)
	Indeterminate (neither Pre or Postmenopausal)	5 (5.7)	5 (2.9)	5 (8.1)	5 (4.4)
	Peri (6-12 months since last menstrual period)	3 (3.4)	7 (4.0)	3 (4.8)	5 (4.4)
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	58 (65.9)	126 (72.4)	43 (69.4)	80 (70.0)
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	22 (25.0)	36 (20.7)	11 (17.7)	24 (21.6)
	Indeterminate	0 (0.0)	1 (0.6)	0 (0.0)	1 (0.9)
breast carcinoma progesterone receptor status (%)					
	Negative	41 (44.1)	17 (9.6)	64 (94.1)	38 (32.8)
	Positive	52 (55.9)	159 (89.8)	4 (5.9)	77 (66.9)
breast carcinoma estrogen receptor status (%)	Indeterminate	0 (0.0)	0 (0.0)	0 (0.0)	2 (1.7)
	Negative	37 (39.8)	1 (0.6)	60 (88.2)	13 (11.1)
	Positive	56 (60.2)	176 (99.4)	8 (11.8)	101 (87.0)
lab proc her2 neu immuno-histochemistry receptor status (%)	Equivocal	7 (8.9)	34 (21.7)	13 (21.7)	17 (15.0)
	Indeterminate	1 (1.3)	1 (0.6)	0 (0.0)	0 (0.0)
	Negative	55 (69.6)	99 (63.1)	39 (65.0)	51 (45.0)
	Positive	16 (20.3)	23 (14.6)	8 (13.3)	43 (38.0)
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 1.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]
number of lymphnodes positive by he (median [IQR])		0.00 [0.00, 3.00]	0.00 [0.00, 2.00]	0.00 [0.00, 1.00]	1.00 [0.00, 3.00]

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	iClusterBayes1	iClusterBayes2	iClusterBayes4	iClusterBayes5
er level cell percentage category (median [IQR])		1.00 [1.00, 9.00]	1.00 [1.00, 11.00]	1.00 [1.00, 2.00]	1.00 [1.00, 2.00]
progesterone receptor level cell percent category (median [IQR])		1.00 [1.00, 2.00]	1.00 [1.00, 7.00]	1.00 [1.00, 2.00]	1.00 [1.00, 2.00]
distant metastasis present ind2 (%)	NO	33 (97.1)	54 (98.2)	15 (93.8)	27 (93.8)
	YES	1 (2.9)	1 (1.8)	1 (6.2)	2 (6.9)
stage event pathologic stage (%)	Stage I	9 (9.5)	19 (10.6)	4 (6.2)	8 (6.8)
	Stage IA	10 (10.5)	20 (11.1)	4 (6.2)	4 (3.4)
	Stage IB	0 (0.0)	2 (1.1)	0 (0.0)	1 (0.8)
	Stage IIA	23 (24.2)	69 (38.3)	33 (50.8)	31 (26.6)
	Stage IIB	21 (22.1)	32 (17.8)	16 (24.6)	37 (31.2)
	Stage IIIA	10 (10.5)	20 (11.1)	5 (7.7)	19 (16.1)
	Stage IIIB	4 (4.2)	4 (2.2)	0 (0.0)	4 (3.4)
	Stage IIIC	13 (13.7)	6 (3.3)	2 (3.1)	7 (5.9)
	Stage IV	2 (2.1)	4 (2.2)	1 (1.5)	5 (4.2)
	Stage X	3 (3.2)	4 (2.2)	0 (0.0)	2 (1.7)