

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	wMKL1	wMKL3	wMKL4	wMKL5	wMKL6	wMKL7	wMKL8	p	test
n		75	11	73	98	129	3	75		
Subtype (%)	wMKL1	75 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	<0.001	exact
	wMKL3	0 (0.0)	11 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	wMKL4	0 (0.0)	0 (0.0)	73 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	wMKL5	0 (0.0)	0 (0.0)	0 (0.0)	98 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	wMKL6	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	129 (100.0)	0 (0.0)	0 (0.0)		
	wMKL7	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	3 (100.0)	0 (0.0)		
	wMKL8	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	75 (100.0)		
vital status (%)	Alive	1 (100.0)	1 (100.0)	0 (NaN)	4 (80.0)	3 (100.0)	0 (NaN)	2 (100.0)	1.000	exact
	Dead	0 (0.0)	0 (0.0)	0 (NaN)	1 (20.0)	0 (0.0)	0 (NaN)	0 (0.0)		
days to birth (median [IQR])		-21793.00 [-, 24527.00, -18364.50]	-24439.00 [-, 27577.50, -18934.50]	-22723.00 [-, 25010.00, -19533.00]	-19298.00 [-, 22707.00, -16817.00]	-22710.00 [-, 25628.50, -18512.50]	-27121.00 [-, 28067.50, -22828.00]	-23465.50 [-, 26794.25, -19742.75]	<0.001	nonnorm
days to death (median [IQR])		1739.50 [995.25, 2965.00]	NA [NA, NA]	1430.00 [468.00, 1781.00]	1032.00 [754.00, 1692.00]	1134.50 [829.00, 2360.50]	NA [NA, NA]	1500.00 [973.50, 2493.00]	0.652	nonnorm
days to last followup (median [IQR])		987.00 [582.50, 2190.50]	754.00 [376.50, 2172.00]	808.00 [516.75, 1354.00]	996.00 [440.00, 2109.00]	955.50 [380.50, 1488.00]	762.00 [762.00, 762.00]	803.00 [487.00, 1139.50]	0.775	nonnorm
race list (%)	ASIAN	3 (4.3)	0 (0.0)	3 (4.8)	6 (6.6)	15 (16.5)	0 (0.0)	2 (3.6)	0.002	exact
	BLACK OR AFRICAN-AMERICAN	8 (11.4)	3 (27.3)	5 (7.9)	25 (27.5)	12 (13.2)	0 (0.0)	4 (7.3)		
history of neoadjuvant treatment (%)	WHITE	59 (84.3)	8 (72.7)	55 (87.3)	60 (65.9)	64 (70.3)	2 (100.0)	49 (89.1)		
	No	73 (97.3)	11 (100.0)	72 (98.6)	98 (100.0)	126 (97.7)	3 (100.0)	74 (98.7)	0.661	exact
	Yes	2 (2.7)	0 (0.0)	1 (1.4)	0 (0.0)	3 (2.3)	0 (0.0)	1 (1.3)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	wMKL1	wMKL3	wMKL4	wMKL5	wMKL6	wMKL7	wMKL8	p	test
age at initial pathologic diagnosis (median [IQR])		59.00 [50.00, 66.50]	66.00 [51.00, 75.00]	62.00 [53.00, 68.00]	52.00 [46.00, 62.00]	60.00 [50.00, 70.00]	74.00 [62.00, 76.50]	63.00 [53.50, 72.50]	<0.001	nonnorm
ethnicity (%)	HISPANIC OR LATINO	3 (4.7)	0 (0.0)	4 (6.8)	1 (1.2)	1 (1.2)	0 (0.0)	1 (2.0)	0.385	exact
	NOT HISPANIC OR LATINO	61 (95.3)	11 (100.0)	55 (93.2)	82 (98.8)	84 (98.8)	2 (100.0)	49 (98.0)		
primary lymph node presentation as assessment (%)	NO	1 (2.2)	0 (0.0)	1 (2.5)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0.828	exact
	YES	44 (97.8)	9 (100.0)	39 (97.5)	47 (100.0)	33 (100.0)	2 (100.0)	38 (100.0)		
histological type (%)	Infiltrating	44 (58.7)	3 (27.3)	46 (63.0)	89 (90.8)	117 (90.7)	1 (33.3)	40 (53.3)	<0.001	exact
	Ductal Carcinoma									
	Infiltrating Lobular Carcinoma	27 (36.0)	4 (36.4)	19 (26.0)	2 (2.0)	5 (3.9)	2 (66.7)	27 (36.0)		
	Medullary Carcinoma	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	Metaplastic Carcinoma	0 (0.0)	0 (0.0)	0 (0.0)	2 (2.0)	0 (0.0)	0 (0.0)	1 (1.3)		
	Mixed Histology (please specify)	1 (1.3)	0 (0.0)	2 (2.7)	2 (2.0)	1 (0.8)	0 (0.0)	2 (2.7)		
	Mucinous Carcinoma	0 (0.0)	3 (27.3)	2 (2.7)	0 (0.0)	1 (0.8)	0 (0.0)	1 (1.3)		
	Other, specify	3 (4.0)	1 (9.1)	4 (5.5)	2 (2.0)	5 (3.9)	0 (0.0)	4 (5.3)		
menopausal status (%)	Indeterminate (neither Pre or Postmenopausal)	2 (3.0)	0 (0.0)	1 (1.5)	7 (7.7)	8 (6.3)	0 (0.0)	2 (2.7)	0.267	exact

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	wMKL1	wMKL3	wMKL4	wMKL5	wMKL6	wMKL7	wMKL8	p	test
	Peri (6-12 months since last menstrual period)	5 (7.6)	0 (0.0)	2 (3.0)	4 (4.4)	5 (4.0)	1 (33.3)	1 (1.4)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	47 (71.2)	8 (72.7)	51 (76.1)	54 (59.3)	85 (67.5)	2 (66.7)	60 (81.1)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	12 (18.2)	3 (27.3)	13 (19.4)	26 (28.6)	28 (22.2)	0 (0.0)	11 (14.9)		
breast carcinoma progesterone receptor status (%)	Indeterm	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	2 (1.6)	0 (0.0)	0 (0.0)	<0.001	exact
	Negative	11 (14.7)	3 (27.3)	10 (13.9)	86 (90.5)	41 (32.3)	1 (33.3)	8 (11.3)		
	Positive	64 (85.3)	8 (72.7)	62 (86.1)	9 (9.5)	84 (66.1)	2 (66.7)	63 (88.7)		
breast carcinoma estrogen receptor status (%)	Indeterminate	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	2 (1.6)	0 (0.0)	0 (0.0)	<0.001	exact
	Negative	5 (6.7)	1 (9.1)	1 (1.4)	84 (88.4)	16 (12.6)	0 (0.0)	4 (5.6)		
	Positive	70 (93.3)	10 (90.9)	71 (98.6)	11 (11.6)	109 (85.8)	3 (100.0)	67 (94.4)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	wMKL1	wMKL3	wMKL4	wMKL5	wMKL6	wMKL7	wMKL8	p	test
lab proc her2 neu im- muno- histo- chem- istry recep- tor status (%)	Equivocal	9 (14.1)	1 (10.0)	16 (25.0)	12 (14.5)	20 (16.9)	1 (33.3)	12 (18.5)	0.001	exact
	Indeterminate	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	2 (3.1)		
	Negative	42 (65.6)	8 (80.0)	41 (64.1)	61 (73.5)	50 (42.4)	1 (33.3)	41 (63.1)		
	Positive	13 (20.3)	1 (10.0)	7 (10.9)	10 (12.0)	48 (40.7)	1 (33.3)	10 (15.4)		
number of lymph- nodes positive by ihc (me- dian [IQR])		0.00 [0.00, 0.75]	0.00 [0.00, 0.00]	0.00 [0.00, 0.50]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	2.00 [2.00, 2.00]	0.00 [0.00, 0.00]	0.081	nonnorm
number of lymph- nodes positive by he (me- dian [IQR])		0.00 [0.00, 3.00]	1.00 [0.00, 3.75]	0.00 [0.00, 2.00]	0.00 [0.00, 1.00]	0.50 [0.00, 2.00]	6.00 [3.00, 9.00]	0.00 [0.00, 1.00]	0.243	nonnorm
er level cell per- cent- age cate- gory (me- dian [IQR])		1.00 [1.00, 11.00]	9.00 [2.50, 11.00]	1.00 [1.00, 11.00]	1.00 [1.00, 2.00]	1.00 [1.00, 2.00]	1.00 [1.00, 6.00]	1.00 [1.00, 11.00]	<0.001	nonnorm
progesterone recep- tor level cell per- cent cate- gory (me- dian [IQR])		1.00 [1.00, 4.00]	5.00 [1.00, 10.00]	1.00 [1.00, 6.00]	1.00 [1.00, 2.00]	1.00 [1.00, 2.00]	1.00 [1.00, 6.00]	1.00 [1.00, 5.50]	0.001	nonnorm
distant metas- tasis present ind2 (%)	NO	28 (93.3)	5 (100.0)	22 (95.7)	26 (92.9)	24 (100.0)	1 (100.0)	23 (100.0)	0.626	exact
	YES	2 (6.7)	0 (0.0)	1 (4.3)	2 (7.1)	0 (0.0)	0 (0.0)	0 (0.0)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	wMKL1	wMKL3	wMKL4	wMKL5	wMKL6	wMKL7	wMKL8	p	test
stage event patho- logic stage (%)	Stage I	9 (12.2)	1 (9.1)	9 (12.5)	6 (6.3)	7 (5.4)	0 (0.0)	8 (10.8)	0.025	exact
	Stage IA	8 (10.8)	0 (0.0)	8 (11.1)	8 (8.4)	4 (3.1)	0 (0.0)	10 (13.5)		
	Stage IB	0 (0.0)	0 (0.0)	1 (1.4)	0 (0.0)	2 (1.6)	0 (0.0)	0 (0.0)		
	Stage IIA	21 (28.4)	4 (36.4)	20 (27.8)	42 (44.2)	35 (27.1)	1 (33.3)	33 (44.6)		
	Stage IIB	13 (17.6)	2 (18.2)	18 (25.0)	23 (24.2)	38 (29.5)	0 (0.0)	12 (16.2)		
	Stage IIIA	6 (8.1)	3 (27.3)	9 (12.5)	9 (9.5)	22 (17.1)	1 (33.3)	4 (5.4)		
	Stage IIIB	1 (1.4)	0 (0.0)	2 (2.8)	2 (2.1)	6 (4.7)	0 (0.0)	1 (1.4)		
	Stage IIIC	10 (13.5)	1 (9.1)	3 (4.2)	3 (3.2)	7 (5.4)	1 (33.3)	3 (4.1)		
	Stage IV	4 (5.4)	0 (0.0)	0 (0.0)	2 (2.1)	5 (3.9)	0 (0.0)	1 (1.4)		
	Stage X	2 (2.7)	0 (0.0)	2 (2.8)	0 (0.0)	3 (2.3)	0 (0.0)	2 (2.7)		