

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	MONET1	MONET2	MONET3	p	test
n		177	263	185		
Subtype (%)	MONET1	177 (100.0)	0 (0.0)	0 (0.0)	<0.001	
	MONET2	0 (0.0)	263 (100.0)	0 (0.0)		
	MONET3	0 (0.0)	0 (0.0)	185 (100.0)		
vital status (%)	Alive	45 (93.8)	63 (94.0)	55 (93.2)	1.000	exact
	Dead	3 (6.2)	4 (6.0)	4 (6.8)		
days to birth (median [IQR])		-20118.00 [-23647.50, -17981.00]	-21559.00 [-24868.00, -17691.00]	-21802.00 [-25224.75, -18215.75]	0.202	nonnorm
days to death (median [IQR])		912.00 [547.50, 2232.50]	1590.50 [779.50, 2075.25]	1694.00 [723.00, 2469.00]	0.665	nonnorm
days to last followup (median [IQR])		753.50 [445.50, 1645.25]	798.00 [504.00, 1611.00]	714.00 [463.00, 1528.00]	0.692	nonnorm
race list (%)	AMERICAN INDIAN OR ALASKA NATIVE	1 (0.6)	0 (0.0)	0 (0.0)	<0.001	exact
	ASIAN	10 (5.7)	10 (3.9)	12 (6.6)		
	BLACK OR AFRICAN AMERICAN	57 (32.8)	32 (12.5)	35 (19.3)		
	WHITE	106 (60.9)	215 (83.7)	134 (74.0)		

Table 1: Summarization of clinical variables stratified by subtype
(continued)

	level	MONET1	MONET2	MONET3	p	test
history of neoadju- vant treatment (%)	No	175 (99.4)	261 (99.2)	181 (98.4)	0.592	exact
	Yes	1 (0.6)	2 (0.8)	3 (1.6)		
age at initial pathologic diagnosis (median [IQR])		55.00 [49.00, 64.00]	59.00 [48.00, 67.50]	59.00 [49.00, 68.00]	0.239	nonnorm
ethnicity (%)	HISPANIC OR LATINO	8 (4.9)	10 (4.3)	9 (5.4)	0.882	
	NOT HISPANIC OR LATINO	155 (95.1)	223 (95.7)	159 (94.6)		
primary lymph node pre- sentation assess- ment (%)	NO	9 (6.1)	4 (1.8)	13 (9.8)	0.004	
	YES	139 (93.9)	215 (98.2)	120 (90.2)		
histological type (%)	Infiltrating Carcinoma NOS	1 (0.6)	0 (0.0)	0 (0.0)	<0.001	exact
	Infiltrating Ductal Carcinoma	146 (83.0)	135 (51.3)	155 (83.8)		
	Infiltrating Lobular Carcinoma	9 (5.1)	94 (35.7)	15 (8.1)		
	Medullary Carcinoma	5 (2.8)	0 (0.0)	0 (0.0)		
	Metaplastic Carcinoma	6 (3.4)	1 (0.4)	0 (0.0)		
	Mixed Histology (please specify)	1 (0.6)	15 (5.7)	5 (2.7)		
	Mucinous Carcinoma	0 (0.0)	7 (2.7)	3 (1.6)		
	Other, specify	8 (4.5)	11 (4.2)	7 (3.8)		

Table 1: Summarization of clinical variables stratified by subtype
(continued)

	level	MONET1	MONET2	MONET3	p	test
menopause status (%)	Indeterminat (neither Pre or Post-menopausal)	6 (3.7)	5 (2.1)	3 (1.8)	0.605	exact
	Peri (6-12 months since last menstrual period)	8 (4.9)	7 (2.9)	5 (2.9)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	116 (71.6)	166 (68.9)	123 (71.9)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	32 (19.8)	63 (26.1)	40 (23.4)		
breast carcinoma progesterone receptor status (%)	Indeterminat	1 (0.6)	1 (0.4)	0 (0.0)	<0.001	exact
	Negative	121 (74.2)	28 (11.2)	35 (20.2)		
	Positive	41 (25.2)	221 (88.4)	138 (79.8)		
breast carcinoma estrogen receptor status (%)	Negative	114 (69.5)	8 (3.2)	8 (4.6)	<0.001	
	Positive	50 (30.5)	242 (96.8)	165 (95.4)		

Table 1: Summarization of clinical variables stratified by subtype
(continued)

	level	MONET1	MONET2	MONET3	p	test
lab proc her2 neu immuno- histochem- istry receptor status (%)	Equivocal	31 (22.0)	42 (19.7)	37 (24.2)	0.009	exact
	Indeterminat	3 (2.1)	2 (0.9)	5 (3.3)		
	Negative	91 (64.5)	147 (69.0)	78 (51.0)		
	Positive	16 (11.3)	22 (10.3)	33 (21.6)		
number of lymphn- odes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.040	nonnorm
number of lymphn- odes positive by he (median [IQR])		0.00 [0.00, 2.00]	1.00 [0.00, 2.00]	1.00 [0.00, 4.00]	<0.001	nonnorm
er level cell percent- age category (%)	<10%	29 (51.8)	3 (2.2)	3 (2.9)	<0.001	exact
	10-19%	5 (8.9)	3 (2.2)	12 (11.8)		
	20-29%	2 (3.6)	2 (1.5)	0 (0.0)		
	30-39%	1 (1.8)	3 (2.2)	1 (1.0)		
	40-49%	0 (0.0)	1 (0.7)	2 (2.0)		
	50-59%	1 (1.8)	3 (2.2)	4 (3.9)		
	60-69%	0 (0.0)	3 (2.2)	4 (3.9)		
	70-79%	4 (7.1)	22 (16.4)	14 (13.7)		
	80-89%	1 (1.8)	10 (7.5)	7 (6.9)		
	90-99%	13 (23.2)	84 (62.7)	55 (53.9)		
progesterone receptor level cell percent category (%)	<10%	36 (73.5)	19 (14.6)	24 (28.6)	<0.001	exact
	10-19%	1 (2.0)	8 (6.2)	9 (10.7)		
	20-29%	0 (0.0)	6 (4.6)	2 (2.4)		
	30-39%	0 (0.0)	3 (2.3)	5 (6.0)		
	40-49%	0 (0.0)	5 (3.8)	6 (7.1)		
	50-59%	0 (0.0)	7 (5.4)	3 (3.6)		

Table 1: Summarization of clinical variables stratified by subtype
(continued)

	level	MONET1	MONET2	MONET3	p	test
	60-69%	2 (4.1)	5 (3.8)	5 (6.0)		
	70-79%	4 (8.2)	17 (13.1)	8 (9.5)		
	80-89%	0 (0.0)	18 (13.8)	5 (6.0)		
	90-99%	6 (12.2)	42 (32.3)	17 (20.2)		
distant metastasis present ind2 (%)	NO	59 (95.2)	120 (98.4)	74 (98.7)	0.439	exact
	YES	3 (4.8)	2 (1.6)	1 (1.3)		
stage event pathologic stage (%)	Stage I	15 (8.5)	27 (10.3)	8 (4.4)	0.004	exact
	Stage IA	11 (6.2)	26 (10.0)	12 (6.6)		
	Stage IB	0 (0.0)	3 (1.1)	0 (0.0)		
	Stage II	1 (0.6)	3 (1.1)	1 (0.5)		
	Stage IIA	78 (44.3)	70 (26.8)	54 (29.5)		
	Stage IIB	36 (20.5)	70 (26.8)	42 (23.0)		
	Stage III	1 (0.6)	0 (0.0)	1 (0.5)		
	Stage IIIA	20 (11.4)	37 (14.2)	43 (23.5)		
	Stage IIIB	3 (1.7)	5 (1.9)	6 (3.3)		
	Stage IIIC	7 (4.0)	18 (6.9)	11 (6.0)		
	Stage IV	3 (1.7)	2 (0.8)	2 (1.1)		
	Stage X	1 (0.6)	0 (0.0)	3 (1.6)		