

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	RWR-F1	RWR-F2	p	test
n		358	267		
Subtype (%)	RWR-F1	358 (100.0)	0 (0.0)	<0.001	
	RWR-F2	0 (0.0)	267 (100.0)		
vital status (%)	Alive	99 (95.2)	64 (91.4)	0.353	exact
	Dead	5 (4.8)	6 (8.6)		
days to birth (median [IQR])		-21335.00 [-24251.25, -17702.00]	-21326.50 [-24846.00, -18102.75]	0.848	nonnorm
days to death (median [IQR])		1365.00 [574.00, 2235.00]	1388.00 [573.00, 2551.00]	0.683	nonnorm
days to last followup (median [IQR])		851.50 [499.50, 1760.25]	670.50 [440.50, 1444.00]	0.044	nonnorm
race list (%)	AMERICAN INDIAN OR ALASKA NATIVE	0 (0.0)	1 (0.4)	<0.001	exact
	ASIAN	15 (4.3)	17 (6.5)		
	BLACK OR AFRICAN AMERICAN	53 (15.1)	71 (27.1)		
	WHITE	282 (80.6)	173 (66.0)		
history of neoadjuvant treatment (%)	No	356 (99.7)	261 (98.1)	0.089	exact
	Yes	1 (0.3)	5 (1.9)		
age at initial pathologic diagnosis (median [IQR])		58.00 [48.00, 65.75]	58.00 [49.00, 67.00]	0.888	nonnorm
ethnicity (%)	HISPANIC OR LATINO	17 (5.3)	10 (4.1)	0.638	
	NOT HISPANIC OR LATINO	303 (94.7)	234 (95.9)		
primary lymph node presentation assessment (%)	NO	11 (3.8)	15 (7.0)	0.170	
	YES	275 (96.2)	199 (93.0)		
histological type (%)	Infiltrating Carcinoma	0 (0.0)	1 (0.4)	<0.001	exact
	NOS				
	Infiltrating Ductal Carcinoma	224 (62.6)	212 (79.7)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	RWR-F1	RWR-F2	p	test
menopause status (%)	Infiltrating Lobular Carcinoma	95 (26.5)	23 (8.6)	0.189	
	Medullary Carcinoma	0 (0.0)	5 (1.9)		
	Metaplastic Carcinoma	2 (0.6)	5 (1.9)		
	Mixed Histology (please specify)	13 (3.6)	8 (3.0)		
	Mucinous Carcinoma	9 (2.5)	1 (0.4)		
	Other, specify	15 (4.2)	11 (4.1)		
	Indeterminate (neither Pre or Postmenopausal)	9 (2.7)	5 (2.1)		
	Peri (6-12 months since last menstrual period)	10 (3.0)	10 (4.1)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	224 (67.7)	181 (74.5)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	88 (26.6)	47 (19.3)		
breast carcinoma progesterone receptor status (%)	Indeterminate	1 (0.3)	1 (0.4)	<0.001	exact
breast carcinoma estrogen receptor status (%)	Negative	55 (16.2)	129 (52.2)	<0.001	
	Positive	283 (83.5)	117 (47.4)		
	Negative	22 (6.5)	108 (43.5)		
lab proc her2 neu immuno-histochemistry receptor status (%)	Positive	317 (93.5)	140 (56.5)	0.083	exact
	Equivocal	61 (21.0)	49 (22.7)		
	Indeterminate	4 (1.4)	6 (2.8)		
	Negative	193 (66.3)	123 (56.9)		
	Positive	33 (11.3)	38 (17.6)		
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.003	nonnorm

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	RWR-F1	RWR-F2	p	test
number of lymphnodes positive by he (median [IQR])		1.00 [0.00, 3.00]	0.50 [0.00, 3.00]	0.143	nonnorm
er level cell percentage category (%)	<10%	9 (4.7)	26 (26.0)	<0.001	exact
	10-19%	10 (5.2)	10 (10.0)		
	20-29%	0 (0.0)	4 (4.0)		
	30-39%	2 (1.0)	3 (3.0)		
	40-49%	2 (1.0)	1 (1.0)		
	50-59%	7 (3.6)	1 (1.0)		
	60-69%	5 (2.6)	2 (2.0)		
	70-79%	29 (15.1)	11 (11.0)		
	80-89%	14 (7.3)	4 (4.0)		
	90-99%	114 (59.4)	38 (38.0)		
progesterone receptor level cell percent category (%)	<10%	37 (20.4)	42 (51.2)	0.001	exact
	10-19%	11 (6.1)	7 (8.5)		
	20-29%	4 (2.2)	4 (4.9)		
	30-39%	6 (3.3)	2 (2.4)		
	40-49%	9 (5.0)	2 (2.4)		
	50-59%	10 (5.5)	0 (0.0)		
	60-69%	10 (5.5)	2 (2.4)		
	70-79%	24 (13.3)	5 (6.1)		
	80-89%	18 (9.9)	5 (6.1)		
	90-99%	52 (28.7)	13 (15.9)		
distant metastasis present ind2 (%)	NO	150 (96.8)	103 (99.0)	0.406	exact
	YES	5 (3.2)	1 (1.0)		
stage event pathologic stage (%)	Stage I	33 (9.3)	17 (6.4)	0.199	exact
	Stage IA	32 (9.0)	17 (6.4)		
	Stage IB	3 (0.8)	0 (0.0)		
	Stage II	3 (0.8)	2 (0.8)		
	Stage IIA	98 (27.6)	104 (39.2)		
	Stage IIB	91 (25.6)	57 (21.5)		
	Stage III	2 (0.6)	0 (0.0)		
	Stage IIIA	59 (16.6)	41 (15.5)		
	Stage IIIB	7 (2.0)	7 (2.6)		
	Stage IIIC	20 (5.6)	16 (6.0)		
	Stage IV	4 (1.1)	3 (1.1)		
	Stage X	3 (0.8)	1 (0.4)		