

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	NEMO1	NEMO2	NEMO3	NEMO4	NEMO5	NEMO6	NEMO7	NEMO8	NEMO9	p	test
n		308	113	96	4	16	11	7	32	38		
Subtype (%)	NEMO1	308 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	<0.001	exact
	NEMO2	0 (0.0)	113 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	NEMO3	0 (0.0)	0 (0.0)	96 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	NEMO4	0 (0.0)	0 (0.0)	0 (0.0)	4 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	NEMO5	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	16 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	NEMO6	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	11 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	NEMO7	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	7 (100.0)	0 (0.0)	0 (0.0)		
	NEMO8	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	32 (100.0)	0 (0.0)		
	NEMO9	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	38 (100.0)		
vital status (%)		7 (8.0)	1 (3.0)	2 (4.4)	0 (0.0)	0 (0.0)	0 (0.0)	1 (25.0)	0 (0.0)	0 (0.0)	0.843	exact
	Alive	76 (87.4)	28 (84.8)	40 (88.9)	1 (100.0)	4 (100.0)	2 (100.0)	3 (75.0)	6 (100.0)	3 (100.0)		
	Dead	4 (4.6)	4 (12.1)	3 (6.7)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
days to birth (median [IQR])		-21356.00 [-17756.75]	-19621.50 [-17467.25]	-21301.50 [-25118.00, -16670.75]	-20174.50 [-26112.00, -14246.25]	-22115.50 [-25038.25, -19485.75]	-22966.00 [-30254.00, -18987.00]	-23266.00 [-24486.00, -21519.00]	-22416.00 [-25998.00, -20067.50]	-22340.00 [-25793.00, -19849.00]	0.028	nonnorm
days to death (median [IQR])		1683.50 [677.00, 2483.25]	785.00 [524.00, 2192.00]	2443.00 [1779.00, 2766.00]	NA [NA, NA]	1759.00 [1562.00, 2155.00]	266.00 [266.00, 266.00]	NA [NA, NA]	769.50 [567.25, 971.75]	959.00 [371.50, 1600.00]	0.131	nonnorm
days to last followup (median [IQR])		847.00 [503.00, 1662.00]	701.00 [488.50, 1478.50]	785.00 [515.50, 2187.00]	614.00 [604.50, 876.50]	795.00 [294.00, 1273.50]	448.00 [368.00, 1611.00]	1326.00 [811.00, 1566.00]	594.00 [412.50, 1621.50]	506.50 [387.25, 758.25]	0.293	nonnorm
race list (%)		5 (1.6)	2 (1.8)	5 (5.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (2.6)	<0.001	exact
	AMERICAN INDIAN OR ALASKA NATIVE	0 (0.0)	1 (0.9)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	ASIAN	11 (3.6)	4 (3.5)	6 (6.2)	0 (0.0)	0 (0.0)	1 (9.1)	0 (0.0)	5 (15.6)	5 (13.2)		
	BLACK OR AFRICAN AMERICAN	44 (14.3)	41 (36.3)	18 (18.8)	2 (50.0)	3 (18.8)	5 (45.5)	1 (14.3)	8 (25.0)	2 (5.3)		
	WHITE	248 (80.5)	65 (57.5)	67 (69.8)	2 (50.0)	13 (81.2)	5 (45.5)	6 (85.7)	19 (59.4)	30 (78.9)	0.100	exact
history of neoadjuvant treatment (%)	No	306 (99.4)	112 (99.1)	94 (97.9)	4 (100.0)	16 (100.0)	11 (100.0)	7 (100.0)	32 (100.0)	35 (92.1)		
	Yes	2 (0.6)	0 (0.0)	1 (1.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	3 (7.9)	0.034	nonnorm
age at initial pathologic diagnosis (median [IQR])		58.00 [48.00, 66.00]	53.00 [47.00, 63.00]	58.00 [45.00, 67.50]	55.00 [38.50, 71.25]	60.00 [53.00, 68.00]	62.00 [51.50, 82.50]	63.00 [58.50, 66.50]	61.00 [53.75, 71.00]	60.50 [53.25, 69.75]		
ethnicity (%)		32 (10.4)	10 (8.8)	11 (11.5)	0 (0.0)	0 (0.0)	2 (18.2)	0 (0.0)	4 (12.5)	2 (5.3)	0.828	exact
	HISPANIC OR LATINO	11 (3.6)	6 (5.3)	6 (6.2)	0 (0.0)	0 (0.0)	0 (0.0)	1 (14.3)	2 (6.2)	1 (2.6)		
	NOT HISPANIC OR LATINO	265 (86.0)	97 (85.8)	79 (82.3)	4 (100.0)	16 (100.0)	9 (81.8)	6 (85.7)	26 (81.2)	35 (92.1)		

Table 1: Summarization of clinical variables stratified by subtype (continued)

	level	NEMO1	NEMO2	NEMO3	NEMO4	NEMO5	NEMO6	NEMO7	NEMO8	NEMO9	p	test
primary lymph node presentation assessment (%)		55 (17.9)	25 (22.1)	25 (26.0)	2 (50.0)	3 (18.8)	1 (9.1)	1 (14.3)	4 (12.5)	9 (23.7)	0.262	exact
	NO	9 (2.9)	4 (3.5)	5 (5.2)	0 (0.0)	0 (0.0)	1 (9.1)	0 (0.0)	3 (9.4)	4 (10.5)		
histological type (%)	YES	244 (79.2)	84 (74.3)	66 (68.8)	2 (50.0)	13 (81.2)	9 (81.8)	6 (85.7)	25 (78.1)	25 (65.8)	<0.001	exact
		0 (0.0)	1 (0.9)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	Infiltrating Carcinoma NOS	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (9.1)	0 (0.0)	0 (0.0)	0 (0.0)		
	Infiltrating Ductal Carcinoma	175 (56.8)	91 (80.5)	81 (84.4)	4 (100.0)	11 (68.8)	7 (63.6)	6 (85.7)	30 (93.8)	31 (81.6)		
	Infiltrating Lobular Carcinoma	100 (32.5)	2 (1.8)	6 (6.2)	0 (0.0)	0 (0.0)	2 (18.2)	1 (14.3)	1 (3.1)	6 (15.8)		
	Medullary Carcinoma	0 (0.0)	4 (3.5)	1 (1.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	Metaplastic Carcinoma	1 (0.3)	2 (1.8)	0 (0.0)	0 (0.0)	4 (25.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	Mixed Histology (please specify)	16 (5.2)	3 (2.7)	2 (2.1)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	Mucinous Carcinoma	5 (1.6)	0 (0.0)	3 (3.1)	0 (0.0)	0 (0.0)	1 (9.1)	0 (0.0)	0 (0.0)	1 (2.6)		
	Other, specify	11 (3.6)	10 (8.8)	3 (3.1)	0 (0.0)	1 (6.2)	0 (0.0)	0 (0.0)	1 (3.1)	0 (0.0)		
menopause status (%)		24 (7.8)	12 (10.6)	5 (5.2)	1 (25.0)	1 (6.2)	0 (0.0)	0 (0.0)	2 (6.2)	6 (15.8)	0.583	exact
	Indeterminate (neither Pre or Post-menopausal)	7 (2.3)	5 (4.4)	1 (1.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (2.6)		
	Peri (6-12 months since last menstrual period)	9 (2.9)	5 (4.4)	4 (4.2)	0 (0.0)	0 (0.0)	1 (9.1)	0 (0.0)	0 (0.0)	1 (2.6)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	193 (62.7)	71 (62.8)	60 (62.5)	2 (50.0)	13 (81.2)	7 (63.6)	6 (85.7)	26 (81.2)	27 (71.1)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	75 (24.4)	20 (17.7)	26 (27.1)	1 (25.0)	2 (12.5)	3 (27.3)	1 (14.3)	4 (12.5)	3 (7.9)		
breast carcinoma progesterone receptor status (%)		16 (5.2)	6 (5.3)	12 (12.5)	0 (0.0)	1 (6.2)	0 (0.0)	1 (14.3)	3 (9.4)	0 (0.0)	<0.001	exact
	Indeterminate	1 (0.3)	1 (0.9)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	Negative	27 (8.8)	87 (77.0)	16 (16.7)	1 (25.0)	9 (56.2)	7 (63.6)	3 (42.9)	19 (59.4)	15 (39.5)		
breast carcinoma estrogen receptor status (%)	Positive	264 (85.7)	19 (16.8)	68 (70.8)	3 (75.0)	6 (37.5)	4 (36.4)	3 (42.9)	10 (31.2)	23 (60.5)	<0.001	exact
		16 (5.2)	5 (4.4)	12 (12.5)	0 (0.0)	1 (6.2)	0 (0.0)	1 (14.3)	3 (9.4)	0 (0.0)		
	Negative	10 (3.2)	80 (70.8)	4 (4.2)	0 (0.0)	6 (37.5)	3 (27.3)	0 (0.0)	17 (53.1)	10 (26.3)		
	Positive	282 (91.6)	28 (24.8)	80 (83.3)	4 (100.0)	9 (56.2)	8 (72.7)	6 (85.7)	12 (37.5)	28 (73.7)		

Table 1: Summarization of clinical variables stratified by subtype (continued)

	level	NEMO1	NEMO2	NEMO3	NEMO4	NEMO5	NEMO6	NEMO7	NEMO8	NEMO9	p	test
lab proc her2 neu immuno- histochem- istry receptor status (%)		60 (19.5)	22 (19.5)	21 (21.9)	0 (0.0)	3 (18.8)	2 (18.2)	2 (28.6)	5 (15.6)	3 (7.9)	<0.001	exact
	Equivocal	58 (18.8)	19 (16.8)	26 (27.1)	1 (25.0)	0 (0.0)	2 (18.2)	0 (0.0)	3 (9.4)	1 (2.6)		
	Indeterminate	3 (1.0)	1 (0.9)	4 (4.2)	1 (25.0)	1 (6.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	Negative	157 (51.0)	58 (51.3)	40 (41.7)	2 (50.0)	10 (62.5)	7 (63.6)	5 (71.4)	16 (50.0)	21 (55.3)		
	Positive	30 (9.7)	13 (11.5)	5 (5.2)	0 (0.0)	2 (12.5)	0 (0.0)	0 (0.0)	8 (25.0)	13 (34.2)		
number of lymphn- odes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	1.00 [0.50, 1.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.25]	0.061	nonnorm
number of lymphn- odes positive by he (median [IQR])		1.00 [0.00, 3.00]	0.00 [0.00, 2.00]	1.00 [0.00, 4.00]	1.00 [0.00, 2.25]	0.00 [0.00, 1.00]	0.00 [0.00, 4.00]	0.00 [0.00, 1.00]	2.00 [0.00, 5.00]	1.00 [0.00, 3.00]	0.036	nonnorm
er level cell percent- age category (%)		148 (48.1)	80 (70.8)	50 (52.1)	0 (0.0)	10 (62.5)	5 (45.5)	2 (28.6)	20 (62.5)	18 (47.4)	<0.001	exact
	<10%	5 (1.6)	20 (17.7)	0 (0.0)	0 (0.0)	1 (6.2)	0 (0.0)	0 (0.0)	3 (9.4)	6 (15.8)		
	10-19%	8 (2.6)	1 (0.9)	8 (8.3)	0 (0.0)	1 (6.2)	0 (0.0)	0 (0.0)	1 (3.1)	1 (2.6)		
	20-29%	1 (0.3)	2 (1.8)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (2.6)		
	30-39%	3 (1.0)	1 (0.9)	0 (0.0)	0 (0.0)	0 (0.0)	1 (9.1)	0 (0.0)	0 (0.0)	0 (0.0)		
	40-49%	1 (0.3)	0 (0.0)	0 (0.0)	0 (0.0)	1 (6.2)	0 (0.0)	0 (0.0)	1 (3.1)	0 (0.0)		
	50-59%	5 (1.6)	1 (0.9)	1 (1.0)	0 (0.0)	0 (0.0)	1 (9.1)	0 (0.0)	0 (0.0)	0 (0.0)		
	60-69%	5 (1.6)	0 (0.0)	2 (2.1)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	70-79%	25 (8.1)	3 (2.7)	3 (3.1)	1 (25.0)	0 (0.0)	2 (18.2)	1 (14.3)	2 (6.2)	3 (7.9)		
	80-89%	11 (3.6)	0 (0.0)	6 (6.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (2.6)		
	90-99%	96 (31.2)	5 (4.4)	26 (27.1)	3 (75.0)	3 (18.8)	2 (18.2)	4 (57.1)	5 (15.6)	8 (21.1)		
progesterone receptor level cell percent category (%)		155 (50.3)	87 (77.0)	58 (60.4)	0 (0.0)	13 (81.2)	6 (54.5)	2 (28.6)	20 (62.5)	21 (55.3)	<0.001	exact
	<10%	23 (7.5)	22 (19.5)	12 (12.5)	1 (25.0)	1 (6.2)	3 (27.3)	3 (42.9)	6 (18.8)	8 (21.1)		
	10-19%	11 (3.6)	0 (0.0)	5 (5.2)	2 (50.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	20-29%	7 (2.3)	0 (0.0)	1 (1.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	30-39%	6 (1.9)	1 (0.9)	1 (1.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	40-49%	5 (1.6)	0 (0.0)	4 (4.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	2 (5.3)		
	50-59%	6 (1.9)	0 (0.0)	2 (2.1)	0 (0.0)	1 (6.2)	1 (9.1)	0 (0.0)	0 (0.0)	0 (0.0)		
	60-69%	7 (2.3)	1 (0.9)	2 (2.1)	0 (0.0)	1 (6.2)	0 (0.0)	1 (14.3)	0 (0.0)	0 (0.0)		
	70-79%	22 (7.1)	1 (0.9)	2 (2.1)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	2 (6.2)	2 (5.3)		
	80-89%	21 (6.8)	0 (0.0)	2 (2.1)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	90-99%	45 (14.6)	1 (0.9)	7 (7.3)	1 (25.0)	0 (0.0)	1 (9.1)	1 (14.3)	4 (12.5)	5 (13.2)		
distant metastasis present ind2 (%)		175 (56.8)	77 (68.1)	56 (58.3)	3 (75.0)	10 (62.5)	7 (63.6)	1 (14.3)	17 (53.1)	20 (52.6)	0.247	exact
	NO	130 (42.2)	35 (31.0)	40 (41.7)	1 (25.0)	6 (37.5)	4 (36.4)	6 (85.7)	14 (43.8)	17 (44.7)		
	YES	3 (1.0)	1 (0.9)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (3.1)	1 (2.6)		
stage event pathologic stage (%)		2 (0.6)	1 (0.9)	1 (1.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (2.6)	0.146	exact
	Stage I	30 (9.7)	6 (5.3)	5 (5.2)	0 (0.0)	2 (12.5)	1 (9.1)	0 (0.0)	2 (6.2)	4 (10.5)		
	Stage IA	25 (8.1)	10 (8.8)	8 (8.3)	1 (25.0)	0 (0.0)	2 (18.2)	1 (14.3)	2 (6.2)	0 (0.0)		
	Stage IB	3 (1.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	Stage II	3 (1.0)	1 (0.9)	0 (0.0)	0 (0.0)	1 (6.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	Stage IIA	87 (28.2)	54 (47.8)	28 (29.2)	1 (25.0)	3 (18.8)	3 (27.3)	2 (28.6)	12 (37.5)	12 (31.6)		
	Stage IIB	77 (25.0)	20 (17.7)	24 (25.0)	0 (0.0)	5 (31.2)	2 (18.2)	3 (42.9)	7 (21.9)	10 (26.3)		
	Stage III	2 (0.6)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	Stage IIIA	47 (15.3)	11 (9.7)	22 (22.9)	2 (50.0)	4 (25.0)	0 (0.0)	1 (14.3)	8 (25.0)	5 (13.2)		
	Stage IIIB	4 (1.3)	3 (2.7)	2 (2.1)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (3.1)	4 (10.5)		
	Stage IIIC	24 (7.8)	4 (3.5)	4 (4.2)	0 (0.0)	0 (0.0)	2 (18.2)	0 (0.0)	0 (0.0)	2 (5.3)		
	Stage IV	2 (0.6)	2 (1.8)	2 (2.1)	0 (0.0)	1 (6.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)		
	Stage X	2 (0.6)	1 (0.9)	0 (0.0)	0 (0.0)	0 (0.0)	1 (9.1)	0 (0.0)	0 (0.0)	0 (0.0)		