

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	iClusterBayes1	iClusterBayes2	iClusterBayes3	iClusterBayes4
n		64	306	61	69
Subtype (%)	iClusterBayes1	64 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)
	iClusterBayes2	0 (0.0)	306 (100.0)	0 (0.0)	0 (0.0)
	iClusterBayes3	0 (0.0)	0 (0.0)	61 (100.0)	0 (0.0)
	iClusterBayes4	0 (0.0)	0 (0.0)	0 (0.0)	69 (100.0)
	iClusterBayes5	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
vital status (%)	Alive	16 (94.1)	86 (94.5)	18 (100.0)	16 (94.1)
	Dead	1 (5.9)	5 (5.5)	0 (0.0)	1 (5.9)
days to birth (median [IQR])		-20249.00 [-23612.00, -17743.50]	-21291.00 [-24301.75, -17607.00]	-21608.00 [-25247.50, -18678.00]	-20048.00 [-23612.00, -18084.50]
days to death (median [IQR])		570.00 [372.50, 742.75]	1590.50 [575.50, 2397.00]	723.00 [419.50, 2232.50]	1759.00 [914.50, 2965.00]
days to last followup (median [IQR])		722.00 [437.50, 1559.00]	862.50 [503.50, 1689.00]	765.00 [528.00, 1224.00]	767.00 [437.50, 1714.25]
race list (%)	AMERICAN INDIAN OR ALASKA NATIVE	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
	ASIAN	4 (6.5)	12 (4.0)	3 (5.1)	3 (4.3)
	BLACK OR AFRICAN AMERICAN	20 (32.3)	44 (14.7)	11 (18.6)	27 (39.3)
	WHITE	38 (61.3)	243 (81.3)	45 (76.3)	39 (56.4)
history of neoadjuvant treatment (%)	No	63 (100.0)	305 (100.0)	60 (98.4)	69 (100.0)
	Yes	0 (0.0)	0 (0.0)	1 (1.6)	0 (0.0)
age at initial pathologic diagnosis (median [IQR])		55.00 [48.00, 64.25]	58.00 [47.25, 66.00]	59.00 [50.00, 69.00]	54.00 [49.00, 64.25]
ethnicity (%)	HISPANIC OR LATINO	2 (3.3)	15 (5.5)	0 (0.0)	4 (6.3)
	NOT HISPANIC OR LATINO	58 (96.7)	260 (94.5)	57 (100.0)	59 (93.7)
primary lymph node presentation assessment (%)	NO	3 (5.6)	7 (2.9)	2 (4.0)	3 (5.3)
	YES	51 (94.4)	234 (97.1)	48 (96.0)	54 (94.7)

Table 1: Summarization of clinical variables stratified by subtype (continued)

	level	iClusterBayes1	iClusterBayes2	iClusterBayes3	iClusterBayes4
histological type (%)	Infiltrating Carcinoma NOS	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.5)
	Infiltrating Ductal Carcinoma	46 (71.9)	194 (63.4)	39 (63.9)	55 (80.0)
	Infiltrating Lobular Carcinoma	9 (14.1)	79 (25.8)	13 (21.3)	1 (1.5)
	Medullary Carcinoma	0 (0.0)	0 (0.0)	0 (0.0)	5 (7.4)
	Metaplastic Carcinoma	3 (4.7)	0 (0.0)	1 (1.6)	3 (4.4)
	Mixed Histology (please specify)	0 (0.0)	11 (3.6)	6 (9.8)	0 (0.0)
	Mucinous Carcinoma	0 (0.0)	9 (2.9)	1 (1.6)	0 (0.0)
	Other, specify	6 (9.4)	13 (4.2)	1 (1.6)	3 (4.4)
menopause status (%)	Indeterminate (neither Pre or Postmenopausal)	2 (3.2)	8 (2.8)	1 (1.7)	2 (3.3)
	Peri (6-12 months since last menstrual period)	2 (3.2)	9 (3.2)	0 (0.0)	6 (9.8)
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	48 (77.4)	187 (66.5)	46 (79.3)	43 (70.0)
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	10 (16.1)	77 (27.4)	11 (19.0)	10 (16.7)
	Indeterminate	1 (1.7)	1 (0.3)	0 (0.0)	0 (0.0)
breast carcinoma progesterone receptor status (%)	Negative	46 (78.0)	41 (14.0)	17 (29.3)	59 (93.3)
	Positive	12 (20.3)	251 (85.7)	41 (70.7)	4 (6.7)
breast carcinoma estrogen receptor status (%)	Negative	44 (73.3)	10 (3.4)	8 (13.8)	53 (84.0)
	Positive	16 (26.7)	283 (96.6)	50 (86.2)	10 (16.0)
lab proc her2 neu immuno-histochemistry receptor status (%)	Equivocal	8 (16.0)	53 (21.2)	22 (44.0)	14 (25.0)
	Indeterminate	0 (0.0)	4 (1.6)	2 (4.0)	2 (3.7)
	Negative	36 (72.0)	169 (67.6)	19 (38.0)	36 (66.7)
	Positive	6 (12.0)	24 (9.6)	7 (14.0)	2 (3.7)

Table 1: Summarization of clinical variables stratified by subtype (continued)

	level	iClusterBayes1	iClusterBayes2	iClusterBayes3	iClusterBayes4
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.00 [0.00, 0.00]
number of lymphnodes positive by he (median [IQR])		0.00 [0.00, 2.00]	1.00 [0.00, 3.00]	0.00 [0.00, 4.00]	0.00 [0.00, 2.00]
er level cell percentage category (%)	<10%	10 (45.5)	6 (3.6)	1 (3.7)	13 (76.2)
	10-19%	2 (9.1)	8 (4.7)	4 (14.8)	1 (5.9)
	20-29%	0 (0.0)	0 (0.0)	1 (3.7)	2 (11.1)
	30-39%	0 (0.0)	2 (1.2)	1 (3.7)	1 (5.9)
	40-49%	0 (0.0)	2 (1.2)	0 (0.0)	0 (0.0)
	50-59%	0 (0.0)	6 (3.6)	0 (0.0)	0 (0.0)
	60-69%	1 (4.5)	4 (2.4)	1 (3.7)	0 (0.0)
	70-79%	3 (13.6)	25 (14.8)	3 (11.1)	0 (0.0)
	80-89%	0 (0.0)	10 (5.9)	1 (3.7)	0 (0.0)
	90-99%	6 (27.3)	106 (62.7)	15 (55.6)	0 (0.0)
progesterone receptor level cell percent category (%)	<10%	12 (63.2)	31 (19.4)	7 (31.8)	15 (100.0)
	10-19%	0 (0.0)	11 (6.9)	4 (18.2)	0 (0.0)
	20-29%	1 (5.3)	3 (1.9)	2 (9.1)	0 (0.0)
	30-39%	0 (0.0)	6 (3.8)	1 (4.5)	0 (0.0)
	40-49%	0 (0.0)	8 (5.0)	0 (0.0)	0 (0.0)
	50-59%	0 (0.0)	8 (5.0)	0 (0.0)	0 (0.0)
	60-69%	0 (0.0)	9 (5.6)	0 (0.0)	0 (0.0)
	70-79%	2 (10.5)	21 (13.1)	3 (13.6)	0 (0.0)
	80-89%	1 (5.3)	14 (8.8)	1 (4.5)	0 (0.0)
	90-99%	3 (15.8)	49 (30.6)	4 (18.2)	0 (0.0)
distant metastasis present ind2 (%)	NO	18 (85.7)	132 (97.8)	24 (100.0)	24 (100.0)
	YES	3 (14.3)	3 (2.2)	0 (0.0)	0 (0.0)
stage event pathologic stage (%)	Stage I	5 (7.8)	28 (9.2)	2 (3.3)	6 (8.8)
	Stage IA	5 (7.8)	31 (10.2)	6 (10.0)	4 (5.9)
	Stage IB	0 (0.0)	3 (1.0)	0 (0.0)	0 (0.0)
	Stage II	0 (0.0)	3 (1.0)	0 (0.0)	1 (1.5)
	Stage IIA	28 (43.8)	81 (26.7)	20 (33.3)	33 (48.1)
	Stage IIB	15 (23.4)	79 (26.1)	11 (18.3)	14 (20.6)
	Stage III	1 (1.6)	0 (0.0)	0 (0.0)	0 (0.0)
	Stage IIIA	6 (9.4)	52 (17.2)	11 (18.3)	5 (7.4)
	Stage IIIB	0 (0.0)	6 (2.0)	2 (3.3)	2 (2.9)
	Stage IIIC	3 (4.7)	15 (5.0)	8 (13.3)	1 (1.5)
	Stage IV	1 (1.6)	3 (1.0)	0 (0.0)	1 (1.5)

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

level	iClusterBayes1	iClusterBayes2	iClusterBayes3	iClusterBayes4
Stage X	0 (0.0)	2 (0.7)	0 (0.0)	1 (1.5)