

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	CIMLR1	CIMLR2	p	test
n		500	125		
Subtype (%)	CIMLR1	500 (100.0)	0 (0.0)	<0.001	
	CIMLR2	0 (0.0)	125 (100.0)		
vital status (%)	Alive	137 (92.6)	26 (100.0)	0.374	exact
	Dead	11 (7.4)	0 (0.0)		
days to birth (median [IQR])		-20716.00 [-24135.50, -17590.25]	-22688.00 [-25957.50, -19591.25]	0.001	nonnorm
days to death (median [IQR])		1365.00 [577.00, 2417.00]	1688.00 [343.50, 2333.00]	0.701	nonnorm
days to last followup (median [IQR])		816.00 [508.25, 1633.25]	602.00 [413.25, 1545.00]	0.043	nonnorm
race list (%)	AMERICAN INDIAN OR ALASKA NATIVE	1 (0.2)	0 (0.0)	0.143	exact
	ASIAN	22 (4.5)	10 (8.1)		
	BLACK OR AFRICAN AMERICAN	94 (19.3)	30 (24.2)		
	WHITE	371 (76.0)	84 (67.7)		
history of neoadjuvant treatment (%)	No	494 (99.2)	123 (98.4)	0.346	exact
	Yes	4 (0.8)	2 (1.6)		
age at initial pathologic diagnosis (median [IQR])		56.00 [47.75, 65.25]	61.00 [53.00, 71.00]	0.001	nonnorm
ethnicity (%)	HISPANIC OR LATINO	25 (5.5)	2 (1.8)	0.152	
	NOT HISPANIC OR LATINO	426 (94.5)	111 (98.2)		
primary lymph node presentation assessment (%)	NO	16 (4.0)	10 (10.1)	0.028	
	YES	385 (96.0)	89 (89.9)		
histological type (%)	Infiltrating Carcinoma NOS	0 (0.0)	1 (0.8)	0.136	exact
	Infiltrating Ductal Carcinoma	347 (69.4)	89 (71.8)		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	CIMLR1	CIMLR2	p	test
menopause status (%)	Infiltrating Lobular Carcinoma	93 (18.6)	25 (20.2)	0.022	exact
	Medullary Carcinoma	3 (0.6)	2 (1.6)		
	Metaplastic Carcinoma	5 (1.0)	2 (1.6)		
	Mixed Histology (please specify)	20 (4.0)	1 (0.8)		
	Mucinous Carcinoma	9 (1.8)	1 (0.8)		
	Other, specify	23 (4.6)	3 (2.4)		
	Indeterminate (neither Pre or Postmenopausal)	11 (2.4)	3 (2.6)		
	Peri (6-12 months since last menstrual period)	16 (3.5)	4 (3.5)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	312 (68.0)	93 (80.9)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	120 (26.1)	15 (13.0)		
breast carcinoma progesterone receptor status (%)	Indeterminate	2 (0.4)	0 (0.0)	<0.001	exact
breast carcinoma estrogen receptor status (%)	Negative	123 (26.3)	61 (51.3)	<0.001	
	Positive	342 (73.2)	58 (48.7)		
	Negative	83 (17.7)	47 (39.5)		
lab proc her2 neu immuno-histochemistry receptor status (%)	Positive	385 (82.3)	72 (60.5)	0.038	exact
	Equivocal	95 (23.6)	15 (14.3)		
	Indeterminate	6 (1.5)	4 (3.8)		
	Negative	250 (62.2)	66 (62.9)	0.250	nonnorm
	Positive	51 (12.7)	20 (19.0)		
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]		

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	CIMLR1	CIMLR2	p	test
number of lymph nodes positive by HER2 (median [IQR])		1.00 [0.00, 3.00]	0.00 [0.00, 1.00]	<0.001	nonnorm
HER2 level cell percentage category (%)	<10%	19 (8.1)	16 (28.1)	0.015	exact
	10-19%	18 (7.7)	2 (3.5)		
	20-29%	2 (0.9)	2 (3.5)		
	30-39%	4 (1.7)	1 (1.8)		
	40-49%	3 (1.3)	0 (0.0)		
	50-59%	7 (3.0)	1 (1.8)		
	60-69%	5 (2.1)	2 (3.5)		
	70-79%	33 (14.0)	7 (12.3)		
	80-89%	14 (6.0)	4 (7.0)		
	90-99%	130 (55.3)	22 (38.6)		
progesterone receptor level cell percent category (%)	<10%	52 (24.8)	27 (50.9)	0.039	exact
	10-19%	16 (7.6)	2 (3.8)		
	20-29%	8 (3.8)	0 (0.0)		
	30-39%	8 (3.8)	0 (0.0)		
	40-49%	8 (3.8)	3 (5.7)		
	50-59%	8 (3.8)	2 (3.8)		
	60-69%	10 (4.8)	2 (3.8)		
	70-79%	25 (11.9)	4 (7.5)		
	80-89%	18 (8.6)	5 (9.4)		
	90-99%	57 (27.1)	8 (15.1)		
distant metastasis present in d2 (%)	NO	207 (97.6)	46 (97.9)	1.000	exact
	YES	5 (2.4)	1 (2.1)		
stage event pathologic stage (%)	Stage I	40 (8.1)	10 (8.1)	0.225	exact
	Stage IA	42 (8.5)	7 (5.6)		
	Stage IB	3 (0.6)	0 (0.0)		
	Stage II	5 (1.0)	0 (0.0)		
	Stage IIA	149 (30.0)	53 (42.7)		
	Stage IIB	121 (24.4)	27 (21.8)		
	Stage III	2 (0.4)	0 (0.0)		
	Stage IIIA	83 (16.7)	17 (13.7)		
	Stage IIIB	9 (1.8)	5 (4.0)		
	Stage IIIC	33 (6.7)	3 (2.4)		
	Stage IV	6 (1.2)	1 (0.8)		
	Stage X	3 (0.6)	1 (0.8)		