

Summary_of_clinical_variables

Contents

Table 1: Summarization of clinical variables stratified by subtype

	level	CS1	CS2	p	test
n		244	381		
Subtype (%)	CS1	244 (100.0)	0 (0.0)	<0.001	
	CS2	0 (0.0)	381 (100.0)		
vital status (%)	Alive	56 (91.8)	107 (94.7)	0.520	exact
	Dead	5 (8.2)	6 (5.3)		
days to birth (median [IQR])		-21608.00 [-25029.00, -18346.00]	-21056.00 [-24219.00, -17683.00]	0.275	nonnorm
days to death (median [IQR])		1694.00 [639.00, 2798.00]	912.00 [564.50, 2012.00]	0.192	nonnorm
days to last followup (median [IQR])		661.00 [439.00, 1417.00]	860.00 [492.00, 1742.00]	0.025	nonnorm
race list (%)	AMERICAN INDIAN OR ALASKA NATIVE	1 (0.4)	0 (0.0)	0.106	exact
	ASIAN	16 (6.7)	16 (4.3)		
	BLACK OR AFRICAN AMERICAN	55 (22.9)	69 (18.5)		
	WHITE	168 (70.0)	287 (77.2)		
history of neoadjuvant treatment (%)	No	238 (97.5)	379 (100.0)	0.003	exact
	Yes	6 (2.5)	0 (0.0)		
age at initial pathologic diagnosis (median [IQR])		58.50 [49.75, 68.00]	57.00 [48.00, 65.00]	0.306	nonnorm
ethnicity (%)	HISPANIC OR LATINO	9 (4.1)	18 (5.2)	0.705	
	NOT HISPANIC OR LATINO	209 (95.9)	328 (94.8)		
primary lymph node presentation assessment (%)	NO	15 (7.7)	11 (3.6)	0.072	
	YES	180 (92.3)	294 (96.4)		
histological type (%)	Infiltrating Carcinoma	0 (0.0)	1 (0.3)	<0.001	exact
	NOS	189 (77.8)	247 (64.8)		
	Infiltrating Ductal Carcinoma				

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	CS1	CS2	p	test
menopause status (%)	Infiltrating Lobular Carcinoma	28 (11.5)	90 (23.6)	0.623	
	Medullary Carcinoma	5 (2.1)	0 (0.0)		
	Metaplastic Carcinoma	3 (1.2)	4 (1.0)		
	Mixed Histology (please specify)	10 (4.1)	11 (2.9)		
	Mucinous Carcinoma	1 (0.4)	9 (2.4)		
	Other, specify	7 (2.9)	19 (5.0)		
	Indeterminate (neither Pre or Postmenopausal)	4 (1.8)	10 (2.8)		
	Peri (6-12 months since last menstrual period)	8 (3.6)	12 (3.4)		
	Post (prior bilateral ovariectomy OR >12 mo since LMP with no prior hysterectomy)	162 (73.3)	243 (68.8)		
	Pre (<6 months since LMP AND no prior bilateral ovariectomy AND not on estrogen replacement)	47 (21.3)	88 (24.9)		
breast carcinoma progesterone receptor status (%)	Indeterminate	0 (0.0)	2 (0.6)	0.001	exact
breast carcinoma estrogen receptor status (%)	Negative	90 (40.4)	94 (25.9)	<0.001	
	Positive	133 (59.6)	267 (73.6)		
	Negative	68 (30.5)	62 (17.0)		
lab proc her2 neu immuno-histochemistry receptor status (%)	Positive	155 (69.5)	302 (83.0)	<0.001	exact
	Equivocal	46 (23.5)	64 (20.6)		
	Indeterminate	5 (2.6)	5 (1.6)		
	Negative	104 (53.1)	212 (68.2)		
	Positive	41 (20.9)	30 (9.6)		
number of lymphnodes positive by ihc (median [IQR])		0.00 [0.00, 0.00]	0.00 [0.00, 0.00]	0.200	nonnorm

Table 1: Summarization of clinical variables stratified by subtype (*continued*)

	level	CS1	CS2	p	test
number of lymphnodes positive by he (median [IQR])		1.00 [0.00, 3.25]	1.00 [0.00, 2.00]	0.576	nonnorm
er level cell percentage category (%)	<10%	18 (18.2)	17 (8.8)	0.006	exact
	10-19%	10 (10.1)	10 (5.2)		
	20-29%	4 (4.0)	0 (0.0)		
	30-39%	3 (3.0)	2 (1.0)		
	40-49%	1 (1.0)	2 (1.0)		
	50-59%	1 (1.0)	7 (3.6)		
	60-69%	2 (2.0)	5 (2.6)		
	70-79%	12 (12.1)	28 (14.5)		
	80-89%	8 (8.1)	10 (5.2)		
	90-99%	40 (40.4)	112 (58.0)		
progesterone receptor level cell percent category (%)	<10%	34 (42.0)	45 (24.7)	0.084	exact
	10-19%	7 (8.6)	11 (6.0)		
	20-29%	4 (4.9)	4 (2.2)		
	30-39%	2 (2.5)	6 (3.3)		
	40-49%	3 (3.7)	8 (4.4)		
	50-59%	1 (1.2)	9 (4.9)		
	60-69%	3 (3.7)	9 (4.9)		
	70-79%	6 (7.4)	23 (12.6)		
	80-89%	8 (9.9)	15 (8.2)		
	90-99%	13 (16.0)	52 (28.6)		
distant metastasis present ind2 (%)	NO	99 (100.0)	154 (96.2)	0.085	exact
	YES	0 (0.0)	6 (3.8)		
stage event pathologic stage (%)	Stage I	15 (6.2)	35 (9.3)	0.039	exact
	Stage IA	10 (4.1)	39 (10.3)		
	Stage IB	0 (0.0)	3 (0.8)		
	Stage II	2 (0.8)	3 (0.8)		
	Stage IIA	90 (37.2)	112 (29.6)		
	Stage IIB	52 (21.5)	96 (25.4)		
	Stage III	1 (0.4)	1 (0.3)		
	Stage IIIA	42 (17.4)	58 (15.3)		
	Stage IIIB	8 (3.3)	6 (1.6)		
	Stage IIIC	18 (7.4)	18 (4.8)		
	Stage IV	3 (1.2)	4 (1.1)		
	Stage X	1 (0.4)	3 (0.8)		