

Seasonality Detection Methods: A Comparative Study

fdars Package

today

Contents

Introduction	1
Detection Methods	2
AIC Comparison (Fourier vs P-spline)	2
FFT Confidence	2
ACF Confidence	2
Variance Strength	3
Spectral Strength	3
Automatic Basis Selection	3
Simulation Studies	3
Simulation 1: Varying Seasonal Strength	3
Setup	3
Code	4
Configuration	4
Detection thresholds (calibrated to ~5% FPR on pure noise)	4
Generate seasonal curve	4
Results	6
F1 Score vs Trend Strength	6
False Positive Rate by Trend Strength	7
Simulation 3: Multiple Trend Types	7
Setup	7
Code	7
Threshold Guidelines	9
Conclusion	9
Appendix: File Listing	10

Introduction

This report describes and compares six methods for detecting seasonality in functional time series data. We evaluate each method's performance across different scenarios including varying seasonal strengths, non-linear trends, and different trend types.

The goal is to answer: **Given a time series, how can we reliably determine if it contains a seasonal pattern?**

Detection Methods

AIC Comparison (Fourier vs P-spline)

Concept: If data is seasonal, a Fourier basis should fit better than P-splines because Fourier bases naturally capture periodic patterns.

Mathematical formulation:

For a curve $y(t)$, we fit two models:

1. **Fourier basis:** $\hat{y}(t) = \sum_{k=0}^K a_k \cos(2\pi kt) + b_k \sin(2\pi kt)$
2. **P-spline:** $\hat{y}(t) = \sum_{j=1}^J c_j B_j(t)$ with penalty $\lambda \int [\hat{y}''(t)]^2 dt$

We compute AIC for each:

$$\text{AIC} = n \log(\text{RSS}/n) + 2 \cdot \text{edf}$$

where RSS is the residual sum of squares and edf is the effective degrees of freedom.

Detection rule: Seasonality detected if $\text{AIC}_{\text{P-spline}} - \text{AIC}_{\text{Fourier}} > 0$

Interpretation: When Fourier has lower AIC, the periodic structure is significant enough to justify the global periodic assumption over the local flexibility of splines.

FFT Confidence

Concept: Use Fast Fourier Transform to detect dominant frequencies. Strong peaks in the periodogram indicate periodic components.

Mathematical formulation:

Given a time series y_1, y_2, \dots, y_n , compute the discrete Fourier transform:

$$Y_k = \sum_{j=1}^n y_j e^{-2\pi i(j-1)(k-1)/n}$$

The periodogram (power spectrum) is:

$$P_k = |Y_k|^2$$

Detection score:

$$\text{Confidence} = \frac{\max_k P_k}{\text{mean}(P_k)}$$

Detection rule: Seasonality detected if Confidence > 6.0

Interpretation: A high ratio indicates one frequency dominates, suggesting periodicity rather than random noise.

ACF Confidence

Concept: Autocorrelation at the seasonal lag should be high for seasonal data.

Mathematical formulation:

The autocorrelation function at lag h is:

$$\rho_h = \frac{\sum_{t=1}^{n-h} (y_t - \bar{y})(y_{t+h} - \bar{y})}{\sum_{t=1}^n (y_t - \bar{y})^2}$$

For seasonal data with period p , we expect ρ_p to be significantly positive.

Detection rule: Seasonality detected if ACF confidence > 0.25

Interpretation: High autocorrelation at the seasonal lag indicates the pattern repeats.

Variance Strength

Concept: Decompose variance into seasonal and residual components. High seasonal variance ratio indicates seasonality.

Mathematical formulation:

Decompose the series: $y_t = T_t + S_t + R_t$ (trend + seasonal + residual)

The seasonal strength is:

$$SS_{\text{var}} = 1 - \frac{\text{Var}(R_t)}{\text{Var}(y_t - T_t)}$$

Detection rule: Seasonality detected if $SS_{\text{var}} > 0.2$

Interpretation: Values close to 1 mean the seasonal component dominates; values close to 0 mean residual noise dominates.

Important: The `period` parameter must be in the same units as `argvals`. For data normalized to $[0,1]$ with 5 annual cycles, use `period = 0.2`.

Spectral Strength

Concept: Measure the proportion of spectral power at the seasonal frequency.

Mathematical formulation:

Using the periodogram P_k , identify the seasonal frequency $f_s = 1/\text{period}$.

$$SS_{\text{spectral}} = \frac{\sum_{k \in \mathcal{S}} P_k}{\sum_k P_k}$$

where \mathcal{S} includes the seasonal frequency and its harmonics.

Detection rule: Seasonality detected if $SS_{\text{spectral}} > 0.3$

Interpretation: High values indicate spectral energy is concentrated at seasonal frequencies.

Automatic Basis Selection

Concept: Let the model selection process decide—if Fourier basis is selected over P-splines, the data is likely seasonal.

Method: Uses `select.basis.auto()` with AIC criterion.

Note: The internal FFT-based seasonal hint has a threshold that is too low (2.0 instead of ~6.0), causing high false positive rates. Use `use.seasonal.hint = FALSE`.

Simulation Studies

Simulation 1: Varying Seasonal Strength

Setup

This simulation tests how well each method detects seasonality at different signal strengths.

Parameters:

- 11 seasonal strength levels: 0.0, 0.1, ..., 1.0
- 50 curves per strength level
- 5 years of monthly data (60 observations)
- Noise standard deviation: 0.3

Signal model:

$$y(t) = s \cdot [\sin(2\pi \cdot 5t) + 0.3 \cos(4\pi \cdot 5t)] + \epsilon, \quad \epsilon \sim N(0, 0.3^2)$$

where s is the seasonal strength (0 = no seasonality, 1 = full seasonality).

Ground truth: A curve is classified as “truly seasonal” if $s \geq 0.2$.

Code

```
“{r} #| eval: false #| code-fold: true #| code-summary: “Show simulation code”  
library(fdars) library(ggplot2) library(tidyr) library(dplyr)  
set.seed(42)
```

Configuration

```
n_strengths <- 11 n_curves_per_strength <- 50 n_years <- 5 n_months <- n_years * 12 noise_sd <- 0.3
```

Detection thresholds (calibrated to ~5% FPR on pure noise)

```
detection_thresholds <- list( aic_comparison = 0, fft_confidence = 6.0, acf_confidence = 0.25,  
strength_variance = 0.2, strength_spectral = 0.3 )  
seasonal_strengths <- seq(0, 1, length.out = n_strengths) t <- seq(0, 1, length.out = n_months)
```

Generate seasonal curve

```
generate_seasonal_curve <- function(t, strength, noise_sd = 0.3) { n_cycles <- length(t) / 12 seasonal <-  
strength * sin(2 * pi * n_cycles * t) seasonal <- seasonal + strength * 0.3 * cos(4 * pi * n_cycles * t) noise  
<- rnorm(length(t), sd = noise_sd) return(seasonal + noise) }
```

Results

```
![Detection rates by seasonal strength](seasonality_detection_comparison.pdf){#fig-detection-rates width=
```

How to interpret:

- The x-axis shows the true seasonal strength (0 = pure noise, 1 = strong seasonality)
- The y-axis shows what percentage of curves each method classified as "seasonal"
- The vertical dashed line at 0.2 marks the ground truth threshold
- ****Ideal behavior****: 0% detection below the threshold, 100% above

Classification Performance

Method	F1 Score	Precision	Recall	FPR	Specificity
***Variance Strength**	97.3%	98.2%	96.4%	2.0%	92.0%

Spectral Strength	95.3%	97.4%	93.3%	10.0%	89.0%	
FFT Confidence	94.8%	99.3%	90.7%	4.0%	97.0%	
AIC Comparison	91.5%	94.3%	88.9%	18.0%	76.0%	
ACF Confidence	85.4%	98.3%	75.6%	10.0%	94.0%	

****How to interpret**:**

- ****F1 Score****: Harmonic mean of precision and recall (higher is better)
- ****Precision****: Of curves detected as seasonal, what % are truly seasonal?
- ****Recall****: Of truly seasonal curves, what % did we detect?
- ****FPR****: False Positive Rate - what % of non-seasonal curves were incorrectly flagged?

Precision-Recall Analysis

![Precision-Recall curves](seasonality_detection_details.pdf){#fig-pr-curves width=100%}

****How to interpret**:**

- Curves closer to the top-right corner are better
- The diamond markers show the operating point at the default threshold
- AUC-PR (Area Under the PR Curve) summarizes overall performance

Simulation 2: Non-linear Trend {#sec-sim2}

Setup

This simulation tests robustness when non-linear trends are added to the seasonal signal.

****Parameters**:**

- 6 seasonal strength levels × 6 trend strength levels
- 30 curves per combination
- Non-linear trend: quadratic + cubic + sigmoid components

****Signal model**:**

$y(t) = \text{Trend}(t, \tau) + \text{Seasonal}(t, s) + \epsilon$

where τ is the trend strength and s is the seasonal strength.

****Trend function**:**

$\text{Trend}(t, \tau) = \tau \cdot [2(t-0.5)^2 + 0.5(t-0.3)^3 + 0.3 \cdot \sigma(10(t-0.6)) - 0.5]$

Code

```

```{r}
#| eval: false
#| code-fold: true
#| code-summary: "Show trend simulation code"

Non-linear trend function
generate_nonlinear_trend <- function(t, trend_strength) {
 quadratic <- 2 * (t - 0.5)^2
 cubic <- 0.5 * (t - 0.3)^3
 sigmoid <- 1 / (1 + exp(-10 * (t - 0.6)))

```

```

trend <- trend_strength * (quadratic + cubic + 0.3 * sigmoid - 0.5)
return(trend)
}

Generate curve with trend + seasonal + noise
generate_curve <- function(t, seasonal_strength, trend_strength, noise_sd = 0.3) {
 trend <- generate_nonlinear_trend(t, trend_strength)
 n_cycles <- length(t) / 12
 seasonal <- seasonal_strength * sin(2 * pi * n_cycles * t)
 seasonal <- seasonal + seasonal_strength * 0.3 * cos(4 * pi * n_cycles * t)
 noise <- rnorm(length(t), sd = noise_sd)
 return(trend + seasonal + noise)
}

```

## Results




Figure 1: Detection rates heatmap by seasonal and trend strength

### How to interpret:

- Each cell shows the detection rate for a combination of seasonal strength (x) and trend strength (y)
- Blue = low detection rate, Red = high detection rate
- The dashed line separates non-seasonal (left) from seasonal (right) ground truth

### F1 Score vs Trend Strength

Method	No Trend	Max Trend	F1 Drop
Spectral	96.3%	92.5%	3.9%
FFT	93.7%	91.8%	2.0%
AIC	92.2%	87.0%	5.7%
ACF	87.4%	83.5%	4.5%

### How to interpret:

- **F1 Drop:** How much performance degrades when strong trends are present
- Lower drop = more robust to trends

### False Positive Rate by Trend Strength




Figure 2: FPR when no seasonality is present, across trend strengths

**Key finding:** FFT’s FPR increases dramatically with trend strength because non-linear trends can create spurious peaks in the periodogram.

## Simulation 3: Multiple Trend Types

### Setup

This simulation tests which types of trends cause the most problems for each detection method.

### Trend types tested:

1. **None:** Flat baseline
2. **Linear:**  $f(t) = t - 0.5$
3. **Quadratic:**  $f(t) = (t - 0.5)^2 - 0.25$
4. **Cubic:**  $f(t) = 2(t - 0.5)^3$
5. **Exponential:**  $f(t) = e^{2t}/e^2 - 0.5$
6. **Logarithmic:**  $f(t) = \log(t + 0.1)$  (normalized)
7. **Sigmoid:**  $f(t) = 1/(1 + e^{-10(t-0.5)}) - 0.5$
8. **Slow sine:**  $f(t) = \sin(2\pi t)$  — one cycle over the entire series

### Code

“{r} #| eval: false #| code-fold: true #| code-summary: “Show trend types code”

```
trend_functions <- list(none = function(t, strength) rep(0, length(t)), linear = function(t, strength) strength
* (t - 0.5), quadratic = function(t, strength) strength * ((t - 0.5)^2 - 0.25), cubic = function(t, strength)
strength * 2 * (t - 0.5)^3, exponential = function(t, strength) strength * (exp(2 * t) / exp(2) - 0.5), logarithmic
= function(t, strength) { strength * (log(t + 0.1) - log(0.1)) / (log(1.1) - log(0.1)) - 0.5 * strength }, sigmoid
= function(t, strength) strength * (1 / (1 + exp(-10 * (t - 0.5)))) - 0.5, slow_sine = function(t, strength)
strength * sin(2 * pi * t))
```

### ### Results

```
![F1 scores by trend type](seasonality_detection_trend_types_f1.pdf){#fig-trend-types-f1 width=100%}
```

### ### FPR by Trend Type

```
![Example trend functions](seasonality_detection_trend_types_examples.pdf){#fig-trend-examples width=100%}
```

```
![FPR across trend types and strengths](seasonality_detection_trend_types_fpr.pdf){#fig-trend-types-fpr width=100%}
```

**\*\*How to interpret\*\*:**

- Each panel shows one trend type
- Lines show how FPR changes as trend strength increases
- **\*\*Slow sine is catastrophic for FFT\*\***: 100% FPR because FFT detects the slow oscillation as "seasonal"

### ### Most Problematic Trend Types

Trend Type	FFT FPR	Spectral FPR	Issue
<b>**slow_sine**</b>	<b>**100%**</b>	0%	FFT detects non-seasonal oscillation
quadratic	10%	5%	Minor
sigmoid	5%	5%	Minor
linear	0%	10%	Handled well

### # Key Findings

#### ## Method Ranking

1. **\*\*Variance Strength\*\*** (F1=97.3%, FPR=2%): Best overall when period is known
2. **\*\*Spectral Strength\*\*** (F1=95.3%, FPR=10%): Most robust to different trend types
3. **\*\*FFT Confidence\*\*** (F1=94.8%, FPR=4%): Good but vulnerable to slow oscillations
4. **\*\*AIC Comparison\*\*** (F1=91.5%, FPR=18%): Interpretable but higher FPR
5. **\*\*ACF Confidence\*\*** (F1=85.4%, FPR=10%): Conservative, misses weak seasonality

#### ## Critical Issues Found

1. **\*\*Period units matter\*\***: The `period` parameter in `seasonal\_strength()` must be in argvals units, not time units
2. **\*\*FFT is vulnerable to slow oscillations\*\***: Any periodic signal (even non-seasonal) triggers detection
3. **\*\*Basis Auto has a bug\*\***: Internal FFT threshold of 2.0 is too low (should be ~6.0)

### # Recommendations

#### ## For Unknown Datasets



**\*\*Primary recommendation: Variance Strength\*\***

```
```{r}
#| eval: false

# Calculate period in argvals units
# If argvals is in [0,1] and you expect 5 annual cycles:
period_in_argvals_units <- 1 / 5 # = 0.2

strength <- seasonal_strength(fd,
                             period = period_in_argvals_units,
                             method = "variance",
                             detrend = "linear")

is_seasonal <- strength > 0.2
```

For robustness to unknown trends: Spectral Strength

```
```{r} #| eval: false

strength <- seasonal_strength(fd, period = period_in_argvals_units, method = "spectral", detrend =
"linear") is_seasonal <- strength > 0.3
```

**\*\*Ensemble approach (most robust)\*\*:**

```
```{r}
#| eval: false

var_detected <- seasonal_strength(fd, period, method = "variance") > 0.2
spec_detected <- seasonal_strength(fd, period, method = "spectral") > 0.3
fft_detected <- estimate_period(fd, method = "fft")$confidence > 6.0

# Majority vote
is_seasonal <- (var_detected + spec_detected + fft_detected) >= 2
```

Threshold Guidelines

Method	Threshold	Calibration
Variance Strength	0.2	95th percentile of noise ~0.17
Spectral Strength	0.3	95th percentile of noise ~0.29
FFT Confidence	6.0	95th percentile of noise ~5.7
ACF Confidence	0.25	95th percentile of noise ~0.22
AIC Difference	0	Fourier better → positive difference

Conclusion

For detecting seasonality in functional time series:

1. **Variance Strength** is the most accurate method when the seasonal period is known
2. **Spectral Strength** is most robust to confounding trends and unknown oscillations
3. **FFT Confidence** works well but is vulnerable to slow non-seasonal oscillations
4. **AIC Comparison** provides an interpretable alternative but has higher false positive rates
5. **ACF Confidence** is conservative (low FPR) but misses weak seasonality

The key insight is that simple variance-based decomposition outperforms more complex spectral methods when properly configured with the correct period parameter.

Appendix: File Listing

All simulation scripts and results are in `scripts/seasonal_simulation/`:

- `seasonality_detection_comparison.R` — Main comparison (Simulation 1)
- `seasonality_detection_with_trend.R` — Non-linear trend study (Simulation 2)
- `seasonality_detection_trend_types.R` — Multiple trend types (Simulation 3)
- `seasonal_basis_comparison.R` — AIC comparison study

PDF outputs contain the visualizations referenced in this report.