

found to produce a greater reduction in A1C (-0.30 percentage points; 95% CI -0.42 to -0.19) compared with control (42). Importantly, these digital methods provide outcomes that are comparable with or even better than those seen with traditional in-person care (42). Greater A1C reductions are demonstrated with increased virtual engagement, although data from trials are heterogeneous.

Diabetes care and education specialists (DCES) are effective providers of DSMES. Members of the DSMES team can include a variety of health care professionals such as nurses (registered nurses and nurse practitioners), registered dietitian nutritionists (RDNs), pharmacists, social workers, certified health education specialists, exercise physiologists, psychologists and behavioral health professionals, community health workers, care coordinators or navigators, and others who can tailor curricula to individual needs (44–46). Team members acting in a DCES capacity should have specialized clinical knowledge of diabetes and behavior change principles. In addition, a DCES needs to be knowledgeable about technology-enabled services and can serve as a technology champion within their practice (47). Credentials such as certified DCES (cbdce.org/) and/or board certification in advanced diabetes management (BC-ADM) (diabeteseducator.org/education/certification/bc_adm) demonstrates an individual's specialized training and expertise in diabetes management, education, and support (10), and engagement with qualified professionals has been shown to improve diabetes-related outcomes (48). There is also continued and growing evidence for the role of community health workers, peer educators, peer support, and lay leaders in providing ongoing diabetes self-management support (49,50). In locations where there are no DCES services available, other members of the diabetes care team can provide some aspects of DSMES; however, billing restrictions need to be considered (51).

Social determinants of health (SDOH) are an important aspect of diabetes care and should be assessed and weighed in guiding the design and delivery of DSMES. The DSMES team needs to consider characteristics such as racial identity, ethnic and cultural background, biological sex and gender identity, age, geographic location, technology access, education, literacy, and numeracy (13). Barriers to equitable DSMES access can be mitigated by

assessing the impact of the individual's SDOH and leveraging creative delivery options (e.g., telehealth and online) that will work best for the population in need of DSMES (10). For example, a systematic review and meta-analysis of telehealth DSMES interventions with Black and Hispanic adults with diabetes showed a 0.465% decrease in A1C, demonstrating the importance of considering demographic factors in relation to DSMES interventions (43).

Despite the recognized benefits of DSMES, fewer than 10% of individuals referred for DSMES through their health insurance or Medicare receive it, while about half the participants in an analysis of Behavioral Risk Factors Surveillance System (BRFSS) data report getting diabetes education (22,52). Barriers to DSMES exist at multiple levels, including the health system, payor, clinic, health care professional, and individual, for a myriad of reasons from lack of administrative leadership support to ineffective DSMES referral processes and transportation challenges. Low participation can be due to lack of referrals, logistical issues (e.g., accessibility, timing), cost, and lack of a perceived benefit (53). Thus, in addition to educating referring health care professionals about the benefits of DSMES and the critical times to refer, efforts to identify and address potential barriers at all levels need to be made (2). This was illustrated in a multilevel diabetes care intervention that combined clinical outreach, standardized protocols, and DSMES, with SDOH screening and referrals to social needs support. A 15% increase in receipt of DSMES, including among people on Medicaid, was documented (54). Support from institutional leadership is foundational for DSMES success. Expert stakeholders, including those external to an organization, should also support DSMES by advocating for it and for people with diabetes (10).

Diabetes Technologies

Technology-enabled diabetes self-management solutions (e.g., continuous glucose monitors [CGM], automated insulin delivery [AID] systems, and connected glucose meters) improve A1C most effectively when there is two-way communication between the person with diabetes and the health care team, individualized (e.g., tailored to suit a particular person for their needs) feedback, use of person-generated health data, and education (55). Technology

can facilitate self-management decisions and improve access to DSMES (55). Use of diabetes technologies warrants broader adoption because they can reduce therapeutic inertia and should be explored as part of continuous improvement (55,56). One potential model is virtual environments, which allow people with diabetes to self-represent as avatars and interact in a world with embedded informational resources accessed using principles of gamification. An example of this is from an RCT that tested DSMES in a virtual environment that demonstrated greater weight loss and similar decreases in A1C, blood pressure, cholesterol, and triglycerides compared with DSMES via a standard website (57). These nontraditional versions of DSMES may not always be reimbursed; however, adopting reimbursement policies that increase DSMES access and use will positively affect beneficiaries' clinical outcomes, quality of life, health care use, and costs (10,58).

Of all the newer diabetes technologies, CGM might be the most widely adopted. When combined with individualized DSMES or behavioral interventions, CGM demonstrated greater improvement of glycemic and psychosocial outcomes than CGM alone (59,60). Similarly, DSMES plus intermittently scanned CGM (isCGM) demonstrated increased time in range (70–180 mg/dL [3.9–10.0 mmol/L]), less time above range, and greater reduction in A1C compared with DSMES alone (61). Incorporating a systematic approach for technology assessment, adoption, and integration into the diabetes care plan could help ensure equity in access and standardized application of technology-enabled solutions (9,47,62–64).

Reimbursement

Medicare reimburses DSMES (known as diabetes self-management training [DSMT] by Medicare) when the services are provided in accordance with the National Standards for DSMES (2,10) and is recognized by the American Diabetes Association (ADA) through the Education Recognition Program (professional.diabetes.org/diabetes-education) or by the Association of Diabetes Care & Education Specialists (ADCES) (<https://www.adces.org/diabetes-education-dsmes/diabetes-education-accreditation-program>).

DSMES is also covered by most insurance plans. Ongoing support has been shown to be instrumental for improving outcomes when it is implemented after the completion of formal DSMES. For comprehensive

information about Medicare reimbursement, readers may find the following website useful: www.cdc.gov/diabetes-toolkit/php/reimbursement/medicare-reimbursement-guidelines.html. In brief, the Medicare Part B initial DSMT is a “once-in-a-lifetime” benefit. Individual encounters are reimbursable for the first 10 h (1 h of individual training and 9 h of group training). Two hours of follow-up DSMT are allowed each year after the initial DSMT. If a person has special needs that could interfere with effective group participation, these should be identified on the referral order to allow for individual sessions. For Medicaid, DSMES coverage varies by state, but further guidance can be found at <https://coveragetoolkit.org/evidence-based-programs/diabetes-self-management-education-and-support/>. Additional information addressing implementation of a successful DSMES program can be found in the Centers for Disease Control and Prevention DSMES toolkit at www.cdc.gov/diabetes-toolkit/php/index.html.

Programs recognized by the ADA and accredited by the ADCES were included on the list of telehealth professionals approved by Centers for Medicare & Medicaid Services (CMS) via the Consolidated Appropriations Act of 2023 (65). However, as of 1 October 2025, Medicare recipients must be in a U.S.-based rural office or medical facility for most telehealth services (66,67).

DSMES uses an evidence-based curriculum designed to educate people with diabetes about all elements from the National Standards for DSMES, as described above, that can be delivered and billed by a variety of health care professionals on the diabetes care team. While the overarching healthy eating concepts used in DSMES can be taught by all members of the team, MNT, which is more in-depth and individualized (e.g., tailored to suit a particular person for their sociocultural preferences and needs) and derived from the evidence-based Nutrition Care Process, can only be delivered and billed by RDNs. For Medicare Part B, the MNT benefit includes individual encounters billed 3 h in the first year of the benefit. Each subsequent year can be billed up to 2 h. However, additional hours are available if a subsequent referral identifies a change in treatment. For further information on Medicare coverage of MNT, readers are encouraged to review www.cdc.gov/diabetes-toolkit/php/reimbursement/medical-nutrition-therapy.html.

MEDICAL NUTRITION THERAPY

When the first ADA Standards of Care guidelines were published in 1989, nutrition was mentioned in only two sentences of the entire 4-page document (68). Even today, the science of nutrition for diabetes continues to evolve. There has also been a change in how we talk about nutrition. We continue to encourage health care professionals to shift away from emphasizing macronutrients (i.e., carbohydrates, proteins, and fats) and micronutrients (i.e., vitamins and minerals) and to instead focus on foods. More broadly, we encourage people to think in terms of eating patterns, also known as dietary patterns or food patterns, or the totality of the foods and beverages a person consumes. Additionally, encourage nutrient-dense food choices. Nutrient dense is defined as foods high in micronutrients while being relatively low in calories (e.g., vegetables, fruits, and legumes). This integrative food-based approach aligns with numerous guidelines from multiple professional health societies (69–71) and the *Dietary Guidelines for Americans, 2020–2025* (72). Simply put, people eat food, not nutrients, and nutrition recommendations need to be applicable to what people actually eat. Additionally, macronutrients are not interchangeable entities and vary by nutrient type and quality. For example, carbohydrates include legumes, whole grains, and fruits, which are in the same category as refined grains, but their health effects are quite different (73).

MNT is effective and beneficial to people with diabetes. When delivered by an RDN, MNT is associated with A1C absolute decreases of 1.0–1.9% for people with type 1 diabetes and 0.3–2.0% for people with type 2 diabetes (74). Because diabetes is progressive, behavior modification alone may not be adequate to maintain euglycemia over time. However, even after pharmacotherapy is initiated, MNT continues to be an important component of ongoing diabetes self-management, and RDNs providing diabetes-specific MNT should assess and monitor medication changes in relation to nutrition care plans (46,75). All health care team members should be empowered to reiterate the general and evidence-based nutrition advice promoted herein—limit processed foods and foods high in added salt, sugars, and fats and, when possible, choose whole foods.

For more detailed information on nutrition therapy for people with diabetes,

please refer to the ADA consensus report on nutrition therapy (46). Contained in the report is an important and often repeated tenet, i.e., there is no one-size-fits-all eating pattern for individuals with diabetes, and meal planning should be individualized. Nutrition therapy plays an integral role in overall diabetes management, and each person with diabetes should actively engage in education, self-management, and treatment planning with the health care team and participate in collaborative development of an individualized eating plan (46,75).

All people with diabetes should be referred for individualized MNT provided by an RDN who is experienced and skilled in providing diabetes-specific MNT (76–78), at diagnosis and as needed throughout the life span, like DSMES. Referrals to RDNs are particularly warranted when a person with diabetes is dealing with additional health conditions such as hypertension, dyslipidemia, heart failure, gastrointestinal disorders, chronic kidney disease (CKD), pregnancy-related nutrition concerns, pediatric growth issues, or obesity (79). See **Table 5.1** for nutrition recommendations and **Table 5.2** for nutrition behaviors that should be encouraged.

Eating Patterns and Meal Planning

To better understand the role of nutrition in diabetes, it is important to clarify terminology. Food patterns, eating plans, and approaches are terms that are often used interchangeably, but they are different and relevant in individualizing nutrition care plans (80).

- **Eating pattern, dietary pattern, or food pattern.** The totality of all foods and beverages consumed over a given period of time. An eating pattern can be ascribed to an individual, but it is also the term used in prospective cohort and observational nutrition studies to classify nutrition patterns. Examples include Mediterranean style, Dietary Approaches to Stop Hypertension (DASH), low carbohydrate, vegetarian, and plant based (80).
- **Eating/meal plan (historically referred to as a diet).** An individualized guide to plan when, what, and how much to eat on a daily basis, completed by the person with diabetes and the RDN. The eating plan could incorporate an eating pattern combined with a strategy to direct some of the choices. Eating plans are based on an individual's usual

Table 5.1—Nutrition recommendations

Recommendations	
5.10	Provide individualized medical nutrition therapy by referring people with prediabetes or diabetes to a registered dietitian nutritionist, preferably one who has comprehensive experience in diabetes care. A
5.11	Diabetes medical nutrition therapy can result in cost savings B and improved cardiometabolic outcomes A and should be reimbursed by insurance. E
5.12	Provide an overweight or obesity treatment plan based on their nutrition, physical activity, and behavioral health status for all people with overweight or obesity, aiming for at least 5–7% weight loss. A
5.13	For diabetes prevention and management of people with prediabetes or diabetes, recommend individualized meal plans that keep nutrient quality, total calories, and metabolic goals in mind. B
5.14	Eating patterns should emphasize key nutrition principles (inclusion of nonstarchy vegetables, whole fruits, legumes, lean proteins, whole grains, nuts and seeds, and low-fat dairy or nondairy alternatives) and minimize consumption of red meat, sugar-sweetened beverages, sweets, refined grains, processed and ultraprocessed foods in people with prediabetes and diabetes. B
5.15	Consider reducing carbohydrate intake for some adults with diabetes to improve glycemia. An effective way to achieve this is by limiting consumption of processed foods. B
5.16	Assess intake of supplements, as supplementation with micronutrients (e.g., vitamins and minerals, such as magnesium or chromium) or herbs or spices (e.g., cinnamon and aloe vera) is not recommended for glycemic benefits. C
5.17	Counsel against β -carotene supplementation, as there is evidence of harm for certain individuals and it confers no benefit. B
5.18	Advise adults with diabetes and those at risk for diabetes who consume alcohol to not exceed the recommended daily limits. B Advise abstainers to not start drinking alcohol, even in moderation. B
5.19	Counsel people with diabetes about the signs, symptoms, and self-management of delayed hypoglycemia and the importance of monitoring glucose after drinking alcohol to reduce hypoglycemia risk, especially when using insulin or insulin secretagogues. B
5.20	Counsel people with diabetes to limit sodium consumption to <2,300 mg/day, as clinically appropriate, and the best way to achieve this is through limiting consumption of processed foods. B
5.21	Encourage people with diabetes and those at risk for diabetes to consume water over other beverages. A
5.22	Counsel people with diabetes and those at risk for diabetes that nonnutritive sweeteners can be used in place of sugar-sweetened products if consumed in moderation and for the short term to reduce overall calorie and carbohydrate intake. B
5.23	Counsel and regularly monitor individuals pursuing intentional weight loss to ensure adequate nutritional intake, with particular attention to preventing protein insufficiency and micronutrient deficiencies. E
5.24	Emphasize minimally processed, nutrient-dense, high-fiber sources of carbohydrate (at least 14 g fiber per 1,000 kcal). B
5.25	Advise people with diabetes and those at risk for diabetes to replace sugar-sweetened beverages (including any juices) with water or low-calorie or no-calorie beverages and minimize foods with added sugar to manage glycemia and reduce risk for cardiometabolic disease. B
5.26	Educate individuals with diabetes who are at risk for developing diabetic ketoacidosis and who are treated with sodium–glucose cotransporter inhibition on the risks and signs of ketoacidosis and methods of risk mitigation management, provide them with appropriate tools for ketone measurement (i.e., serum β -hydroxybutyrate), and discourage a ketogenic eating pattern. E
5.27	Provide education on the glycemic impact of carbohydrate, A fat, and protein B tailored to an individual's needs, insulin plan, and preferences for care to optimize mealtime insulin dosing.
5.28	Counsel people using fixed insulin doses about consistent patterns of carbohydrate intake with respect to time and amount while considering the insulin action time, as it can result in improved glycemia and reduce the risk for hypoglycemia. B
5.29	Counsel people with diabetes and those at risk for diabetes to incorporate more plant-based protein sources (e.g., nuts, seeds, and legumes) as part of an overall diverse eating pattern to reduce cardiovascular disease risk. B
5.30	Counsel people with diabetes and those at risk for diabetes to consider an eating plan emphasizing elements of a Mediterranean eating pattern, which is rich in fatty fish, nuts, and seeds, to reduce cardiovascular disease risk A and improve glucose metabolism. B
5.31	Counsel people with diabetes and those at risk for diabetes to limit intake of foods high in saturated fat to help reduce cardiovascular disease risk. B

eating style, cultural background, SDOH, and food preferences.

- **Eating/meal plan approach.** Method to individualize a desired eating pattern and provide practical tools for developing healthy eating patterns. Examples include the plate method, carbohydrate choice,

carbohydrate counting, and highly individualized behavioral approaches (81).

Meal Planning

There is no ideal percentage of calories from carbohydrate, protein, or fat for

people with diabetes. Therefore, macronutrient distribution should be based on an individualized assessment of current eating patterns, preferences, and metabolic goals. Members of the health care team should complement and reinforce MNT by providing evidence-based guidance to help

Table 5.2—Nutrition behaviors to encourage

- Vegetables—especially nonstarchy vegetables that are dark green, red, and orange in color; fresh, frozen, or low-sodium canned are all acceptable vegetable options.
- Legumes—dried beans, peas, and lentils.
- Fruits—especially whole fruit—fresh, frozen, or canned in own juice (or no added sugar) are all acceptable fruit options.
- Foods with at least 3 g of fiber per serving are generally considered higher fiber choices. Whole-grain foods—where culturally appropriate, whole-grain versions of commonly consumed foods, such as 100% whole-wheat breads or pastas and brown rice. When not culturally appropriate, focus more on portion control.
- Water should be the primary beverage of choice.
- For individuals who do not prefer plain water, no-calorie alternatives are the next best choice. Options include adding lemon, lime, berries, or cucumber slices to water; sparkling no-calorie water or flavored no-calorie waters; no-calorie carbonated beverages.
- Plant-based proteins can include legumes (e.g., soybeans, pinto beans, black beans, garbanzo beans, dried peas, and lentils), nuts, and seeds.
- Meats and poultry should be from fresh, frozen, or low-sodium canned and in lean forms (e.g., chicken breast and ground turkey).
- Heart-healthy wild-caught fatty fish such as salmon, tuna, sardines, and mackerel. Fresh, frozen, or low-sodium canned are all acceptable options.
- Use herbs (e.g., basil, fennel, mint, parsley, rosemary, and thyme) and spices (e.g., cinnamon, garam masala, ginger, pepper, and turmeric) to season foods instead of salt or salt-containing preparations.
- Incorporate onions, garlic, celery, carrots, and other vegetables as a base for preparing various homemade foods.
- Cook with vegetable oil (e.g., avocado, canola, and olive) in place of fats high in saturated fat (e.g., butter, coconut oil, lard, and shortening).
- Plan out meals for the week. Grocery shop using a list. Cook on a day off so there are ready-to-eat and ready-to-reheat homemade meals waiting in the fridge or freezer.
- Include family or roommates in meal preparation; share the responsibilities of grocery shopping and cooking and use time off for meal preparation in advance when possible.

people with diabetes make healthy food choices that meet their individualized needs and improve overall health. Ultimately, ongoing diabetes and nutrition education paired with appropriate support to implement and sustain health behaviors are recommended (78).

Research confirms a variety of eating patterns are acceptable for diabetes management (46,74,82,83). Evidence-based eating patterns most frequently recommended include Mediterranean, DASH, low-fat, carbohydrate-restricted, vegetarian, and vegan eating patterns. Until evidence around benefits of specific eating patterns is strengthened, health care professionals should focus on the common core characteristics among healthful patterns: inclusion of nonstarchy vegetables, whole fruits, legumes, whole grains, nuts, seeds, and low-fat dairy products or nondairy replacements and minimizing consumption of red meat, sugar-sweetened beverages, sweets, refined grains, and processed and ultraprocessed foods (84,85). The recent Dietary Approaches to Stop Hypertension for Diabetes (DASH4D) RCT further supports this guidance (86). In their randomized crossover feeding study conducted from 2021 to 2024, adults with type 2 diabetes and

hypertension improved time in range (mean difference 5.2%; $P < 0.001$) and blood pressure. The intervention eating pattern was moderate in carbohydrates (45% of total calories), rich in fruits, vegetables, whole grains, lean proteins, and low-fat dairy, and low in saturated fat, sugar-sweetened beverages, and sodium.

Referral to and ongoing support from an RDN is essential to assess the overall nutrition status of the person with diabetes. RDNs work collaboratively with people to create a personalized meal plan aligned with the overall lifestyle treatment plan, including physical activity, work and life schedules, and medication use. Using shared decision-making to execute the plan is also often part of the nutrition care process.

Eating/M meal Plan Approaches and Methods

Few head-to-head studies have compared different eating approaches. The diabetes plate method (87) is a commonly used visual approach for providing basic meal planning guidance for individuals with type 1 and type 2 diabetes. One RCT found that both the diabetes plate method and carbohydrate counting were effective in helping achieve improved A1C (88). The

diabetes plate method uses a simple graphic (featuring a 9-in plate) to portion foods (one-half of the plate for nonstarchy vegetables, one-quarter of the plate for protein, and one-quarter of the plate for carbohydrates).

Carbohydrate counting is a more advanced skill that helps plan for and track how much carbohydrate is consumed at meals and snacks. In a systematic review and meta-analysis of carbohydrate counting versus other forms of meal planning advice (e.g., standard education, low glycemic index [GI], and fixed carbohydrate quantities), no significant differences were seen in A1C levels compared with standard education (89). In another RCT, a simplified carbohydrate counting tool based on individual glycemic response was noninferior to conventional carbohydrate counting in 85 adults with type 1 diabetes (90). In a randomized crossover trial, carbohydrate counting and qualitative meal size (i.e., low, medium, and high carbohydrate) were compared. Time in range was 74% for carbohydrate counting and 70.5% for the quantitative meal size estimates. Noninferiority was not confirmed for the qualitative method (91). Newer technologies (e.g., smart phone apps and CGM) and

AID systems may reduce the need for precise carbohydrate counting and allow for personalized nutrition approaches (92,93).

Meal planning approaches should be customized to the individual, including their numeracy and food literacy level (88). Health numeracy refers to understanding and using numbers and numerical concepts in relation to health and self-management. Food literacy generally describes proficiency in food-related knowledge and skills that ultimately affect health, although specific definitions vary across initiatives (94,95).

Nutrition Therapy Goals for All People With Diabetes

1. To promote and support healthful eating patterns, emphasizing a variety of nutrient-dense foods in appropriate portion sizes, contributing to improved overall health, and to:
 - achieve and maintain body weight goals
 - attain individualized glycemic, blood pressure, and lipid goals
 - delay or prevent the complications of diabetes
2. To address individual nutrition needs based on personal and cultural preferences, health literacy and numeracy, access to healthful foods, willingness and ability to make behavioral changes, and existing barriers to change
3. To maintain the pleasure of eating by providing nonjudgmental messages about food choices while also reducing or limiting certain foods only when indicated by scientific evidence
4. To provide an individual with diabetes the practical tools for developing healthy eating patterns rather than focusing on individual macronutrients, micronutrients, or single foods

Carbohydrates

Studies examining the optimal amount of carbohydrate intake for people with diabetes are inconclusive, although monitoring carbohydrate intake is a key strategy in reaching glucose goals in people with type 1 and type 2 diabetes (96,97).

The amount of carbohydrate intake for people with type 2 diabetes has been a focus of research for many years. Systematic reviews and meta-analyses of RCTs report carbohydrate-restricted eating patterns, particularly those considered very

low carbohydrate (<26% total energy), were effective in reducing A1C in the short term (<6 months), with less difference in eating patterns beyond 1 year (80,98,99). However, in a 12-week RCT among adults with prediabetes and type 2 diabetes, a well-formulated ketogenic eating pattern (20–50 g total carbohydrate/day and keeping protein to ~1.5 g/kg ideal body weight/day, with the remainder of energy from fat) did not significantly improve A1C and increased LDL cholesterol compared with a low-carbohydrate Mediterranean eating pattern (99). Additionally, a systematic review and meta-analysis of six RCTs at least 12 months in duration and including a total of 524 participants with type 2 diabetes reported that a low-carbohydrate eating pattern was beneficial for lipids but not glycemic management (standardized mean difference -0.11 , 95% CI -0.33 to 0.11 , $P = 0.32$) (100).

Reduction in body weight and the wide range of definitions for low-carbohydrate eating plans are important challenges in interpreting carbohydrate-restricted research studies (101–103). As studies on low-carbohydrate eating plans generally indicate challenges with long-term sustainability (104), it is important to reassess and individualize meal plan guidance regularly for those interested in this approach.

Health care professionals should maintain consistent medical oversight of individuals following very-low-carbohydrate eating plans and recognize that insulin and other diabetes medications may need to be adjusted to prevent hypoglycemia, and blood pressure will need to be monitored. In addition, very-low-carbohydrate eating plans are not currently recommended for individuals who are pregnant or lactating, children, people who have kidney disease, or people with or at risk for disordered eating (46).

Very-low-carbohydrate eating plans should be avoided in those taking sodium–glucose cotransporter 2 (SGLT2) inhibitors because of the potential risk of ketoacidosis (105,106). Numerous case reports have now been published illustrating that diabetic ketoacidosis (DKA) can occur in people with type 1 and type 2 diabetes using SGLT2 inhibitors in combination with very-low-carbohydrate or ketogenic eating patterns. Additionally, excessive alcohol intake should be avoided when taking SGLT2 inhibitors (105). Maintaining adequate hydration is also very important.

Regardless of carbohydrate quantity in the meal plan, the focus should be on high-quality, minimally processed, nutrient-dense, high-fiber carbohydrate sources. Fiber modulates gut microbiota composition and increases gut microbial diversity. Although there is still much to be elucidated about the gut microbiome and chronic disease, higher-fiber eating patterns are advantageous (107). Both children and adults with diabetes are encouraged to minimize intake of refined carbohydrates with added sugars, fat, and sodium and instead focus on carbohydrates from vegetables, legumes, fruits, dairy (milk and yogurt) or fortified nondairy alternatives, and whole grains. People with diabetes and those at risk for diabetes are encouraged to consume a minimum of 14 g of fiber/1,000 kcal, with at least half of grain consumption being whole, intact grains, according to the *Dietary Guidelines for Americans, 2020–2025* (72). Regular intake of sufficient fiber is associated with lower all-cause mortality in people with diabetes, and prospective cohort studies have found fiber intake is inversely associated with risk for type 2 diabetes (108, 109). Consumption of sugar-sweetened beverages and processed food products with large amounts of refined grains and added sugars is strongly discouraged (72), as these can displace healthier, more nutrient-dense foods and increase inflammation (110).

The literature on GI and glycemic load (GL) in individuals with diabetes is complex, often with varying definitions of low- and high-GI foods (111–113). The GI ranks carbohydrate foods on their postprandial glycemic response, and GL considers both the GI of foods and the amount of carbohydrate eaten. Studies report mixed effects of GI and GL on fasting glucose levels and A1C, with one systematic review finding no significant effect on A1C (112) while others demonstrated A1C reductions of 0.15% (111) to 0.5% (114,115). More recently, however, a meta-analysis of large cohorts ($\geq 100,000$ participants) reported that when people had larger intakes of high-GI foods, there was increased incidence of type 2 diabetes (risk ratio 1.27 [95% CI 1.21–1.34]; $P < 0.0001$), total cardiovascular disease (CVD) (1.15 [1.11–1.19]; $P < 0.0001$), diabetes-related cancer (1.05 [1.02–1.08]; $P = 0.0010$), and all-cause mortality (1.08 [1.05–1.12]; $P < 0.0001$) (113). It is important to note that low GI or low GL is synonymous with higher-fiber eating patterns.

Individuals with type 1 or type 2 diabetes taking insulin at mealtime should be offered comprehensive and ongoing education about nutrition content and the need to couple insulin administration with carbohydrate intake. For people whose meal schedule or carbohydrate consumption is variable, education on the relationship between carbohydrate intake and insulin needs is important. In addition, assessing food literacy, numeracy, interest, and capability should be evaluated, especially if teaching more advanced methods of MNT diabetes management such as insulin-to-carbohydrate ratios. Teaching insulin-to-carbohydrate ratios for meal planning can assist individuals with effectively modifying insulin dosing from meal to meal to improve glycemic management (74,96). Consumption of fat and protein can affect early and delayed postprandial glycemia (116), and it appears to have a dose-dependent response (117,118). However, more research is needed to determine the optimal insulin dose and delivery strategy. A cautious approach to increasing insulin doses for high-fat and/or high-protein mixed meals is recommended to address delayed hyperglycemia that may occur after eating (46,119). For individuals using an insulin pump, a split bolus feature (part of the bolus delivered immediately and the remainder over a programmed duration of time) may provide better insulin coverage for high-fat and/or high-protein mixed meals (120,121).

Insulin dosing decisions should be confirmed with a structured approach to blood glucose monitoring or CGM to evaluate individual responses and guide insulin dose adjustments. Checking glucose 3 h after eating may help determine if additional insulin adjustments are required (i.e., increasing or stopping bolus) (120,121). For individuals on a fixed daily insulin schedule, meal planning should emphasize a relatively fixed carbohydrate consumption pattern with respect to both time and amount while considering insulin action. Attention to hunger and satiety cues also helps (46).

Most commercially available AID systems still require basic diabetes management skills, including carbohydrate counting and understanding of the effect of protein and fat on postprandial glucose response, but the algorithms included in the systems work with less accurate carbohydrate entry (122,123). The most advanced AID system provides

adaptive closed-loop algorithms enabling fully autonomous insulin delivery automatically titrating all therapeutic insulins, including basal, correction, and prandial insulins. For more on AID and carbohydrates, see section 7, "Diabetes Technology."

Protein

There is no evidence that adjusting the daily protein intake above or below the recommended amount for the general public (typically 0.8–1.5 g/kg body weight/day or 15–20% of total calories) will improve health, and research is inconclusive regarding the ideal amount of dietary protein to optimize either glycemic management or CVD risk (72,124). Therefore, protein intake goals should be individualized based on current eating patterns. Some research has found successful management of type 2 diabetes with meal plans including slightly higher levels of protein (20–30%), which may contribute to increased satiety (125).

Historically, low-protein eating plans were advised for individuals with diabetes-related CKD (with albuminuria and/or reduced estimated glomerular filtration rate [eGFR]); however, current evidence does not suggest that people with CKD need to restrict protein to less than the generally recommended protein intake (126). Reducing the amount of protein below the recommended daily allowance of 0.8 g/kg is not recommended because it does not alter glycemic measures, cardiovascular risk measures, or the rate at which eGFR declines and may increase risk for malnutrition (126).

Growing evidence suggests higher plant protein intake and replacement of animal protein with plant protein is associated with lower risk of all-cause and cardiovascular mortality. A meta-analysis of 13 RCTs showed that replacing animal proteins with plant proteins leads to small improvements in A1C and fasting glucose in adults with type 2 diabetes (127). A 2023 systematic review and meta-analysis of 13 RCTs and 7 cohort studies concluded that there is limited suggestive evidence to support replacing animal protein with plant-based protein based on a moderate degree of bias in cohort studies (128). However, a prospective observational study of more than 11,000 community-dwelling adults over 22 years of follow-up reported that those with higher intakes of plant foods and lower intakes of animal foods had

lower diabetes risk (129). Plant proteins are lower in saturated fat, higher in fiber, and also support planetary health (130).

Fats

There is no optimal percentage of calories from fat for people with or at risk for diabetes, and macronutrient distribution should be individualized according to the individual's eating patterns, cultural and personal preferences, and metabolic goals (46). The type of fat consumed is more important than total amount of fat when looking at metabolic goals and CVD risk, and the percentage of total calories from saturated fats should be limited (72,131–133). Multiple RCTs including people with type 2 diabetes have reported that a Mediterranean eating pattern can improve both glycemic management and blood lipids (134–136). The Mediterranean eating pattern is based on traditional eating patterns in the countries bordering the Mediterranean Sea. Although eating styles vary by country and culture (i.e., customs and behaviors of a particular group of people or other social group), they share a number of common features, including consumption of fresh fruits and vegetables, whole grains, beans, and nuts/seeds; olive oil as the primary fat source; low to moderate amounts of fish, eggs, and poultry; and limited added sugars, sugary beverages, sodium, highly processed foods, refined carbohydrates, saturated fats, and fatty or processed meats.

People with diabetes should be advised to follow the same guidelines as the general population for the recommended intakes of saturated fat, cholesterol, and *trans* fat (72). In a 12-week double-blinded randomized controlled feeding study among 61 adults with overweight and obesity, without diabetes, higher intakes of saturated fat, compared with polyunsaturated fat, were found to increase liver fat deposition (137). A 2021 systematic review and meta-analysis including over 22,500 prospective study participants followed for 9.8 years reported that replacing saturated fats with other macronutrients, such as polyunsaturated fats, was associated with reduced CVD occurrence (138). *Trans* fats should be avoided. Importantly, it should be noted that as foods high in saturated fats are progressively decreased, they should be replaced with foods high in unsaturated fats and not with refined carbohydrate foods (139).

Evidence does not conclusively support recommending n-3 (eicosapentaenoic acid and docosahexaenoic acid) supplements for people with diabetes for the prevention or treatment of cardiovascular events (46,140). In individuals with type 2 diabetes, two systematic reviews with n-3 and n-6 fatty acids concluded that the dietary supplements did not improve glycemic management (141,142). In the ASCEND (A Study of Cardiovascular Events in Diabetes) trial, when compared with placebo, supplementation with n-3 fatty acids at a dose of 1 g/day did not lead to cardiovascular benefit in people with diabetes without evidence of CVD (143). However, results from the Reduction of Cardiovascular Events with Icosapent Ethyl-Intervention Trial (REDUCE-IT) found that supplementation with 4 g/day pure eicosapentaenoic acid significantly lowered the risk of adverse cardiovascular events. REDUCE-IT included 8,179 participants, of whom over 50% had diabetes, and found a 5% absolute reduction in cardiovascular events for individuals with established atherosclerotic CVD already treated with a statin with residual hypertriglyceridemia (135–499 mg/dL [1.52–5.63 mmol/L]) (144). See section 10, “Cardiovascular Disease and Risk Management,” for more information.

Sodium

As for the general population, people with diabetes are advised to limit their sodium consumption to <2,300 mg/day (46,145). Sodium intake has been shown to mediate glucose metabolism in a number of studies and affect eGFR, so limiting sodium intake is a valuable strategy for people with diabetes with or without kidney disease (145,146). In their post hoc analysis of the DASH-sodium RCT, Morales-Alvarez et al. (147) reported that participants randomized to the low-sodium DASH eating pattern (containing ~1,150 mg sodium/day [50 mmol sodium/day]) had change in eGFR of $-3.10 \text{ mL/min/1.73 m}^2$ (95% CI -5.46 to -0.73) after 4 weeks compared with 3,450 mg sodium/day (150 mmol sodium/day).

The DASH4D trial, which was a randomized 4-period crossover community-based feeding study, reported that in comparison with the eating pattern formulated with higher levels of sodium, the DASH4D eating pattern with lower sodium (1,500 mg/day at 2,000 kcal) resulted in improved systolic blood pressure

by 4.6 mmHg (95% CI, 7.2–2.0; $P < 0.001$) and diastolic blood pressure by 2.3 mmHg (95% CI, 3.7–0.9) (86). It was noted that the largest blood pressure reduction occurred during the first 3 weeks of each treatment period, and the effect of sodium reduction appeared stronger than the effect of the DASH4D eating pattern.

Limiting sodium intake is most easily achieved through focusing on whole, fresh foods. Additionally, it is important to reduce consumption of processed and ultraprocessed foods, which are major contributors of sodium intake. Encouraging people to avoid adding salt to foods and during cooking can also help. People providing nutrition advice should consider palatability, availability, affordability, clinical appropriateness, and the difficulty of achieving low-sodium recommendations in a nutritionally adequate eating plan.

Micronutrients and Other Supplements

Despite lack of evidence of benefit from dietary supplements, consumers continue to take them. Estimates show that up to 59% of people with diabetes in the U.S. use supplements (148). Without underlying deficiency, there is no benefit from herbal or other (i.e., vitamin or mineral) supplementation for people with diabetes (46,149).

U.S. federal law broadly defines dietary supplements as products having one or more dietary ingredients, including vitamins, minerals, herbs or other botanicals, amino acids, enzymes, tissues from organs or glands, or extracts of these (150). Dietary supplements are not regulated like other over-the-counter medications or prescription drugs in the U.S. (151). In combination with the strong views on dietary supplements (both positive and negative), this can contribute to consumer confusion (152). Readers can consult the U.S. Food and Drug Administration (FDA) Dietary Supplement Ingredient Directory to locate information about ingredients used in dietary supplements and any action taken by the agency with regard to that ingredient (153).

Specific nutrient supplementation for people with diabetes is generally not recommended outside the presence of deficiencies or overt malnutrition. Routine antioxidant supplementation (such as vitamins E and C) is not recommended due to lack of evidence of efficacy and concern related to long-term safety. Based

on the 2022 U.S. Preventative Services Task Force statement, the harms of β -carotene outweigh the benefits for the prevention of CVD or cancer. β -Carotene was associated with increased lung cancer and cardiovascular mortality risk (154).

Vitamin D in the context of diabetes has generated much research, but universal vitamin D supplementation for people with type 1 or type 2 diabetes without deficiency is not recommended at this time. Although post hoc analyses of the Vitamin D and Type 2 Diabetes Study (D2d) prospective RCT and Diabetes Prevention and Active Vitamin D (DPVD) and some meta-analyses suggest a potential benefit in specific populations (155–157), other studies have found no benefit or mixed results (158–160). Furthermore, adopting healthy lifestyle habits, including the eating patterns recommended herein, are strongly advised. Additional research is needed to define individual characteristics, clinical indicators, and appropriate dosages if and when vitamin D supplementation might benefit people with type 1 or type 2 diabetes.

There is also insufficient evidence to support routine use of herbal supplements and micronutrients, such as cinnamon (161), curcumin (e.g., turmeric), aloe vera, or chromium, to improve glycemia in people with type 1 or type 2 diabetes (46).

While not a dietary supplement, metformin is associated with vitamin B12 deficiency based on findings from the Diabetes Prevention Program Outcomes Study (DPPOS). Therefore, periodic testing of vitamin B12 levels should be considered in people taking metformin, particularly in those with anemia or peripheral neuropathy (162) (see section 9, “Pharmacologic Approaches to Glycemic Treatment”).

For special populations, including pregnant or lactating individuals, older adults, vegetarians, vegans, and people following very-low-calorie or low-carbohydrate eating patterns, a multivitamin may be necessary (163).

Alcohol

Long-term effects of alcohol consumption for people with diabetes are unknown. The World Health Organization declared there is no safe amount of alcohol intake (164,165). Associated risks of alcohol consumption include hypoglycemia and/or delayed hypoglycemia (particularly for those

using insulin or insulin secretagogue therapies), weight gain, and hyperglycemia (for those consuming excessive amounts) (46,166). People with diabetes should be educated about these risks and encouraged to monitor glucose frequently before and after drinking alcohol to minimize such risks. People with diabetes who consume alcohol can follow the same guidelines as people without diabetes consistent with the *Dietary Guidelines for Americans, 2020–2025* (72), which does not promote alcohol consumption in people who do not already drink. To reduce risk of alcohol-related harms, adults can choose not to drink or to drink in moderation by limiting intake to ≤ 2 drinks a day for men or ≤ 1 drink a day for women (one drink is equal to a 12-oz beer, a 5-oz glass of wine, or 1.5 oz of distilled spirits) (72). Recent meta-analyses have reported the previously recognized J-shaped relationship between alcohol intake and health risks likely varies by sex, obesity status, genetics, and alcohol intake behaviors (167,168).

Nonnutritive Sweeteners and Water

The FDA has approved many nonnutritive sweeteners (NNS) (containing few or no calories; commonly referred to as artificial sweeteners) for consumption by the general public, including people with diabetes (46,169). However, the safety and role of NNS continue to be sources of concern and confusion for the public.

For some people with diabetes who are accustomed to regularly consuming sugar-sweetened foods or beverages (e.g., regular soda pop, juice drinks, and other items sweetened with cane sugar or high-fructose corn syrup), NNS may be an acceptable substitute for nutritive sweeteners (those containing calories, such as sugar, honey, and agave syrup) when consumed in moderation (i.e., consuming no more than the acceptable daily intake) (170). NNS do not appear to have a significant effect on glycemic management (171,172), and they can reduce overall calorie and carbohydrate intake as long as individuals are not compensating with additional calories from other food sources (46,173). A meta-analysis and systematic review of RCTs found no evidence that NNS raise liver enzymes (174).

There is mixed evidence from systematic reviews and meta-analyses for NNS use with regard to weight management, with some finding benefit for weight loss

(175–177) while other research suggests an association with weight gain (178,179). This may be explained by reverse causality and residual confounding variables (179). The addition of NNS to eating plans poses no benefit for weight loss or reduced weight gain without energy restriction (180). In a systematic review and meta-analysis using low-calorie and no-calorie sweetened beverages as an intended substitute for sugar-sweetened beverages, a small improvement in body weight and cardiometabolic risk factors was seen without evidence of harm and had a direction of benefit similar to that seen with water (181). While health care professionals should promote water as the healthiest beverage option, people with overweight or obesity and diabetes may also use a variety of no-calorie or low-calorie sweetened products so that they do not feel deprived (181).

Health care professionals should encourage reductions in foods and beverages with added sugars and promote reducing overall sugar intake and calories with or without the use of NNS. Assuring people with diabetes that NNS have undergone extensive safety evaluation by regulatory agencies and are continually monitored can allay unnecessary concern for harm. Health care professionals can regularly assess individual use of NNS based on the acceptable daily intake (amount of a substance considered safe to consume each day over a person's life) and recommend moderation. Readers are directed to the FDA for additional information on NNS including safety, graphics, and the acceptable daily intake levels for the six major ones used in the U.S. ([fda.gov/food/food-additives-petitions/aspartame-and-other-sweeteners-food](https://www.fda.gov/food/food-additives-petitions/aspartame-and-other-sweeteners-food)).

Weight Management

Management and reduction of weight is important for people with type 1 diabetes, type 2 diabetes, or prediabetes with overweight or obesity. To support weight loss and improve A1C, CVD risk factors, and well-being in adults with overweight or obesity and prediabetes or diabetes, MNT and DSMES services should include an individualized eating plan resulting in an energy deficit in combination with enhanced physical activity (46). Lifestyle intervention programs should be intensive and have frequent follow-up to achieve significant reductions in excess body weight and

improve clinical indicators. Behavior modification goals should address physical activity, calorie restriction, healthy weight management strategies, and motivation. There is strong and consistent evidence that modest, sustained weight loss can delay the progression from prediabetes to type 2 diabetes (78,182,183) (see section 3, "Prevention or Delay of Diabetes and Associated Comorbidities") and is beneficial for type 2 diabetes management (see section 8, "Obesity and Weight Management for the Prevention and Treatment of Diabetes").

In prediabetes, the weight loss goal is at least 5–7% from baseline body weight, and it is higher for reducing risk of progression to type 2 diabetes. Some RCTs report less than 5% weight loss in adults with diabetes and overweight or obesity following a lifestyle behavioral intervention, but this limited amount of weight loss has not been shown to improve glycemia, lipids, or blood pressure; rather, a minimum weight loss of 5% or more seems necessary to achieve metabolic improvements (184). In conjunction with support for healthy lifestyle behaviors, medication-assisted weight loss can be considered for people at risk for type 2 diabetes when needed to achieve and sustain 7–10% weight loss (185,186) (see section 8, "Obesity and Weight Management for the Prevention and Treatment of Diabetes"). People with prediabetes and healthy weight range should also be considered for behavioral interventions to help establish routine aerobic and resistance exercise (182,187,188) and healthy eating patterns. Services delivered by health care professionals familiar with diabetes and its management, such as an RDN, have been found to be effective (77).

For many individuals with overweight or obesity alongside type 2 diabetes, at least 5% weight loss is needed to achieve beneficial outcomes in glycemic management, lipids, and blood pressure (189). However, any magnitude of weight loss is recommended. It also should be noted that the clinical benefits of weight loss are progressive, and more intensive weight loss goals (i.e., 15%) may be appropriate to maximize benefit depending on need, feasibility, and safety (190,191). Long-term sustainability of weight loss remains a challenge (192). For example, in some South Asian adult populations, traditional interventions have not been as effective in prevention or remission of type 2 diabetes, so

those groups will benefit from more culturally tailored interventional approaches (193).

Medications can augment MNT to support weight loss, weight loss maintenance, and improve cardiovascular outcomes. Newer medications (e.g., glucagon-like peptide 1 receptor agonists [GLP-1 RAs]) may be more viable, positively affect cardiovascular outcomes, and produce weight reduction beyond 10–15% (194–198). For more information on the nutritional considerations important for people undergoing significant weight loss, see *MONITORING NUTRITION INTAKE*, below.

Overweight and obesity are increasingly prevalent in people with type 1 diabetes and present clinical challenges regarding diabetes treatment and CVD risk factors (199,200). There is some evidence that GLP-1 RAs are useful in achieving weight loss among adults with type 1 diabetes, although with a higher risk of nausea and ketosis (201).

Regardless of diabetes type, weight loss maintenance is challenging (184,202) but has well-recognized long-term benefits. Weight loss maintenance physiology is complex and involves many hormonal, psychosocial, behavioral, and environmental factors. Following a weight loss of at least 8%, a subsequent “weight loss maintenance” intervention was reported to be only moderately beneficial, as it helped sustain physical health improvements but not glucose metabolism improvements (203). However, in another RCT with long-term, real-world, clinic-based follow-up of 10 years, Tomah et al. (204) reported lasting glycemic benefits in their cohort with an average weight loss of 7.7 ± 0.9 kg ($-6.9 \pm 0.8\%$) maintained for 10 years.

Starting a conversation about weight management should be based on motivational interviewing techniques (205), beginning with first asking the individual if they want to discuss their weight. Health care professionals should never assume a person with overweight or obesity wants to discuss weight at a medical appointment, especially if the appointment is for a seemingly unrelated issue (e.g., back pain, which many people do not realize is often secondary to excess body weight). Using person-centered approaches to weight management conversations involves meeting the individual where they are in their life and working with what they and their health care professional agree is the most beneficial approach. Guidance from an RDN

with expertise in motivational interviewing and diabetes and weight management MNT during any comprehensive structured weight loss program is strongly recommended.

Along with routine medical management visits, people with diabetes and prediabetes should be screened during DSMES and MNT encounters for a history of dieting and past or current disordered eating behaviors. Characterizing an individual's past efforts with weight loss and their body weight history can also be very useful. Nutrition therapy should be individualized to help address maladaptive eating behavior (e.g., purging) or compensatory changes in medical treatment plan (e.g., overtreatment of hypoglycemic episodes and reduction in medication dosing to reduce hunger) (46) (see *DISORDERED EATING BEHAVIOR*, below). Caloric restriction may be necessary for glycemic and weight management, but rigid meal plans and strict tracking of food intake and/or body weight can be contraindicated for individuals at increased risk of significant maladaptive eating behaviors (206). If eating disorders are identified during screening with diabetes-specific questionnaires, individuals should be referred to a qualified behavioral health professional (1).

Meal Replacements

Use of partial or total meal replacements is an additional strategy for energy restriction. Meal replacements are prepackaged foods (bars, shakes, and soups) that contain fixed amounts of macronutrients and micronutrients. They can improve nutrient quality and glycemic management and, consequently, reduce portion size and energy intake. In a meta-analysis involving 17 studies incorporating both partial and total meal replacements, greater weight loss and improvements in A1C and fasting blood glucose were demonstrated compared with conventional meal plans (207). Furthermore, meal replacements have been used in several landmark clinical trials, including Look AHEAD (Action for Health in Diabetes) (208), DiRECT (Diabetes Remission Clinical Trial) (209), and PREVIEW (Prevention of Diabetes Through Lifestyle Intervention and Population Studies in Europe and Around the World) (210). Results of these trials showed that partial or total meal replacements can be a potential short-term (i.e., <6 months) strategy for weight loss.

Regardless of the specific eating pattern or meal plan selected, long-term follow-up and support from members of the diabetes care team are needed to optimize self-efficacy and maintain behavioral changes (211).

Fasting and Timing of Food Intake

Chrononutrition is an emerging nutrition and biology subspecialty aimed toward increasing the understanding of how the timing of food ingestion affects metabolic health (212). Glucose metabolism follows a circadian rhythm through diurnal variation of glucose tolerance and peaks during daylight hours when food is consumed. Some preliminary studies show cardiometabolic benefits when food is consumed earlier (213). Similarly, circadian disruptions found in shift workers increase risk of type 2 diabetes (214). This evolving area of research currently lacks conclusive evidence, but future studies are anticipated.

Nonreligious Fasting

The primary forms of nonreligious fasting are intermittent fasting or time-restricted eating. These are popular strategies for weight and glucose management. One of the key distinctions between nonreligious and religious fasting is water intake. See **Fig. 5.1** for further details on how religious and nonreligious fasting practices compare.

Intermittent fasting is an umbrella term that includes three main forms of restricted eating: alternate-day fasting (energy restriction of 500–600 calories on alternate days), the 5:2 eating pattern (energy restriction of 500–600 calories on consecutive or nonconsecutive days with usual intake the other five), and time-restricted eating (daily calorie restriction based on window of time of 8–15 h). Each produces mild to moderate weight loss (3–8% loss from baseline) over short durations (8–12 weeks) with no significant differences in weight loss when compared with continuous calorie restriction (215,216). A 2024 systematic review and meta-analysis of RCTs examined the most common types of fasting in studies lasting 2–52 weeks. The authors concluded that intermittent energy restriction produces small but significant reductions in waist circumference and fat-free mass but were otherwise not superior to continuous energy restriction eating patterns (217). Generally, time-restricted eating or shortening the eating window can be adapted to any

Religious and intermittent fasting: differences and similarities




Figure 5.1—Differences and similarities between religious and intermittent fasting for people with diabetes.

eating pattern and has been shown to be safe for adults with type 1 or type 2 diabetes (218). People with diabetes who are taking insulin and/or secretagogues should be medically monitored during the fasting period (219). Because of the simplicity of intermittent fasting and time-restricted eating, these may be useful strategies for people with diabetes who are looking for practical eating management tools. It should also be noted that the principles of healthy eating still apply during non-fasting times.

Religious Fasting

Recommendations

5.32 Use the updated International Diabetes Federation along with Diabetes and Ramadan International Alliance comprehensive prefasting risk assessment to generate a risk score for the safety of religious fasting. Provide fasting-focused education to minimize risks. **B**

5.33 Assess and optimize treatment plan, dose, and timing for people with diabetes well in advance of religious fasting to reduce risk of hypoglycemia, dehydration, hyperglycemia, and/or ketoacidosis. **B**

Although intermittent fasting and time-restricted eating are specific dietary strategies for energy restriction, religious fasting has been practiced for thousands of years and is part of many faith-based traditions. Duration, frequency, and type of fast vary among different religions (220). For example, Jewish people abstain from any intake for ~25 h during Yom Kippur (221,222). For Muslims, Ramadan fasting lasts for a full month, when abstinence from any food or drink is required from dawn to dusk (223). Individuals with diabetes who fast have an increased risk for hypoglycemia, dehydration, hyperglycemia, and ketoacidosis (224,225).

Prefasting risk assessment is essential to increase level of safety (224,225). Various risk factors need to be considered for every individual wishing to fast. Some of these factors are related to the type of fast, type of diabetes, and/or the individual. Indeed, health care professionals should inquire about any religious fasting for people with diabetes and provide education and support to accommodate their choice. The number of fasting days is an important factor to consider. In Ramadan fasting, a person fasts from dawn to dusk for a lunar

month (29–30 days). It is important for the health care professional to comprehensively assess these risk factors well in advance of the planned fasting date, as some of them are modifiable. Some of these factors are related to the nature of the fasting practice, others are related to diabetes, and others might be due to individual factors such as the physical intensity of their work and/or the nutrition choices when they break their fast. The International Diabetes Federation along with Diabetes and Ramadan International Alliance adopted a risk calculator for the various risk factors (224,226). Several clinical studies from different countries have been published that assess the validity of the fasting risk score and the ease of use of it (226–229). The accumulation of these risk factors provides a risk score as low, moderate, or high (**Table 5.3**) (224). While the risks of different religious fasting practices may vary, this risk calculator provides some useful guidance for other types of religious fasting.

Prefasting education regarding the importance of increasing the frequency of glucose monitoring for people wishing to fast is very important. Timing of glucose monitoring is also especially important,

Table 5.3—Elements for risk calculation and suggested risk score for people with diabetes who seek to fast during Ramadan

Fasting risk element	Risk score
1. Pregnancy with any type of diabetes	
• Yes	6.5
2. Diabetes type	
• Type 1 diabetes or LADA	1
• Type 2 diabetes or any other type of diabetes	0
3. Duration of diabetes (years)	
• >20 years	1
• 10–20 years	0.5
• <10 years	0
4. Type of diabetes treatment (select all that are relevant)	
• Multiple daily premixed insulin injections	2
• Once-daily premixed insulin	1.5
• Open-loop insulin pump	1.5
• Automated insulin delivery system	1
• Standard basal insulin (NPH, detemir, or glargine 100)	1
• Ultra-long-acting basal insulin (glargine 300 or degludec)	0.75
• Short-acting insulin	0.75
• Glibenclamide or glipizide	0.75
• Modern sulfonylurea (gliclazide, gliclazide MR, glimepride) or repaglanide	0.5
• ≥2 glucose-lowering medications excluding insulin or sulfonylurea	0.25
• Nutrition modification only or monotherapy (excluding insulin or sulfonylurea)	0
5. Presence of hypoglycemia	
• Impaired hypoglycemia awareness	6.5
• Severe* hypoglycemia during last 4 weeks	5
• Hypoglycemia more than once daily	4
• 6–7 episodes of hypoglycemia/week	3
• 3–5 episodes of hypoglycemia/week	2
• 1–2 episodes of hypoglycemia/week	1
• Hypoglycemia <1 time per week	0.5
• No hypoglycemia in last 4 weeks	0
6. Level of A1C	
• >9% (>75 mmol/mol)	1
• 7.5–9% (58–75 mmol/mol)	0.5
• <7.5% (<58 mmol/mol)	0
7. Glucose monitoring	
• Not done	2
• Done suboptimally	1
• Done as indicated	0
• Any type of CGM	–0.5
8. Hyperglycemic emergencies	
• DKA or HHS in the last month	3.5
• DKA or HHS in last 2–3 months	2
• DKA or HHS in last 4–6 months	1
• No DKA or HHS in last 6 months	0
9. Macrovascular** complications	
• Unstable macrovascular disease	6.5
• Stable macrovascular disease	2
• No macrovascular disease	0
10. Microvascular complications	
a. Nephropathy	
• eGFR <30 mL/min	6.5
• eGFR 30–45 mL/min	4
• eGFR 45–60 mL/min	2
• eGFR >60 mL/min	0
b. Neuropathy, foot complications, or diabetic retinopathy	
• 3 microvascular complications	3
• 2 microvascular complications	2
• 1 microvascular complication	1
• 0 microvascular complications	0

Continued on p. S102

as the last few hours of fasting are frequently associated with approximately 50% of hypoglycemic events (230). Consequently, avoiding intense physical activity during the last few hours of fasting seems to be a sensible approach.

During religious fasting, some people change their nutrition habits and overindulge after fasting concludes. In many communities, the meal consumed to break the fast is rich in carbohydrates and includes foods and beverages high in added sugars and fat (224). Indeed, in one recent study 16.5% of people with type 2 diabetes who fasted for Ramadan reported high blood glucose of >300 mg/dL (>16.6 mmol/L) during fasting days (230). Individualized fluid adjustment and meal advice should be provided with emphasis on higher intake of fiber and replacing added sugars with complex carbohydrates to minimize hypoglycemia and hyperglycemia and emphasis on sustaining adequate daily fluid intake (231).

Treatment before and after fasting should be culturally sensitive and individualized. Specific recommendations for diabetes management during religious fasting in different faiths are available (224,225). In general, for people planning to fast for long hours and multiple consecutive days, choice of treatment should prioritize drugs with low hypoglycemia risk. Hypoglycemia risk while fasting in people using insulin, sulfonylureas, and other insulin secretagogues is higher than that for individuals treated with other types of diabetes medications (224). The safety of SGLT2 inhibitors was assessed in several studies during Ramadan fasting. These studies did not show significant change in kidney function, dehydration rates, or ketosis (232). Guidelines do not advise any change in SGLT2 inhibitor dose during fasting; however, they advise against initiating SGLT2 inhibitors close to the start of fasting days to avoid excessive thirst (224). **Table 5.4** summarizes the effect of fasting on different treatment options and the possible change in doses or timing for people with diabetes.

Technology could be an important tool to enhance safety during fasting. Several studies have investigated the use of monitoring technology during Ramadan fasting (e.g., flash glucose monitoring and CGM) and confirmed that these tools are able to support high-risk groups wishing to fast, especially if combined with Ramadan-focused education (232–234). Meanwhile, the use of insulin pumps

Table 5.3—Continued

Fasting risk element	Risk score
11. Cognitive function, frailty, and age	
• Impaired cognitive function	6.5
• Advanced frailty	6.5
• Mild to moderate frailty	4
• Age >70 years with no home support	1
• Normal cognitive function and no frailty	0
12. Physical labor	
• High intensity	4
• Moderate intensity	2
• Low intensity	0
13. Fasting-focused education	
• Yes	0
• No	1
14. Fasting hours	
• ≥16 h	1
• <16 h	0

Based on risk scoring, people with diabetes can be categorized as having:

- Score 0–3.0: low risk, fasting is probably safe
- Score 3.5–6.0: moderate risk, fasting safety is uncertain
- Score >6.0: high risk, fasting is probably unsafe

A1D, automated insulin delivery; CGM, continuous glucose monitoring; DKA, diabetic ketoacidosis; eGFR, estimated glomerular filtration rate; HHS, hyperglycemic hyperosmolar state; LADA, latent autoimmune diabetes in adults (type 1 diabetes). *Hypoglycemia requiring assistance for treatment. **Macrovascular disease includes cardiac, cerebral, or peripheral. Adapted from Hassanein et al. (224).

has been associated with low rates of hypoglycemia during fasting in people with type 1 diabetes. Diabetes technologies should be considered as a useful adjunct to risk calculation and/or nutrition planning and education during religious fasting for people with diabetes (224).

Monitoring Nutrition Intake

It is important to monitor for adequate nutritional intake to prevent nutrient deficiencies, especially in people with diabetes who are seeking intentional weight loss. Health care professionals should advise a healthy, whole foods–based eating pattern with sufficient micro- and macronutrients, protein, fiber, and fluid intake alongside regular resistance training to maintain lean body mass and prevent nutritional deficiencies (235,236). Referral to an RDN is also important to optimize nutrition status, especially if they have experience conducting a nutrition-focused physical exam (235,237) (see section 8, “Obesity and Weight Management for the Prevention and Treatment of Diabetes”).

Food Insecurity and Access

Food insecurity is a household-level economic and social condition of limited or

uncertain access to adequate food (238). In 2023, 13.5% of Americans were food insecure (238), and food insecurity affects 16% of adults with diabetes compared with 9% of adults without diabetes (239). There is a complex bidirectional association between food insecurity and co-occurring diabetes. Food security screening should happen at all levels of the health care system. Any member of the health care team can screen for food insecurity using the Hunger Vital Sign. Households are considered at risk if they answer either or both of the following statements as “often true” or “sometimes true” (compared with “never true”) (240):

- “Within the past 12 months, we worried whether our food would run out before we got money to buy more.”
- “Within the past 12 months, the food we bought just didn’t last, and we didn’t have money to get more.”

If screening is positive for food insecurity, efforts should be made to refer to appropriate programs and resources. See section 1, “Improving Care and Promoting Health in Populations,” for more information concerning the social determinants of health and related issues like food insecurity and access.

PHYSICAL ACTIVITY

Recommendations

5.34 Evaluate baseline physical activity and sedentary time for all people with diabetes and those at risk for diabetes. For people who do not meet activity guidelines, encourage an increase in physical activities above baseline with the goal of meeting activity guidelines. **B** Counsel that prolonged sitting should be interrupted at least every 30 min for blood glucose and other benefits. **C**

5.35 Counsel children and adolescents with type 1 diabetes **C** or type 2 diabetes **B** to engage in 60 min/day or more of moderate- or vigorous-intensity aerobic activity, with muscle-strengthening and bone-strengthening activities at least 3 days/week, and to limit the amount of time being spent sedentary, including recreational screen time. **C**

5.36 Counsel most adults with type 1 diabetes **C** and type 2 diabetes **B** to engage in 150 min or more of moderate- to vigorous-intensity aerobic activity per week, spread over at least 3 days/week, with no more than 2 consecutive days without activity. Shorter durations (minimum 75 min/week) of vigorous-intensity or interval training may be sufficient for more physically fit individuals.

5.37 Counsel adults with type 1 diabetes **C** and type 2 diabetes **B** to engage in 2–3 sessions/week of resistance exercise on nonconsecutive days.

5.38 Counsel most older adults with diabetes to engage in flexibility training and balance training 2–3 times/week. **C**

5.39 Counsel all people with diabetes who are treated with obesity pharmacotherapy or metabolic surgery that meeting physical activity recommendations, in particular muscle-strengthening exercises, may be beneficial for maintaining lean body mass. **C**

Physical activity includes all movement that increases energy use, and exercise is a more specific form of physical activity that is structured and designed to improve physical fitness. Both physical activity and exercise demonstrate numerous benefits for people with and at

Table 5.4—Changes in medications during fasting

Medication name	Risk of hypoglycemia	Timing	Total daily dose
Metformin, SGLT2 inhibitor, DPP-4 inhibitor, GLP-1 receptor agonist, acarbose, or pioglitazone	Low	• If once daily, then take at main mealtime. • If twice daily, then split dose between the two meals. • If once weekly, no change of time.	• No change
New generation of sulfonylurea (glimepiride and gliclazide)	Low to moderate	• If once daily, then take at main mealtime. • If twice daily, then split dose between the two meals.	• Reduce dose if glucose levels are within individualized goal range and if no hypoglycemia or hyperglycemia is present at baseline.
Older generation of sulfonylurea (glyburide)	Moderate to high	• Take at time of main meal	• Replace with newer-generation sulfonylurea or reduce dose by 50%.
Basal insulin	Moderate to high	• For longer-acting basal analogs (glargine 300 or degludec), no need to change timing. • For other basal insulins, take at beginning of breaking fast meal.	• Choose the insulin with lower risk of hypoglycemia among the class. • Reduce dose by 25–35% if not well managed.
Prandial insulin	High	• At mealtime	• Reduce dose of insulin for the meal followed by fasting (35–50%). • For other meals, insulin dose should match carbohydrate intake.
Mixed insulin and insulin coformulations	High	• If once daily, then take at main mealtime. • If twice daily, then split dose between the two meals	• Reduce dose of insulin for the meal followed by fasting (35–50%). • For other meals, no change of dose.

DPP-4, dipeptidyl peptidase 4; GLP-1, glucagon-like peptide 1; SGLT2, sodium–glucose cotransporter 2.

risk for diabetes and are important for the diabetes management plan. Higher levels of total and leisure-time physical activity are associated with a lower risk of cardiovascular and overall mortality in people with diabetes (241,242). Leisure-time activity may also help prevent type 2 diabetes (243) and reduce A1C in those with diabetes (244). Moreover, data from the DPPOS noted that self-reported habitual physical activity is positively associated with 6-min walking distance, which suggests long-term benefits of regular physical activity engagement (245). Exercise and physical activity have favorable effects on glycemia, cardiovascular risk factors, weight loss, body composition, mobility, mortality, psychosocial well-being, and physical function in people with diabetes (242,246–254).

Given the benefits of physical activity and exercise for people with diabetes, they should be recommended and prescribed to all individuals who are at risk for or have diabetes as part of the diabetes care plan, unless otherwise contraindicated. Specific recommendations and precautions on the type of activity will vary by diabetes type, age, and presence of complications. Health care professionals should support people with diabetes to set

stepwise goals toward meeting the recommended exercise volume. As individuals intensify their exercise program, medical monitoring may be indicated to ensure safety and evaluate the effects on glycemic management. Exercise and activity plans should be tailored to meet the specific needs of each individual (255), and different strategies can be used to increase engagement (256,257). Individuals with diabetes may experience obstacles to exercise, such as not having enough time or inadequate access to equipment or education for safe participation. The care team should help identify these obstacles and individualize approaches to improve long-term adoption of physical activity and exercise as a key part of the diabetes care plan.

Furthermore, activity plans can be modified to best suit the fitness level of the individual, which may vary due to ability level or complications. The plan might also need to be modified over time due to changes in health status, goals, or preferences or if the therapeutic response reaches a plateau. For this reason, individuals with diabetes benefit from a team-based approach, including working with an exercise physiologist, physical therapist,

or personal trainer, among others, where available and affordable (258). The ADA position statement “Physical Activity/Exercise and Diabetes” reviews the evidence for the benefits of exercise in people with type 1 and type 2 diabetes and offers more specific guidance (255).

Physical Activity and Glycemic Management

In people with type 2 diabetes, exercise training has demonstrated multiple cardio-metabolic improvements, including but not limited to improved glycemia, cardiovascular function, and lipid profiles (259,260). A meta-analysis primarily consisting of aerobic training studies revealed that structured exercise interventions of at least 8 weeks have been shown to lower A1C by 0.66% in people with type 2 diabetes, even without a significant change in BMI (247). Data also suggest that combining aerobic and resistance exercise may provide greater glycemic benefits compared with performing either mode of exercise alone (260).

In people with type 1 diabetes, the data for improving A1C are somewhat unclear (261,262). Interestingly, adding resistance training to an exercise plan for adults with type 1 diabetes who were

already aerobically active did not impact glycemia, even with improvements in waist circumference and strength (263). Real-world data from the Type 1 Diabetes and Exercise Initiative (T1DEXI) show that aerobic exercise reduces glycemia after an acute exercise session compared with high-intensity interval exercise and resistance exercise; however, all three modes of exercise improve time in range over a 24-h period (264). Furthermore, meeting daily step count of ~7,000–10,000 steps/day may support marginal improvements in time in range and reduced insulin requirements (265). Though there may be some benefits in glycemic response to exercise, this variability should be taken into consideration when recommending the type, intensity, and duration of exercise and insulin adjustments for a given individual with type 1 diabetes to prevent hypoglycemia and hyperglycemia (266,267).

Individuals of childbearing potential with preexisting diabetes, particularly type 2 diabetes, and those at risk for or presenting with gestational diabetes mellitus should be encouraged to engage in regular moderate-intensity physical activity prior to and during their pregnancies, as tolerated (255). Regular exercise can reduce the odds of developing gestational diabetes mellitus by nearly 40% (268), and acute and long-term prenatal exercise can positively impact glycemia, with a larger effect noted in people with diabetes (269). For more information, see section 15, “Management of Diabetes in Pregnancy.”

Evaluating Pre-exercise Risk and Baseline Physical Activity

As discussed more fully in section 10, “Cardiovascular Disease and Risk Management,” the best protocol for assessing asymptomatic people with diabetes for coronary artery disease remains unclear. The ADA consensus report “Screening for Coronary Artery Disease in Patients With Diabetes” (270) concluded that routine testing is not recommended. However, health care professionals should perform a careful history prior to exercise engagement, assess cardiovascular risk factors, be aware of the atypical presentation of coronary artery disease, such as a recently reported or measured decrease in exercise tolerance, and determine if testing for cardiac abnormality (e.g., arrhythmia) is warranted. Certainly, those with high risk should be encouraged to start with short periods of low-intensity exercise and

slowly increase the duration and intensity as tolerated. Additionally, individuals with complications may warrant a more thorough evaluation prior to engaging in a physical activity or exercise plan to ensure the plan is safe and appropriate. Health care professionals should assess for conditions that might contraindicate certain types of exercise or predispose to injury, such as inadequately managed hypertension, untreated proliferative retinopathy, autonomic neuropathy, orthostatic hypotension, peripheral neuropathy, balance impairment, and a history of foot ulcers or Charcot foot.

An evaluation of baseline physical activity and time spent in sedentary behavior is recommended to help guide initial increases in physical activity levels. When discussing physical activity, health care professionals should consider inquiring about motivators (e.g., health benefits, enjoyable activities with family) and potential barriers (e.g., lack of time, fear of hypoglycemia, no safe place to perform activity) to exercise and activity participation (271,272). Understanding these barriers may help with devising practical solutions for sustainable exercise participation, as complete uptake of the recommended total exercise volume may be initially challenging for some people with diabetes (273,274). People who do not meet activity guidelines should be encouraged to increase physical activity levels above baseline by performing activities such as walking, yoga, housework, gardening, swimming, and dancing (275). Sedentary behaviors (e.g., time seated at work, on a computer, tablet, smartphone, or watching television) should be discussed regularly. People with diabetes should be counseled to reduce sedentary time and break up sedentary activities at least every 30 min by briefly standing, walking, or performing other light physical activities to support glycemic and other cardiometabolic benefits, such as vascular function (276–279).

Exercise for Children and Adolescents

Similar to children and adolescents in general, children and adolescents with diabetes or who are at risk for diabetes should be encouraged to engage in regular physical activity, including at least 60 min of moderate-to-vigorous aerobic activity every day and muscle- and bone-strengthening activities at least 3 days per week to support development and health

benefits (254,280). Exercise programs promoting nutrition modification, increasing physical activity levels, and reducing sedentary time in children and adolescents at risk for type 2 diabetes have been shown to reduce risk of type 2 diabetes development (281–283), and vigorous-intensity physical activity participation is associated with lower CVD risk in youth with type 2 diabetes (284). In general, children and adolescents with type 1 diabetes benefit from being physically active, with meta-analyses reporting significant associations between physical activity and lower A1C (285) and an absolute A1C reduction of nearly 0.8% (286). Thus, an active lifestyle should be recommended to all children and adolescents with diabetes to support health outcomes and health-related quality of life (287,288). Children and adolescents are recommended to limit sedentary time, including recreational screen time, to less than 2 h per day (289–291). See section 14, “Children and Adolescents,” for details.

Frequency and Type of Physical Activity

A comprehensive physical activity and exercise plan should encompass multiple modes of activities, including aerobic, resistance, flexibility, balance, and leisure-time activities. Each activity type offers unique benefits for diabetes management, overall health, and quality of life. Age, previous physical activity experiences, fitness level, and goals should be considered when discussing physical activity and exercise plans. For more specific guidance, see the ADA position statement “Physical Activity/Exercise and Diabetes” (255).

Aerobic Activity

Aerobic activities encompass prolonged rhythmic activities using large muscle groups, such as walking, running, and cycling (255), and provide multiple cardiometabolic benefits for people with diabetes (241,247,259,260,292). Adults with diabetes are encouraged to engage in 150 min or more of moderate- to vigorous-intensity aerobic activity weekly (254,255). Aerobic activity bouts should last at least 10 min, with the goal of ~30 min/day or more most days of the week to accumulate at least 150 min per week (e.g., 30 min/day, 5 days per week). Daily exercise, or at least not allowing more than 2 days to elapse between exercise sessions, is recommended to decrease insulin resistance,

regardless of diabetes type (293,294). Over time, aerobic activities should progress in intensity, frequency, and/or duration to meet the recommended exercise volume. Though exercise at higher intensities can facilitate greater reductions in A1C (295), many adults may be unable or unwilling to participate in high-intensity exercise. In those cases, diabetes care professionals should encourage engagement in moderate exercise for the recommended duration for A1C reduction and other cardiometabolic benefits (241,259,292).

Resistance Activity

Resistance activity refers to movements working against an external force for the purpose of improving skeletal muscle strength and size, which typically includes free weight, weight machine, body weight, or resistance band movements to target the major muscle groups. Resistance training can improve glycemia, strength, hypertrophy, bone mineral density, and cardiometabolic health (296–298), and higher relative muscle mass is associated with better insulin sensitivity and lower diabetes risk (298). Adults with diabetes are encouraged to participate in 2–3 sessions/week of resistance exercise on non-consecutive days (254,255). Although it has been suggested that high-intensity resistance exercise training provides a greater benefit for glycemic management when compared with lower intensities (299), resistance training of any intensity is recommended to improve strength, balance, and the ability to engage in activities of daily living throughout the lifespan.

Flexibility and Balance Activities

Flexibility and balance activities serve to improve the range of motion around joints and reduce the risk of falls, respectively (254,255). Older adults with diabetes are encouraged to participate in flexibility and balance training 2–3 times/week. A variety of activities, including yoga, tai chi, and resistance training, can significantly improve A1C, flexibility, muscle strength, balance, and gait (254,300–304). Older adults in particular can benefit from flexibility and resistance training for maintenance of range of motion, strength, and balance (255,305,306) (Fig. 5.2).

High-Intensity Interval Training

High-intensity interval training (HIIT) involves short bursts of aerobic training performed

between 65% and 90% $\text{VO}_{2\text{peak}}$ (a measure of maximal aerobic capacity) or 75% and 95% heart rate peak for 10 s to 4 min with 12 s to 5 min of active or passive recovery. HIIT is a potentially time-efficient modality that can elicit significant physiologic and metabolic adaptations for individuals with type 1 and type 2 diabetes, including improvements in glycemia, β -cell function, central adiposity, and cardiac function (307–310).

When matched for energy expenditure, a 4-month HIIT-style walking program provided greater improvement in glycemia, physical fitness, and body composition compared with a continuous-walking program in people with type 2 diabetes (311). In contrast, a 12-week study in people with type 2 diabetes noted that HIIT cycling and continuous cycling programs, matched for energy expenditure, were not different in efficacy compared with each other or with a nonexercise control group (312). Both cycling protocols demonstrated improvements in aerobic capacity and A1C compared with baseline, but only continuous cycling showed improvements in cardiovascular response and visceral adiposity compared with baseline (312).

Because an acute bout of HIIT can lead to transient increases in postexercise hyperglycemia, individuals with type 1 diabetes may need to use bolus correction (313) and individuals with type 2 diabetes are encouraged to monitor blood glucose when starting HIIT (314). In type 1 diabetes, weeks of HIIT reduce A1C and insulin requirements and improve cardiometabolic risk profiles (310). Variability in glucose may occur with an increased risk in delayed hypoglycemia, so careful monitoring of glucose during and after HIIT is advised (310). Though HIIT may be an attractive mode of activity due to lowered required time commitment, individuals may be unable or unwilling to participate in activity at a high intensity. It may be of interest to intersperse HIIT within the broader activity plan with considerations for adequate recovery to prevent potential exercise-related injury (293).

Other Considerations

Exercise During Obesity Treatment

As many people with diabetes and obesity are often treated with obesity pharmacotherapy or metabolic surgery, additional emphasis on meeting physical activity guidelines, particularly muscle-strengthening activities, is warranted. Both obesity pharmacotherapy and metabolic surgery lead

to reductions in body weight, which often induces loss of both fat mass (adipose tissue) and fat-free mass (nonadipose tissues). This has raised concern about the loss of fat-free mass (particularly skeletal muscle and bone) and its potential long-term consequences on strength, physical function, resting energy expenditure, and the potential development or worsening of frailty, sarcopenia, and sarcopenic obesity. Engagement in exercise programs prior to and after metabolic surgery may support muscle mass and strength (315,316), reduce weight recurrence and promote better weight maintenance (317,318), and improve insulin sensitivity (319). Even modest increases in physical activity after metabolic surgery may contribute to greater fat mass loss and better retention of skeletal muscle mass (320). Though aerobic exercise may be beneficial, combined aerobic and resistance training may produce greater weight loss and better improvements in functional capacity, muscle mass, and strength compared with aerobic exercise alone (316). In addition to metabolic surgery, combining obesity pharmacotherapy with exercise programs may lead to some synergistic benefits. Combination of structured exercise training plus a GLP-1 RA has demonstrated improvements in β -cell function, glucose tolerance, and insulin sensitivity (321,322), reductions in abdominal adiposity and inflammation (323), and more favorable effects on body composition such as fat mass loss, fat-free mass maintenance, and reduced waist circumference (321–324). Conversely, a 16-week RCT reported that combined exercise training and GLP-1 RA therapy did not show synergistic benefits on cardiac function but rather showed potential inhibitory effects on left ventricular diastolic function in people with type 2 diabetes (325). However, this was a predefined sub-study that was not designed to detect changes in myocardial function. As data in this area for people with diabetes are emerging, more studies of longer duration and with newer-generation medications are needed.

Hypoglycemia

For individuals taking insulin and/or insulin secretagogues, physical activity may cause hypoglycemia if the medication dose or carbohydrate consumption is not adjusted for the exercise session and postbout effect on glucose. Individuals on these therapies may need to ingest carbohydrates if

Importance of 24-hour physical behaviors for type 2 diabetes

SITTING/BREAKING UP PROLONGED SITTING

- Limit sitting. Breaking up prolonged sitting (at least every 30 min) with short regular bouts of slow walking or simple resistance exercises can improve glucose metabolism.



STEPPING

- An increase of only 500 steps/day is associated with 2-9% decreased risk of cardiovascular morbidity and all-cause mortality.
- A 5- to 6-min brisk-intensity walk per day equates to ~4 years' greater life expectancy.



SLEEP

Aim for consistent, uninterrupted sleep, even on weekends.



Quantity - Long (>8 h) and short (<6 h) sleep durations negatively impact A1C.



Quality - Irregular sleep results in poorer glycemic levels, likely influenced by the increased prevalence of insomnia, obstructive sleep apnea, and restless leg syndrome in people with type 2 diabetes.



Chronotype - Evening chronotypes (i.e., night owl: go to bed late and get up late) may be more susceptible to inactivity and poorer glycemic levels than morning chronotypes (i.e., early bird: go to bed early and get up early).

SWEATING (MODERATE-TO-VIGOROUS ACTIVITY)

- Encourage ≥150 min/week of moderate-intensity physical activity (i.e., uses large muscle groups, rhythmic in nature) OR ≥75 min/week vigorous-intensity activity spread over ≥3 days/week, with no more than 2 consecutive days of inactivity. Supplement with two to three resistance, flexibility, and/or balance sessions.
- As little as 30 min/week of moderate-intensity physical activity improves metabolic profiles.



PHYSICAL FUNCTION

Physical function/ frailty/sarcopenia

- The frailty phenotype in type 2 diabetes is unique, often encompassing obesity alongside physical frailty, at an earlier age. The ability of people with type 2 diabetes to undertake simple functional exercises in middle-age is similar to that in those over a decade older.



STRENGTHENING

Resistance exercise (i.e., any activity that uses the person's own body weight or works against a resistance) also improves insulin sensitivity and glucose levels; activities like tai chi and yoga also encompass elements of flexibility and balance.



	Glucose/ insulin	Blood pressure	A1C	Lipids	Physical function	Depression	Quality of life
SITTING/BREAKING UP PROLONGED SITTING	↓	↓	↓	↓	↑	↓	↑
STEPPING	↓	↓	↓	↓	↑	↓	↑
SWEATING (MODERATE-TO-VIGOROUS ACTIVITY)	↓	↓	↓	↓	↑	↓	↑
STRENGTHENING	↓	↓	↓	↓	↑	↓	↑
ADEQUATE SLEEP DURATION	↓	↓	↓	↓	?	↓	↑
GOOD SLEEP QUALITY	↓	↓	↓	↓	?	↓	↑
CHRONOTYPE/CONSISTENT TIMING	↓	?	↓	?	?	↓	?

IMPACT OF PHYSICAL BEHAVIORS ON CARDIOMETABOLIC HEALTH IN PEOPLE WITH TYPE 2 DIABETES

- ↑ Higher levels of improvement (physical function, quality of life) ↓ Lower levels of improvement (glucose/insulin, blood pressure, A1C, lipids, depression)
 ? No data available
 ↑ Green arrows = strong evidence ↑ Yellow arrows = medium-strength evidence ↑ Red arrows = limited evidence

Figure 5.2—Importance of 24-h physical behaviors for type 2 diabetes. Adapted from Davies et al. (633).

pre-exercise glucose levels are <90 mg/dL (<5.0 mmol/L), depending on whether they are able to lower insulin doses during the workout (such as with an insulin pump

or reduced pre-exercise insulin dosage), the time of day exercise is done, and the intensity and duration of the activity (267). Due to heterogeneity of glycemic responses

to physical activity, the approach to carbohydrate intake and medication adjustments around the activity or exercise session should be individualized. More

guidance on carbohydrate intake and medication considerations can be found in the ADA position statement “Physical Activity/Exercise and Diabetes” (255). In some people with diabetes, hypoglycemia after exercise may occur and last for several hours due to increased insulin sensitivity. Hypoglycemia is not common in those who are not treated with insulin or insulin secretagogues, and no routine preventive measures for hypoglycemia are usually advised in these cases. Intense activities, such as HIIT, may actually raise glucose levels instead of lowering them, especially if pre-exercise glucose is elevated (267). Because of variation in glycemic response to exercise, people with diabetes should be taught to check blood glucose levels and/or monitor CGM values during and after exercise, how to understand the effect of exercise on glucose, about the potential prolonged effects (depending on intensity and duration), and how to manage and use AID technology around and during the time of exercise (266,326). See section 6, “Glycemic Goals, Hypoglycemia, and Hyperglycemic Crises,” for more information on hypoglycemia prevention and management.

Exercise in the Presence of Complications

See section 11, “Chronic Kidney Disease and Risk Management,” and section 12, “Retinopathy, Neuropathy, and Foot Care,” for more information on these long-term complications. A meta-analysis demonstrated that high versus low levels of physical activity were associated with lower CVD incidence and mortality (summary risk ratio 0.84 [95% CI 0.77–0.92], $n = 7$, and 0.62 [0.55–0.69], $n = 11$) and fewer microvascular complications (0.76 [0.67–0.86], $n = 8$). Dose-response meta-analyses showed that physical activity was associated with lower risk of diabetes-related complications even at lower activity levels (327).

Retinopathy

If proliferative diabetic retinopathy or severe nonproliferative diabetic retinopathy is present, then vigorous-intensity aerobic or resistance exercise may be contraindicated because of the risk of triggering vitreous hemorrhage or retinal detachment (328). Consultation with an ophthalmologist prior to engaging in an intense exercise plan may be appropriate.

Peripheral Neuropathy

Decreased pain sensation and a higher pain threshold in the extremities can result

in an increased risk of skin breakdown, infection, and Charcot joint destruction with some forms of exercise. Therefore, a thorough assessment should be done to ensure that neuropathy does not alter kinesthetic or proprioceptive sensation during physical activity, particularly in those with more severe neuropathy. Moderate-intensity walking may not lead to an increased risk of foot ulcers or reulceration in those with peripheral neuropathy who use proper footwear (329,330). In addition, 150 min/week of moderate exercise improved outcomes in people with prediabetic neuropathy (331). All individuals with peripheral neuropathy should wear proper footwear and examine their feet daily to detect lesions early. Anyone with a foot injury or open sore should be restricted to non-weight-bearing activities.

Autonomic Neuropathy

Autonomic neuropathy can increase the risk of exercise-induced injury or adverse events through decreased cardiac responsiveness to exercise, postural hypotension, impaired thermoregulation, impaired night vision due to impaired papillary reaction, and greater susceptibility to hypoglycemia (332). Cardiovascular autonomic neuropathy is also an independent risk factor for cardiovascular death and silent myocardial ischemia (333). Therefore, individuals with diabetic autonomic neuropathy should undergo cardiac investigation before beginning physical activity more intense than that to which they are accustomed.

Chronic Kidney Disease

Physical activity can acutely increase urinary albumin excretion. However, there is no evidence that vigorous-intensity exercise accelerates the rate of progression of CKD, and there appears to be no need for specific exercise restrictions for people with CKD in general (328).

SMOKING CESSATION: TOBACCO, E-CIGARETTES, AND CANNABIS

Recommendations

5.40 Ask people with diabetes routinely about the use of tobacco or vape products. **A** Advise complete avoidance of tobacco and vaping. **A** For individuals who use these products, provide or refer for combination treatment consisting of tobacco and/or vape product(s) cessation counseling and pharmacologic therapy. **A**

5.41 Advise people with type 1 diabetes **C** and those with other forms of diabetes at risk for diabetic ketoacidosis not to use recreational cannabis in any form. **E**

A causal link between cigarette smoking and diabetes has been well established for over a decade (334). Results from epidemiologic, case-control, and cohort studies support the causal link between cigarette smoking and multiple health risks that profoundly affect the morbidity and mortality of people with diabetes (334). People with diabetes who smoke and are exposed to second-hand smoke have a heightened risk of macrovascular complications (e.g., cardiovascular and peripheral vascular disease), microvascular complications (e.g., kidney disease and visual impairment), elevated A1C, and premature death compared with those who do not smoke and are not exposed to second-hand smoke (335,336). Findings from a systematic review and meta-analysis show a dose-response relation for current smoking and the risk for type 2 diabetes; this risk decreases as the time since quitting increases (337).

Routine (every visit with every person), thorough assessment of all types of tobacco use is essential to prevent tobacco product initiation and promote cessation. Evidence demonstrates quitting smoking reduces or reverses adverse health effects in addition to increasing life expectancy by up to 10 years (338). However, tobacco use among adults with chronic conditions remains higher than the general population, despite recent declines in smoking among middle-aged and older adults with diabetes (339). Numerous RCTs demonstrate the efficacy and cost-effectiveness of both intensive and brief counseling on smoking cessation, including the use of telephone quit lines and web-based interventions, in reducing tobacco use and maintaining abstinence from smoking (338,340). The World Health Organization recommends both counseling and pharmacologic therapy to assist with smoking cessation in nonpregnant adults (341). A secondary data analysis of the Evaluating Adverse Events in a Global Smoking Cessation Study (EAGLES), a randomized, double-blind, triple-dummy, placebo-controlled and active-controlled trial, found varenicline to be the most efficacious pharmacotherapy for people with diabetes compared