Applied Linear Algebra in Data Analysis

Positive Definite Matrices and Matrix Norms

Sivakumar Balasubramanian

Christian Medical College Vellore
Department of Bioengineering

Positive Definite Matrices

There are matrices that are positive in one sense.

Positive definite matrices

We know that $\mathbf{x}^\top \mathbf{x} > 0$ for all $\mathbf{x} \in \mathbb{R}^n$ and $\mathbf{x} \neq \mathbf{0}$.

What can we say about $\mathbf{x}^\top \mathbf{A}\mathbf{x}$, where $\mathbf{A} \in \mathbb{R}^{n \times n}$?

- Is it positive for all $\mathbf{x} \neq \mathbf{0}$? $\mathbf{I}$, $\begin{bmatrix} 3 & 0 \\ 0 & 2 \end{bmatrix}$
- Can it be zero for some $\mathbf{x} \neq \mathbf{0}$? $\begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}$
- Can it be negative some $\mathbf{x} \neq \mathbf{0}$? $\begin{bmatrix} -2 & 0 \\ 0 & -1 \end{bmatrix}$, $\begin{bmatrix} 1 & -4 \\ -4 & 1 \end{bmatrix}$

Any matrix $\mathbf{A}$ for which $\mathbf{x}^\top \mathbf{A}\mathbf{x} > 0$, for all $\mathbf{x} \neq \mathbf{0}$ is called a *positive definite matrix*.

Positive definite matrices are commonly encountered in practice: optimization, mechanics (mass matrix, stiffness matrix), stability analysis, co-variance matrices etc.

Positive definite matrices

Are the following matrices positive definite? $\begin{bmatrix} 2 & -6 \\ 2 & 1 \end{bmatrix}, \begin{bmatrix} 5 & -1 \\ 14 & 11 \end{bmatrix}$

Positive definite matrices

The idea of positive definiteness is intimately related to the problem of minimization of a function.

Consider the following function of a single variable $f(x)$. This function reaches a minimum at $x = 0$, when $\frac{df(x)}{dx}\Big|_{x=0} = 0$ and $\frac{d^2f(x)}{dx^2}\Big|_{x=0} > 0$. E.g.,

$$f(x) = 3x^2 \longrightarrow \frac{df(x)}{dx}\Big|_{x=0} = 0, \ \frac{d^2f(x)}{dx^2}\Big|_{x=0} = 3 > 0$$

Positive definite matrices

What about $f(x_1, x_2) = ax_1^2 + 2bx_1x_2 + cx_2^2$? We can extend the previous idea using partial derivatives.

$$\frac{\partial f(x_1, x_2)}{\partial x_1} = 0, \quad \frac{\partial f(x_1, x_2)}{\partial x_2} = 0, \quad \frac{\partial^2 f(x_1, x_2)}{\partial x_1^2} > 0, \quad \frac{\partial^2 f(x_1, x_2)}{\partial x_2^2} > 0$$

This is not enough. $\frac{\partial^2 f(x_1,x_2)}{\partial x_1 \partial x_2}$ must also be taken into account.

Are these functions positive for all x_1, x_2?

1. $x_1^2 + x_1x_2 + x_2^2$
2. $x_1^2 + 2x_1x_2 + x_2^2$
3. $x_1^2 + 3x_1x_2 + x_2^2$?

Positive definite matrices

We can rearrange $ax_1^2 + 2bx_1x_2 + cx_2^2$ in the following manner,

$$f(x_1, x_2) = ax_1^2 + 2bx_1x_2 + cx_2^2 = a\left(x_1 + \frac{b}{a}x_2\right)^2 + \left(c - \frac{b^2}{a}\right)x_2^2$$

$f(\bullet) > 0, \forall x_1, x_2 \neq 0$ when,

$$a > 0 \quad \text{and} \quad c - \frac{b^2}{a} > 0 \implies ac > b^2$$

$$\frac{\partial^2 f(x_1, x_2)}{\partial x_1^2} > 0 \quad \text{and} \quad \frac{\partial^2 f(x_1, x_2)}{\partial x_1^2}\frac{\partial^2 f(x_1, x_2)}{\partial x_2^2} > \left(\frac{\partial^2 f(x_1, x_2)}{\partial x_1 \partial x_2}\right)^2$$

Positive definite matrices

$f(\bullet)$ can be expressed as $\mathbf{x}^\top \mathbf{A}\mathbf{x}$, where $\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$, and $\mathbf{A} = \begin{bmatrix} a & b \\ b & c \end{bmatrix}$. $f(\mathbf{x}) > 0$ for all $\mathbf{x} \neq \mathbf{0}$, when $\mathbf{A}$ is positive definite.

$\mathbf{x}^\top \mathbf{A}\mathbf{x}$ is called a *quadratic form*. For a symmetric matrix $\mathbf{A}$,

$$\mathbf{x}^\top \mathbf{A}\mathbf{x} = \begin{bmatrix} x_1 & x_2 & \cdots & x_n \end{bmatrix} \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{12} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{1n} & a_{2n} & \cdots & a_{nn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \cdots \\ x_n \end{bmatrix} = \sum_{i=1}^{n} a_{ii} x_i^2 + 2 \sum_{i=1}^{n} \sum_{j=1, j \neq i}^{n} a_{ij} x_i x_j$$

Positive definite matrices

In general, $\mathbf{A} \in \mathbb{R}^{n \times n}$ is positive definite, if:

- The eigenvalues of $\mathbf{A}$ are all positive.
- The pivots (without row exchange) are all positive.

Show that any $\mathbf{A}$ is positive definite if the symmetric matrix $\mathbf{A} + \mathbf{A}^\top$ is positive definite.
Note: *This should explain why we have only been talking about symmetric matrices.*

Matrix Norms

How big is a matrix?

Matrix Norm

Since the set matrices $\mathbf{A} \in \mathbb{R}^{p \times q}$ form vector spaces, we can talk about norms of matrices, extending the idea of sizes and distances to spaces of matrices.

If we think of matrices a collection of pq scalars, then we can use the same approach as vectors,

$$\|\mathbf{A}\|_F = \left(\sum_{i=1}^{p} \sum_{j=1}^{q} |a_{ij}|^2 \right)^{\frac{1}{2}}$$

This is called the **Frobinius norm**.

Note that the Frobinius norm is the 2-norm equivalent.

Induced Matrix Norm

There are other norms defined for matrices that are very useful from the point of the view of linear transformation.

These are called **induced matrix norms**, that looks at how matrices map vectors from the range to domain spaces.

Let $\mathbf{A} \in \mathbb{R}^{p \times q} : \mathbb{R}^q \longrightarrow \mathbb{R}^p$, and $\mathbf{x} \in \mathbb{R}^q$, $\mathbf{y} \in \mathbb{R}^p$

$$\|\mathbf{y}\| = \|\mathbf{A}\mathbf{x}\| \leq C \|\mathbf{x}\|, \quad \forall \mathbf{x} \in \mathbb{R}^q, \quad C \geq 0$$

C is the maximum factor by which $\mathbf{A}$ amplifies the vector $\mathbf{x}$.

The induced norm of a matrix is defined as,

$$\|\mathbf{A}\|_p = \max_{\mathbf{x}} \frac{\|\mathbf{A}\mathbf{x}\|_p}{\|\mathbf{x}\|_p} = \max_{\|\mathbf{x}\|_p = 1} \|\mathbf{A}\mathbf{x}\|_p$$

Induced Matrix Norm

Consider a matrix $\mathbf{A} = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} = \begin{bmatrix} \mathbf{a}_1 & \mathbf{a}_2 \end{bmatrix} = \begin{bmatrix} \tilde{\mathbf{a}}_1^\top \\ \tilde{\mathbf{a}}_2^\top \end{bmatrix}$.

$\|\mathbf{A}\|_1 = \max_i \|\mathbf{a}_i\|_1$



$\|\mathbf{A}\|_2 = \sqrt{\lambda_{\max}}$



$\|\mathbf{A}\|_\infty = \max_i \|\tilde{\mathbf{a}}_i\|_1$