

# Applied Linear Algebra in Data Analysis

Tutorial

Sivakumar Balasubramanian

## CONTENTS

1	Concepts in Vector Spaces	2
2	Matrices	3
3	Orthogonality	4
4	Matrix Inverses	5

## 1 CONCEPTS IN VECTOR SPACES

---

1. Which of the following sets forms a vector space?

- a)  $\{\mathbf{x} \mid x_1, x_2 \in \mathbb{R} \text{ and } a_1 x_1 + a_2 x_2 = 0\}$ , where  $a_1, a_2 \in \mathbb{R}$  are fixed constants.
- b)  $\{\mathbf{x} \mid \mathbf{x} \in \mathbb{R}^n \text{ and } \mathbf{a}^\top \mathbf{x} = b\}$ , where  $\mathbf{a} \in \mathbb{R}^n$  and  $b \in \mathbb{R}$  are fixed constants.
- c)  $\{\mathbf{x} \mid \mathbf{x} \in \mathbb{R}^n \text{ and } \mathbf{x}^\top \mathbf{x} = 1\}$ .
- d)  $\{(x[0], x[1], x[2], \dots, x[N-1]) \mid x[i] \in \mathbb{R}, 0 \leq i < N\}$ .

(The set of all real-valued time-domain signals of length  $N$ .  $x[i]$  is the value of the signal at time instant  $i$ .)

2. Consider the vector space of polynomials of order  $n$  or less.

$$\mathcal{P} = \left\{ \sum_{k=0}^n a_k x^k \mid a_k \in \mathbb{R} \right\}, \text{ where, } x \in [0, 1]$$

Show that polynomials of order strictly lower than  $n$  form subspaces of  $\mathcal{P}$ .

3. Is the following function a valid norm of the vector space  $\mathcal{P}$ ?

$$\|\mathbf{p}(x)\| = \sqrt{\sum_{k=0}^n a_k^2}, \quad \mathbf{p} = \sum_{k=0}^n a_k x^k \in \mathcal{P}$$

4. Consider the following function, which is often called the *zero-norm* of a vector  $\mathbf{x} \in \mathbb{R}^n$ .

$$\|\mathbf{x}\|_0 = \sum_{i=1}^n \mathbb{I}(x_i \neq 0), \text{ where, } \mathbb{I}(A) = \begin{cases} 1 & A \text{ is true.} \\ 0 & A \text{ is false.} \end{cases}$$

Is the *zero-norm*, which is often used for quantifying the *sparsity* of a vector, a proper norm?

5. Is the following set of vectors linear independent?

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ -1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 0 \\ -1 \end{bmatrix} \right\}$$

What is the span of this set? Does this set form the basis for its span? Does it form an orthonormal basis?

## 2 MATRICES

---

1. Consider the following matrices:

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -1 & 0 \\ 0 & 2 & -2 & 1 \\ -3 & 1 & 1 & 3 \end{bmatrix} \quad \text{and} \quad \mathbf{B} = \begin{bmatrix} 2 & 1 & 1 \\ 1 & -1 & 1 \\ 3 & 2 & 1 \\ 1 & 2 & 1 \end{bmatrix}$$

Find the product of the two matrices  $\mathbf{C} = \mathbf{AB}$  using the four views of matrix multiplication.

If we change  $b_{23} = 0$ . Can you compute the new matrix  $\mathbf{C}$  without performing the matrix multiplication?

If we increase the value of the elements of the 3<sup>rd</sup> column of  $\mathbf{A}$  by 1, how can we compute the new  $\mathbf{C}$  without performing the matrix multiplication?

If we insert a new row  $\mathbf{1}^\top$  in  $\mathbf{A}$  after the 2<sup>nd</sup> row, how can we compute the new  $\mathbf{C}$  without performing the matrix multiplication?

2. Show that the matrix product  $\mathbf{ABC}$  can be written as a weighted sum of the outer products of the columns of  $\mathbf{A}$  and rows of  $\mathbf{C}$ , with the weights coming from the matrix  $\mathbf{B}$ .
3. Prove the following for the matrices  $\mathbf{A}_1, \mathbf{A}_2, \mathbf{A}_3, \dots, \mathbf{A}_n$ .

$$(\mathbf{A}_1, \mathbf{A}_2, \mathbf{A}_3, \dots, \mathbf{A}_n)^\top = \mathbf{A}_n^\top \mathbf{A}_{n-1}^\top \dots \mathbf{A}_2^\top \mathbf{A}_1^\top$$

4. **Nilpotent matrices.** Show that a strictly triangular matrix  $\mathbf{A} \in \mathbb{R}^{n \times n}$ ,  $\mathbf{A}^n = \mathbf{0}$ .
5. **Matrix Inversion Lemma.** Consider an invertible matrix  $\mathbf{A}$ . The matrix  $\mathbf{A} + \mathbf{uv}^\top$  is invertible if and only if the two vectors  $\mathbf{u}, \mathbf{v} \neq \mathbf{0}$ , and  $\mathbf{v}^\top \mathbf{A}^{-1} \mathbf{u} \neq -1$ . Then, the inverse is given by,

$$(\mathbf{A} + \mathbf{uv}^\top)^{-1} = \mathbf{A}^{-1} - \frac{\mathbf{A}^{-1} \mathbf{uv}^\top \mathbf{A}^{-1}}{1 + \mathbf{v}^\top \mathbf{A}^{-1} \mathbf{u}}$$

6. Prove that  $\text{tr}(\mathbf{AB}) = \text{tr}(\mathbf{BA})$ , where  $\mathbf{A} \in \mathbb{R}^{n \times d}$  and  $\mathbf{B} \in \mathbb{R}^{d \times n}$ .
7. Effect of matrix operation on matrix rank. Let  $\mathbf{A} \in \mathbb{R}^{n \times d}$  and  $\mathbf{B} \in \mathbb{R}^{d \times n}$ , with ranks  $a$  and  $b$  respectively. What is the rank of the following matrices?
  - a)  $\mathbf{A} + \mathbf{B}$
  - b)  $\mathbf{AB}$
8. Show that the rank  $(\mathbf{AB}) = \text{rank}(\mathbf{A})$ , when  $\mathbf{B}$  is square and full rank.
9. Let  $\mathbf{A}, \mathbf{B} \in \mathbb{R}^{n \times n}$ ,  $\mathbf{AB}$  is non-singular if and only if both  $\mathbf{A}$  and  $\mathbf{B}$  are non-singular.
10. Let  $\mathbf{A}$  is a full rank matrix. Show that the *Gram matrix* of the column space,  $\mathbf{A}^\top \mathbf{A}$  is invertible.

### 3 ORTHOGONALITY

---

1. If  $\mathbf{A}$  is an orthogonal matrix, show that  $\mathbf{A}^{-1} = \mathbf{A}^\top$ .
2. If  $\mathbf{P}_S$  is the orthogonal projection matrix onto the subspace  $S$ , then what is the corresponding orthogonal projection matrix onto  $S^\perp$  – the orthogonal complement of  $S$ ?
3. Let  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ . Let  $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n\}$  be an orthonormal basis for  $\mathbb{R}^n$ . Show that the following holds,

$$\mathbf{x}^\top \mathbf{y} = \sum_{i=1}^n (\mathbf{x}^\top \mathbf{u}_i) \cdot (\mathbf{u}_i^\top \mathbf{y})$$

4. Consider the following set of vectors,  $S = \{\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3, \dots, \mathbf{a}_n\}$ , where  $\mathbf{a}_i \in \mathbb{R}^n$ . The set  $S$  is linearly independent. Find the orthogonal components of a vector  $\mathbf{b} \in \mathbb{R}^n$  in the subspace spanned by the sets of vectors  $S_1 = \{\mathbf{a}_i\}_{i=1}^m$  and  $S_1^\perp$ .
5. Consider the set of  $n \times n$  orthogonal matrices,

$$\mathcal{Q} = \left\{ \mathbf{Q} \mid \mathbf{Q} \in \mathbb{R}^{n \times n}, \mathbf{Q}^\top \mathbf{Q} = \mathbf{Q} \mathbf{Q}^\top = \mathbf{I}_n \right\}$$

Is this set a subspace of  $\mathbb{R}^{n \times n}$ ?

Show that the set is closed under matrix multiplication.

6. Consider the linear map,  $\mathbf{y} = \mathbf{A}\mathbf{x}$ , such that  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$  and  $\mathbf{A} \in \mathbb{R}^{n \times n}$ . Let us assume that  $\mathbf{A}$  is full rank. What conditions must  $\mathbf{A}$  satisfy for the following statements to be true,
  - a)  $\|\mathbf{y}\|_2 = \|\mathbf{x}\|_2$ , for all  $\mathbf{x}, \mathbf{y}$  such that  $\mathbf{y} = \mathbf{A}\mathbf{x}$ .
  - b)  $\mathbf{y}_1^\top \mathbf{y}_2 = \mathbf{x}_1^\top \mathbf{x}_2$ , for all  $\mathbf{x}_1, \mathbf{x}_2, \mathbf{y}_1, \mathbf{y}_2$  such that  $\mathbf{y}_1 = \mathbf{A}\mathbf{x}_1$  and  $\mathbf{y}_2 = \mathbf{A}\mathbf{x}_2$ .

**Note:** A linear map  $\mathbf{A}$  with the aforementioned properties preserves lengths and angle between vectors. Such maps are encountered in rigid body mechanics.

## 4 MATRIX INVERSES

---

1. Find a left inverse for the matrix  $\mathbf{A} = \begin{bmatrix} 1 & 1 \\ -1 & 1 \\ 0 & 1 \end{bmatrix}$ . Find the set of all possible left inverses.
2. Show that if the product of two square  $d \times d$  matrices  $\mathbf{A}$  and  $\mathbf{B}$  is the identity matrix  $\mathbf{I}$ , then  $\mathbf{BA} = \mathbf{I}$ .
3. Consider an upper triangular matrix  $\mathbf{R} \in \mathbb{R}^{n \times n}$ . We are interested in solving the following set of  $n$  linear equations,

$$\mathbf{R}\mathbf{x} = \mathbf{e}_i$$

$\mathbf{x} = [x_1 \ x_2 \ x_3 \ \dots \ x_n]^\top$  is the solution to the above equation. Show that  $x_{i+1} = x_{i+2} = \dots = x_n = 0$ .

Show that the solution to this equation is equal to the  $i^{\text{th}}$  column of the inverse of  $\mathbf{R}$ .

4. Find the pseudo-inverse of the matrix  $\mathbf{A} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 1 & 1 \end{bmatrix}$ . Show that the matrix  $\mathbf{AA}^\dagger$  is the orthogonal projection matrix onto the column space of  $\mathbf{A}$ .