

1. Limiting density of interpolation nodes:

- Suppose x_0, x_1, \dots, x_n are $n+1$ points equally spaced from -1 to 1 . If $-1 \leq a < b \leq 1$, what fraction of points lie in the interval $[a, b]$ in the limit as $n \rightarrow \infty$?

Solution: $\frac{b-a}{2n}$ ■

- Give the analogous expression for the case where x_0, x_1, \dots, x_n are Chebyshev nodes.

Solution: The Chebyshev nodes are given by $\cos\left(\frac{j\pi}{n}\right)$, where $j \in \{0, 1, 2, \dots, n\}$. Hence, the number of points is $\frac{n}{\pi} (\lceil \arccos(a) \rceil - \lceil \arccos(b) \rceil) = \frac{(\lceil \arccos(a) \rceil - \lceil \arccos(b) \rceil)}{\pi}$ ■

- Obtain in the limit as $n \rightarrow \infty$, the density of Chebyshev points at $x \in (-1, 1)$. The density is defined as

$$\rho(x) = \lim_{\epsilon \rightarrow 0} \lim_{n \rightarrow \infty} \frac{\text{Number of Chebyshev nodes in the interval } (x-\epsilon, x+\epsilon)}{2\epsilon n}$$

Solution:

$$\lim_{\epsilon \rightarrow 0} \frac{1}{\pi} \frac{\arccos(x-\epsilon) - \arccos(x+\epsilon)}{2\epsilon} = \frac{1}{\pi \sqrt{1-x^2}}$$

■

2. Recall that we interpolated a function $f(x)$ on a set of $n+1$ distinct points

$$x_0 < x_1 < \dots < x_n$$

using Lagrange polynomials as $p_n(x) = \sum_{j=0}^n y_j L_j(x)$, where y_j is the value of the function $f(x)$ at x_j and $L_j(x)$ is the Lagrange polynomial taking a value of 1 at $x = x_j$ and 0 elsewhere. To obtain the derivative of $p_n(x)$ at the data points x_j , we seek a matrix D such that

$$Dy = p'_n$$

where y is the vector of function values and p'_n is a vector of derivatives at x_j .

- Show that $d_{jk} = \left. \frac{dL_k(x)}{dx} \right|_{x=x_j}$

Solution: We have $\frac{d}{dx}(p_n(x)) = \sum_{j=0}^n y_j \frac{dL_j(x)}{dx} \implies \left. \frac{d}{dx}(p_n(x)) \right|_{x=x_k} = \sum_{j=0}^n y_j \left. \frac{dL_j(x)}{dx} \right|_{x=x_k}$ ■

- Obtain an expression for d_{jk} in terms of x_i 's.

Solution: $d_{jk} = \frac{\alpha_k}{\alpha_j (x_j - x_k)}$ for $j \neq k$, where $\alpha_j = \left(\prod_{i=0, i \neq j}^n (x_j - x_i) \right)^{-1}$ ■

3. A general Pade type boundary scheme (at $i=0$) for the first derivative can be written as

$$f'_0 + \alpha f'_1 = \frac{af_0 + bf_1 + cf_2 + df_3}{h}$$

- Obtain a, b, c, d in terms of α , if we would like the scheme to be third order accurate.

Solution: $a = -\frac{11+2\alpha}{6}$, $b = \frac{6-\alpha}{2}$, $c = \frac{2\alpha-3}{2}$, $d = \frac{2-\alpha}{6}$ ■

- What value of α you would pick and why?

Solution: Pick $d=0$, so that the stencil would be more compact, i.e., pick $\alpha=2$. ■

- Find all the coefficients if such a scheme would be fourth-order accurate.

4. The integral $\int_0^\pi \sin(x)dx$ is approximated using Trapezoidal rule with $N + 1$ points and a grid spacing of $\frac{\pi}{N}$, say T_n is the approximation.

- Obtain an expression for T_N as a function of N .

Solution: The grid points are $x_j = \frac{j\pi}{N}$. Hence, we have

$$T_N = \frac{\pi}{N} \sum_{j=1}^{N-1} \sin\left(\frac{j\pi}{N}\right) = \frac{\pi}{N} \frac{\sin((N-1)\pi/(2N))}{\sin(\pi/(2N))} \sin\left(\frac{\pi}{2}\right) = \frac{\pi}{N} \cot(\pi/(2N))$$

■

- Compute the value of $\lim_{N \rightarrow \infty} T_N$ and check if it gives the exact integral.

Solution: Limiting value of T_N is 2, which indeed matches with the integral. ■

5. One would like to derive a Runge Kutta method of order 3. Recall that RK3 method is given by

$$y_{n+1} = y_n + h \sum_{i=1}^3 \gamma_i k_i$$

$$k_1 = f(t_n, y_n)$$

$$k_2 = f(t_n + \alpha_1 h, y_n + h\beta_{21}k_1)$$

$$k_3 = f(t_n + \alpha_2 h, y_n + h\beta_{31}k_1 + h\beta_{32}k_2)$$

Obtain the constraints on γ_i, α_i and β_{ij} .

6. Obtain the stability region and accuracy for the weighted trapezoidal rule:

$$y_{n+1} = y_n + h(\alpha f(t_n, y_n) + (1 - \alpha) f(t_{n+1}, y_{n+1}))$$

where $\alpha \in (0, 1)$.

7. We have access to a uniform random number generator on the interval $[0, 1]$. Let's call this function $\text{rand}()$, i.e., calling $x = \text{rand}()$ will give us a number on the interval $[0, 1]$ generated out of a uniform distribution on the interval $[0, 1]$. Use this random number generating function to generate

- A Bernoulli random variable Z , i.e., the random variable Z takes only two values 0 or 1. It takes value 0 with a probability p and a value 1 with a probability $1 - p$.

Solution: Let U be the uniform random variable generated by $\text{rand}()$ on the interval $[0, 1]$. Consider the random variable Z which takes value 0, when $\text{rand}() \leq p$ and value 1, when $\text{rand}() > p$. Now probability that Z takes 0 is same as the probability that a uniform random variable on the interval $[0, 1]$ is less than p , which is nothing but p , i.e.,

$$\mathbb{P}(Z = 0) = \mathbb{P}(U \leq p) = p$$

$$\mathbb{P}(Z = 1) = \mathbb{P}(U > p) = 1 - p$$

■

- A Binomial random variable Y with parameters n and p , i.e., a random variable which takes values belonging to $\{0, 1, 2, \dots, n\}$, whose probability mass function is given by

$$P_Y(y = k) = \binom{n}{k} p^k (1 - p)^{n-k} \quad \text{for } k \in \{0, 1, 2, \dots, n\}$$

Solution: A binomial random variable with parameters n and p is a sum of n Bernoulli random variable with parameter p . Hence, generate n Bernoulli random variables Z_1, Z_2, \dots, Z_n . Set the binomial random variable as $Z_1 + Z_2 + \dots + Z_n$. ■

8. We have access to a uniform random number generator on the interval $[0, 1]$. Let's call this function $\text{rand}()$, i.e., calling $x = \text{rand}()$ will give us a number on the interval $[0, 1]$ generated out of a uniform distribution on the interval $[0, 1]$. Describe a procedure to use this random number generating function to obtain the value of the integral

$$\int_1^\infty \frac{dx}{1+x^2}$$

Solution: Use a change of variables $x = 1/t$. We then see that the integral becomes

$$\int_0^1 \frac{dt}{1+t^2}$$

Now generate n uniform random numbers on the interval $[0, 1]$, say $\{t_i\}_{i=1}^n$, and approximate the integral as

$$\frac{\sum_{i=1}^n \frac{1}{1+t_i^2}}{n}$$

■