$\mathsf{Module}\ \mathsf{D}$

- Section D.1
- Section D.1
 Section D.2
- Section D.3
- Section D.4
- Section D.
- Section D 6

Module D: Discontinuous functions in ODEs

Section D.1

Section D.1 Section D.2

Section D.3

Section D.4

Section D.

How can we solve and apply ODEs involving functions that are not continuous?

Section D.3 Section D.3 Section D.4

Section |

At the end of this module, students will be able to...

- D1. Laplace Transform. ...compute the Laplace transform of a function
- **D2. Discontinuous ODEs.** ...solve initial value problems for ODEs with discontinuous coefficients
- D3. Modeling non-smooth motion. ...model the motion of an object undergoing discontinuous acceleration
- **D4. Modeling non-smooth oscillators.** ...model mechanical oscillators undergoing discontinuous acceleration

Section D.1 Section D.2 Section D.3 Section D.4

Readiness Assurance Outcomes

Before beginning this module, each student should be able to...

• Partialf ractions

Section D.1 Section D.2

Section D.3 Section D.4

Section D. Section D.

The following resources will help you prepare for this module.

• TODO

Module F

Section D.1

Section D.1 Section D.2

Section D.

Section D.4

Section D.6

Module D Section 1

. . . .

viouule D

Section D.1 Section D.2

Section D.3


Section D.4

Section D.5

Section

Activity D.1.1 (\sim 10 min)

A 1 kg mass is hung from a spring with spring constant $k=1\ N/m$. The mass is at rest, when it is hit with a hammer imparting 3J of energy.



Module

Section D.1

Section D.1 Section D.2


Section D.3

C--ti-- D.F

Section I

Activity D.1.1 (\sim 10 min)

A 1 kg mass is hung from a spring with spring constant $k=1~\mathrm{N/m}$. The mass is at rest, when it is hit with a hammer imparting 3J of energy.



Part 1: Draw a graph of the kinetic energy in the system with respect to time.


Section D.

Section D.4

Section I

Activity D.1.1 (\sim 10 min)

A 1 kg mass is hung from a spring with spring constant $k=1\ N/m$. The mass is at rest, when it is hit with a hammer imparting 3J of energy.



Part 1: Draw a graph of the kinetic energy in the system with respect to time.

Part 2: Write an initial value problem modelling this system.

Definition D.1.2

The **Dirac delta distribution** $\delta(t)$ models the application of instantaneous force. **It is not a function**, but makes sense in definite integrals:

If a, b is any open interval containing 0, then

$$\int_a^b f(t)\delta(t)dt = f(0)$$

for any function f(t) that is continuous around 0.

Vlodule

Section D.1 Section D.2

Section D.3

Section D.4

Section D.

Section

Definition D.1.3

The **unit impulse function** u(t) is given by

$$u(t) = \begin{cases} 1 & t > 0 \\ 0 & t < 0. \end{cases}$$

Note that $u(s) = \int_{-\infty}^{s} \delta(t) dt$; in this fuzzy sense, δ is the derivative of u(t) (which is not differentiable everywhere!)

Section D.1 Section D.2

Section D.3

Section D.4

Section D

Section

Activity D.1.4 (\sim 10 min)

Try to solve the IVP

$$y'' + y = \delta(t)$$

Where does our existing technique break down?

Observation D.1.5

To get around this difficulty, we will apply an **integral transform** called the **Laplace Transform** to our ODE.

- We want to use a definite integral to handle things like δ , which we can only understand via a definite integral.
- \bullet Since we are focused on IVPs, we can integrate starting at 0, but need to go to ∞
- But now we need to worry about convergence—thus we will multiply by a suitable function that decays fast enough to make most functions converge.

Aodule D

Section D.1

Section D.2

Section D.3

Section D.4

Section [

Section I

Activity D.1.6 (\sim 5 min)

Arrange the following functions in order of how fast they decay to zero in the limit at infinity:

- (A) x^{-n} for a positive integer n
- (B) e^{-ax} for a positive integer a
- (C) $\frac{1}{\ln(ax)}$ for a positive integer a
- (D) $\frac{1}{\ln(x^n)}$ for a positive integer n

Definition D.1.7

The **Laplace Transform** of a function f(t) is the function

$$\mathcal{L}{f}(s) = \int_0^\infty e^{-st} f(t) dt.$$

Note that the Laplace transform turns a function of t into a function of s.

Moreover, \mathcal{L} is linear: $\mathcal{L}\{f+g\} = \mathcal{L}\{f\} + \mathcal{L}\{g\}$, and $\mathcal{L}\{cf\} = c\mathcal{L}\{f\}$ for constants c.

Section D.1

- Section D.2 Section D.3
- Section D.4

Activity D.1.8 (\sim 5 min)

Recall that

$$\mathcal{L}\{f\}(s) = \int_0^\infty e^{-st} f(t) dt.$$

Module E

Section D.1

Section D.2 Section D.3

Section D.4

Section D.

Section I

Recall that

Activity D.1.8 (\sim 5 min)

$$\mathcal{L}{f}(s) = \int_0^\infty e^{-st} f(t) dt.$$

Part 1: Compute $\mathcal{L}(\delta(t))$

Section D.1

Activity D.1.8 (\sim 5 min)

Recall that

$$\mathcal{L}{f}(s) = \int_0^\infty e^{-st} f(t) dt.$$

Part 1: Compute $\mathcal{L}(\delta(t))$

Part 2: If a > 0, compute $\mathcal{L}\{\delta(t-a)\}$

Section D.1

Section D.2 Section D.3

Section D.4

Activity D.1.9 (\sim 5 min)

Recall that

$$\mathcal{L}\{f\}(s) = \int_0^\infty e^{-st} f(t) dt.$$

Module E

Section D.1 Section D.2 Section D.3 Activity D.1.9 (~5 min)

Section D.4
Section D.5
Recall that

$$\mathcal{L}\{f\}(s) = \int_0^\infty e^{-st} f(t) dt.$$

Part 1: Compute $\mathcal{L}\{e^t\}$

Section D.1

Section D.2 Section D.3

Section D.4

Activity D.1.9 (\sim 5 min)

Recall that

$$\mathcal{L}{f}(s) = \int_0^\infty e^{-st} f(t) dt.$$

Part 1: Compute $\mathcal{L}\{e^t\}$

Part 2: If a > 0, compute $\mathcal{L}\{e^{at}\}$

Activity D.1.10 (\sim 15 min) Section D.1

Section D.2 Recall that Section D.3 Section D.4

$$\mathcal{L}{f}(s) = \int_0^\infty e^{-st} f(t) dt.$$

Section D.1 Section D.2

Section D.3 Section D.4

Activity D.1.10 (\sim 15 min)

Recall that

$$\mathcal{L}{f}(s) = \int_0^\infty e^{-st} f(t) dt.$$

Part 1: Compute $\mathcal{L}\{1\}$

Section D.3 Section D.4

Activity D.1.10 (\sim 15 min) Section D.1 Section D.2

Recall that

$$\mathcal{L}{f}(s) = \int_0^\infty e^{-st} f(t) dt.$$

Part 1: Compute $\mathcal{L}\{1\}$ Part 2: Compute $\mathcal{L}\{t\}$

Section D.1 Section D.2

Section D.3

Section D.4

Activity D.1.10 (\sim 15 min)

Recall that

$$\mathcal{L}{f}(s) = \int_0^\infty e^{-st} f(t) dt.$$

Part 1: Compute $\mathcal{L}\{1\}$

Part 2: Compute $\mathcal{L}\{t\}$

Part 3: Compute $\mathcal{L}\{t^2\}$

Section D.1

- Section D.3
- Section D.2
- Section D.4

Activity D.1.10 (\sim 15 min)

Recall that

$$\mathcal{L}{f}(s) = \int_0^\infty e^{-st} f(t) dt.$$

- Part 1: Compute $\mathcal{L}\{1\}$
- Part 2: Compute $\mathcal{L}\{t\}$
- Part 3: Compute $\mathcal{L}\{t^2\}$
- Part 4: Compute $\mathcal{L}\{t^3\}$

Section D.1

Section D.2 Section D.3

Section D.4

Activity D.1.10 (\sim 15 min)

Recall that

$$\mathcal{L}{f}(s) = \int_0^\infty e^{-st} f(t) dt.$$

Part 1: Compute $\mathcal{L}\{1\}$

Part 2: Compute $\mathcal{L}\{t\}$

Part 3: Compute $\mathcal{L}\{t^2\}$

Part 4: Compute $\mathcal{L}\{t^3\}$

Part 5: Compute $\mathcal{L}\{t^4\}$

M			

Section D 1

Section D.2

Section D.

Section D.

Section D.6

Module D Section 2

Observation D.2.1

Last week, we encountered the **Laplace Transform** of a function f(t):

$$\mathcal{L}{f}(s) = \int_0^\infty e^{-st} f(t) dt.$$

Recall that the Laplace transform turns a function of t into a function of s.

Moreover, \mathcal{L} is linear: $\mathcal{L}\{f+g\} = \mathcal{L}\{f\} + \mathcal{L}\{g\}$, and $\mathcal{L}\{cf\} = c\mathcal{L}\{f\}$ for constants c.

Observation D.2.2

We computed a few Laplace Transforms:

•
$$\mathcal{L}\{\delta(t-a)\}=e^{-as}$$
 for any $a>0$.

•
$$\mathcal{L}\{e^{at}\}=\frac{1}{s-a}$$

•
$$\mathcal{L}\{t^n\} = \frac{n!}{s^{n+1}}$$
 for any positive integer n .

•
$$\mathcal{L}\{1\} = \frac{1}{s}$$

Section D.1 Section D.2

Section D.3

Section D.4

Activity D.2.3 (\sim 10 min)

Recall

$$\mathcal{L}\{f\}(s) = \int_0^\infty e^{-st} f(t) dt.$$

Module D

Section D.1 Section D.2

Section D.3

Section D.4

Section I

Activity D.2.3 (\sim 10 min)

Recall

$$\mathcal{L}{f}(s) = \int_0^\infty e^{-st} f(t) dt.$$

Part 1: Compute $\mathcal{L}\{\sin(t)\}$.

Activity D.2.3 (\sim 10 min)

Recall

$$\mathcal{L}{f}(s) = \int_0^\infty e^{-st} f(t) dt.$$

Part 1: Compute $\mathcal{L}\{\sin(t)\}$.

Part 2: Compute $\mathcal{L}\{\cos(t)\}$.

So now our list of Laplace transforms is:

•
$$\mathcal{L}\{\delta(t-a)\}=e^{-as}$$
 for any $a>0$.

•
$$\mathcal{L}\{e^{at}\}=\frac{1}{s-a}$$

•
$$\mathcal{L}\{t^n\} = \frac{n!}{s^{n+1}}$$
 for any positive integer n .

•
$$\mathcal{L}\{1\} = \frac{1}{s}$$

•
$$\mathcal{L}\{\sin(t)\}=\frac{1}{s^2+1}$$

•
$$\mathcal{L}\{\cos(t)\} = \frac{s}{s^2+1}$$

Activity D.2.5 (\sim 10 min)

Suppose we want to apply the Laplace transform to an IVP: we will need to know how $\mathcal{L}\{y'\}$ is related to $\mathcal{L}\{y\}$. Recall

$$\mathcal{L}{f}(s) = \int_0^\infty e^{-st} f(t) dt.$$

Activity D.2.5 (\sim 10 min)

Suppose we want to apply the Laplace transform to an IVP: we will need to know how $\mathcal{L}\{y'\}$ is related to $\mathcal{L}\{y\}$. Recall

$$\mathcal{L}{f}(s) = \int_0^\infty e^{-st} f(t) dt.$$

Part 1: Use integration by parts to relate $\mathcal{L}\{y'\}$ to $\mathcal{L}\{y\}$.

Activity D.2.5 (\sim 10 min)

Suppose we want to apply the Laplace transform to an IVP: we will need to know how $\mathcal{L}\{y'\}$ is related to $\mathcal{L}\{y\}$. Recall

$$\mathcal{L}\{f\}(s) = \int_0^\infty e^{-st} f(t) dt.$$

Part 1: Use integration by parts to relate $\mathcal{L}\{y'\}$ to $\mathcal{L}\{y\}$.

Part 2: Use integration by parts (and the fact that $\mathcal{L}\{y'\} = s\mathcal{L}\{y\} - y(0)$) to relate $\mathcal{L}\{y''\}$ to $\mathcal{L}\{y\}$.

Observation D.2.6

We have

$$\mathcal{L}{y'} = s\mathcal{L}{y} - y(0)$$

$$\mathcal{L}{y''} = s^2\mathcal{L}{y} - sy(0) - y'(0)$$

This allows us to easily rewrite expressions like ay'' + by' + cy in terms of $\mathcal{L}\{y\}$.

Section D.2 Section D.3

Section D.4

Activity D.2.7 (\sim 10 min)

Consider the simple IVP

$$y'' + y = \delta(t),$$
 $y(0) = 0, y'(0) = 0.$

Activity D.2.7 (\sim 10 min)

Consider the simple IVP

$$y'' + y = \delta(t),$$
 $y(0) = 0, y'(0) = 0.$

Part 1: Apply the Laplace transform to this IVP, and simplify. Solve for $\mathcal{L}\{y\}$.

Activity D.2.7 (\sim 10 min)

Consider the simple IVP

$$y'' + y = \delta(t),$$
 $y(0) = 0, y'(0) = 0.$

Part 1: Apply the Laplace transform to this IVP, and simplify. Solve for $\mathcal{L}\{y\}$.

Part 2: Find a function
$$y$$
 satisfying $\mathcal{L}\{y\} = \frac{1}{s^2+1}$. We write $y = \mathcal{L}^{-1}\left\{\frac{1}{s^2+1}\right\}$.

Activity D.2.8 (\sim 15 min) Solve the IVP

$$y'' + y = \delta(t),$$
 $y(0) = 1, y'(0) = 2.$

Module F

C. ... D.1

Section D.1 Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Module D Section 3

Observation D.3.1

To solve an IVP using Laplace transforms:

- 1) Apply \mathcal{L} to the ODE. Use the initial condition(s) in computing $\mathcal{L}\{y'\}$, $\mathcal{L}\{y''\}$, etc.
- 2) Solve for $\mathcal{L}\{y\}$.
- 3) Take the inverse transform (using a table) to find the solution y.

Module D Section D.1

Section D.2 Section D.3

Section D.s

Section D.4

Section

Activity D.3.2 (\sim 5 min)

Compute $\mathcal{L}^{-1}\left\{\frac{1}{s+5}\right\}$.

- (a) u(t-5)
- (b) $\delta(t-5)$
- (c) e^{5t}
- (d) e^{-5t}

Section D.2 Section D.3

Section D.4

Activity D.3.3 (\sim 5 min)

Compute
$$\mathcal{L}^{-1}\left\{\frac{e^{-10s}}{s}\right\}$$
.

(a)
$$u(t-10)$$

(b)
$$\delta(t - 10)$$

(c)
$$u(t-10)e^{-t}$$

(d)
$$\delta(t-10)e^{-t}$$

- Section D.1
- Section D.2
- Section D.3
- Section D.4

- Activity D.3.4 (\sim 5 min)
- Compute $\mathcal{L}^{-1}\left\{\frac{2e^{-2s}}{s^2+4}\right\}$.
- (a) $u(t)\sin(2t)$
- (b) $u(t-2)\sin(2t)$
- (c) $u(t-2)\sin(2t-2)$
- (d) $u(t-2)\sin(2t-4)$

Section D.2 Section D.3

Section D.4

Activity D.3.5 (\sim 5 min)

Compute
$$\mathcal{L}^{-1}\left\{\frac{e^{-100s}}{s^2}\right\}$$
.

- (a) u(t)t
- (b) u(t)(t-100)
- (c) u(t-100)t
- (d) u(t-100)(t-100)

Section

Activity D.3.6 (\sim 15 min) Solve the IVP

$$500v' = 20(u(t) - u(t - 100)),$$
 $v(0) = 50.$

Activity D.3.7 (\sim 15 min) Solve the IVP

$$y'' + 4y = \delta(t - 2),$$
 $y(0) = 0, y'(0) = 1$

Module

Section D.1

Section D.1

Section [

Section D.4

Section D.5

Module D Section 4

/lodule

Section D.1

Section D.1 Section D.2

Section D.3

Section D.4

Section D. Section D. **Activity D.4.1** (\sim 30 min)

A spacecraft weighing 500 $\rm kg$ is travelling 50 $\rm m/s.$ At time $\it t=0$, its thrusters (which provide 20 $\rm N$ of force) are turned on and burn for 100 $\rm s.$

Module D Section D.1 Section D.2 Section D.3

Section D.4

Section D Section D Activity D.4.1 (\sim 30 min)

A spacecraft weighing 500 $\rm kg$ is travelling 50 $\rm m/s$. At time t=0, its thrusters (which provide 20 $\rm N$ of force) are turned on and burn for 100 $\rm s$.

Part 1: Write down a function modelling the thrust force on the spacecraft.

Module D Section D.1 Section D.2 Section D.3 Section D.4

Activity D.4.1 (\sim 30 min)

A spacecraft weighing 500 ${\rm kg}$ is travelling 50 ${\rm m/s}$. At time t=0, its thrusters (which provide 20 N of force) are turned on and burn for 100 ${\rm s}$.

Part 1: Write down a function modelling the thrust force on the spacecraft.

Part 2: Write down an IVP modelling the velocity of the spacecraft.

Activity D.4.1 (\sim 30 min)

A spacecraft weighing 500 kg is travelling 50 m/s. At time t = 0, its thrusters (which provide 20 N of force) are turned on and burn for 100 s.

Part 1: Write down a function modelling the thrust force on the spacecraft.

Part 2: Write down an IVP modelling the velocity of the spacecraft.

Part 3: Solve the IVP:

$$500v' = 20(u(t) - u(t - 100)),$$
 $v(0) = 50.$

Activity D.4.1 (\sim 30 min)

A spacecraft weighing 500 ${\rm kg}$ is travelling 50 ${\rm m/s}$. At time t=0, its thrusters (which provide 20 ${\rm N}$ of force) are turned on and burn for 100 ${\rm s}$.

Part 1: Write down a function modelling the thrust force on the spacecraft.

Part 2: Write down an IVP modelling the velocity of the spacecraft.

Part 3: Solve the IVP:

$$500v' = 20(u(t) - u(t - 100)),$$
 $v(0) = 50.$

Part 4: What is its velocity after 200 s?

		D

Section D.2

Section D.5

Module D Section 5

Module D

Section D.1

Section D.1

Section D.

Section D.

ection D.

Section D.6

Module D Section 6