

Module D

Section D.1
Section D.2
Section D.3
Section D.4
Section D.5
Section D.6

Module D: Discontinuous functions in ODEs

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

How can we solve and apply ODEs involving functions that are not continuous?

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

At the end of this module, students will be able to...

D1. Laplace Transform. ...compute the Laplace transform of a function

D2. Discontinuous ODEs. ...solve initial value problems for ODEs with discontinuous coefficients

D3. Modeling non-smooth motion. ...model the motion of an object undergoing discontinuous acceleration

D4. Modeling non-smooth oscillators. ...model mechanical oscillators undergoing discontinuous acceleration

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Readiness Assurance Outcomes

Before beginning this module, each student should be able to...

- Partial fractions

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

The following resources will help you prepare for this module.

- TODO

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Module D Section 1

Module D

Section D.1

Section D.2

Section D.3


Section D.4

Section D.5

Section D.6

Activity D.1.1 (~ 10 min)

A 1 kg mass is hung from a spring with spring constant $k = 1$ N/m. The mass is at rest, when it is hit with a hammer imparting 3J of energy.



Module D

Section D.1

Section D.2

Section D.3


Section D.4

Section D.5

Section D.6

Activity D.1.1 (~ 10 min)

A 1 kg mass is hung from a spring with spring constant $k = 1$ N/m. The mass is at rest, when it is hit with a hammer imparting 3J of energy.



Part 1: Draw a graph of the kinetic energy in the system with respect to time.

Module D

Section D.1

Section D.2

Section D.3


Section D.4

Section D.5

Section D.6

Activity D.1.1 (~ 10 min)

A 1 kg mass is hung from a spring with spring constant $k = 1$ N/m. The mass is at rest, when it is hit with a hammer imparting 3J of energy.



Part 1: Draw a graph of the kinetic energy in the system with respect to time.

Part 2: Write an initial value problem modelling this system.

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Definition D.1.2

The **Dirac delta distribution** $\delta(t)$ models the application of instantaneous force. **It is not a function**, but makes sense in definite integrals:

If a, b is any open interval containing 0, then

$$\int_a^b f(t)\delta(t)dt = f(0)$$

for any function $f(t)$ that is continuous around 0.

Definition D.1.3

The **unit impulse function** $u(t)$ is given by

$$u(t) = \begin{cases} 1 & t > 0 \\ 0 & t < 0. \end{cases}$$

Note that $u(s) = \int_{-\infty}^s \delta(t) dt$; in this fuzzy sense, δ is the derivative of $u(t)$ (which is not differentiable everywhere!)

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.1.4 (*~10 min*)

Try to solve the IVP

$$y'' + y = \delta(t)$$

Where does our existing technique break down?

Observation D.1.5

To get around this difficulty, we will apply an **integral transform** called the **Laplace Transform** to our ODE.

- We want to use a definite integral to handle things like δ , which we can only understand via a definite integral.
- Since we are focused on IVPs, we can integrate starting at 0, but need to go to ∞
- But now we need to worry about convergence—thus we will multiply by a suitable function that decays fast enough to make most functions converge.

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.1.6 (~ 5 min)

Arrange the following functions in order of how fast they decay to zero in the limit at infinity:

- (A) x^{-n} for a positive integer n
- (B) e^{-ax} for a positive integer a
- (C) $\frac{1}{\ln(ax)}$ for a positive integer a
- (D) $\frac{1}{\ln(x^n)}$ for a positive integer n

Definition D.1.7

The **Laplace Transform** of a function $f(t)$ is the function

$$F(s) = \int_0^{\infty} e^{-st} f(t) dt.$$

We will also use the notation $\mathcal{L}(f) = F$.

Note that the Laplace transform turns a function of t into a function of s .

Moreover, \mathcal{L} is linear: $\mathcal{L}(f + g) = \mathcal{L}(f) + \mathcal{L}(g)$, and $\mathcal{L}(cf) = c\mathcal{L}(f)$ for constants c .

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.1.8 (*~5 min*)

Recall that

$$\mathcal{L}(f(t)) = \int_0^{\infty} e^{-st} f(t) \, dt.$$

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.1.8 (*~5 min*)

Recall that

$$\mathcal{L}(f(t)) = \int_0^{\infty} e^{-st} f(t) dt.$$

Part 1: Compute $\mathcal{L}(\delta(t))$

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.1.8 (~ 5 min)

Recall that

$$\mathcal{L}(f(t)) = \int_0^{\infty} e^{-st} f(t) dt.$$

Part 1: Compute $\mathcal{L}(\delta(t))$

Part 2: If $a > 0$, compute $\mathcal{L}(\delta(t - a))$

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.1.9 (*~5 min*)

Recall that

$$\mathcal{L}(f(t)) = \int_0^{\infty} e^{-st} f(t) \, dt.$$

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.1.9 (*~5 min*)

Recall that

$$\mathcal{L}(f(t)) = \int_0^{\infty} e^{-st} f(t) dt.$$

Part 1: Compute $\mathcal{L}(e^t)$

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.1.9 (*~5 min*)

Recall that

$$\mathcal{L}(f(t)) = \int_0^{\infty} e^{-st} f(t) dt.$$

Part 1: Compute $\mathcal{L}(e^t)$

Part 2: If $a > 0$, compute $\mathcal{L}(e^{at})$

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.1.10 (*~15 min*)

Recall that

$$\mathcal{L}(f(t)) = \int_0^{\infty} e^{-st} f(t) dt.$$

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.1.10 (*~15 min*)

Recall that

$$\mathcal{L}(f(t)) = \int_0^{\infty} e^{-st} f(t) dt.$$

Part 1: Compute $\mathcal{L}(1)$

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.1.10 (*~15 min*)

Recall that

$$\mathcal{L}(f(t)) = \int_0^{\infty} e^{-st} f(t) dt.$$

Part 1: Compute $\mathcal{L}(1)$ *Part 2:* Compute $\mathcal{L}(t)$

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.1.10 (*~15 min*)

Recall that

$$\mathcal{L}(f(t)) = \int_0^{\infty} e^{-st} f(t) dt.$$

Part 1: Compute $\mathcal{L}(1)$ *Part 2:* Compute $\mathcal{L}(t)$ *Part 3:* Compute $\mathcal{L}(t^2)$

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.1.10 (*~15 min*)

Recall that

$$\mathcal{L}(f(t)) = \int_0^{\infty} e^{-st} f(t) dt.$$

Part 1: Compute $\mathcal{L}(1)$ *Part 2:* Compute $\mathcal{L}(t)$ *Part 3:* Compute $\mathcal{L}(t^2)$ *Part 4:* Compute $\mathcal{L}(t^3)$

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.1.10 (*~15 min*)

Recall that

$$\mathcal{L}(f(t)) = \int_0^{\infty} e^{-st} f(t) dt.$$

Part 1: Compute $\mathcal{L}(1)$ *Part 2:* Compute $\mathcal{L}(t)$ *Part 3:* Compute $\mathcal{L}(t^2)$ *Part 4:* Compute $\mathcal{L}(t^3)$ *Part 5:* Compute $\mathcal{L}(t^4)$

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Module D Section 2

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Observation D.2.1

Last week, we encountered the **Laplace Transform** of a function $f(t)$:

$$F(s) = \int_0^{\infty} e^{-st} f(t) dt.$$

We also use the notation $\mathcal{L}(f) = F$.

Recall that the Laplace transform turns a function of t into a function of s .

Moreover, \mathcal{L} is linear: $\mathcal{L}(f + g) = \mathcal{L}(f) + \mathcal{L}(g)$, and $\mathcal{L}(cf) = c\mathcal{L}(f)$ for constants c .

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Observation D.2.2

We computed a few Laplace Transforms:

- $\mathcal{L}(\delta(t - a)) = e^{-as}$ for any $a > 0$.
- $\mathcal{L}(e^{at}) = \frac{1}{s-a}$
- $\mathcal{L}(t^n) = \frac{n!}{s^{n+1}}$ for any positive integer n .
- $\mathcal{L}(1) = \frac{1}{s}$

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.2.3 (*~10 min*)

Recall

$$\mathcal{L}(f) = \int_0^{\infty} e^{-st} f(t) dt.$$

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.2.3 (*~10 min*)

Recall

$$\mathcal{L}(f) = \int_0^{\infty} e^{-st} f(t) dt.$$

Part 1: Compute $\mathcal{L}(\sin(t))$.

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.2.3 (*~10 min*)

Recall

$$\mathcal{L}(f) = \int_0^{\infty} e^{-st} f(t) dt.$$

Part 1: Compute $\mathcal{L}(\sin(t))$.*Part 2:* Compute $\mathcal{L}(\cos(t))$.

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Observation D.2.4

So now our list of Laplace transforms is:

- $\mathcal{L}(\delta(t - a)) = e^{-as}$ for any $a > 0$.
- $\mathcal{L}(e^{at}) = \frac{1}{s-a}$
- $\mathcal{L}(t^n) = \frac{n!}{s^{n+1}}$ for any positive integer n .
- $\mathcal{L}(1) = \frac{1}{s}$
- $\mathcal{L}(\sin(t)) = \frac{1}{s^2+1}$
- $\mathcal{L}(\cos(t)) = \frac{s}{s^2+1}$

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.2.5 (*~10 min*)

Suppose we want to apply the Laplace transform to an IVP: we will need to know how $\mathcal{L}(y')$ is related to $\mathcal{L}(y)$. Recall

$$\mathcal{L}(f) = \int_0^{\infty} e^{-st} f(t) dt.$$

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.2.5 (*~10 min*)

Suppose we want to apply the Laplace transform to an IVP: we will need to know how $\mathcal{L}(y')$ is related to $\mathcal{L}(y)$. Recall

$$\mathcal{L}(f) = \int_0^{\infty} e^{-st} f(t) dt.$$

Part 1: Use integration by parts to relate $\mathcal{L}(y')$ to $\mathcal{L}(y)$.

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.2.5 (~ 10 min)

Suppose we want to apply the Laplace transform to an IVP: we will need to know how $\mathcal{L}(y')$ is related to $\mathcal{L}(y)$. Recall

$$\mathcal{L}(f) = \int_0^{\infty} e^{-st} f(t) dt.$$

Part 1: Use integration by parts to relate $\mathcal{L}(y')$ to $\mathcal{L}(y)$.

Part 2: Use integration by parts (and the fact that $\mathcal{L}(y') = s\mathcal{L}(y) - y(0)$) to relate $\mathcal{L}(y'')$ to $\mathcal{L}(y)$.

Observation D.2.6

We have

$$\mathcal{L}(y') = sL(y) - y(0)$$

$$\mathcal{L}(y'') = s^2L(y) - sy(0) - y'(0)$$

This allows us to easily rewrite expressions like $ay'' + by' + cy$ in terms of $\mathcal{L}(y)$.

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.2.7 (*~10 min*)

Consider the simple IVP

$$y'' + y = \delta(t), \quad y(0) = 0, y'(0) = 0.$$

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.2.7 (*~10 min*)

Consider the simple IVP

$$y'' + y = \delta(t), \quad y(0) = 0, y'(0) = 0.$$

Part 1: Apply the Laplace transform to this IVP, and simplify. Solve for $\mathcal{L}(y)$.

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.2.7 (*~10 min*)

Consider the simple IVP

$$y'' + y = \delta(t), \quad y(0) = 0, y'(0) = 0.$$

Part 1: Apply the Laplace transform to this IVP, and simplify. Solve for $\mathcal{L}(y)$.

Part 2: Find a function y satisfying $\mathcal{L}(y) = \frac{1}{s^2+1}$. We write $y = \mathcal{L}^{-1}\left(\frac{1}{s^2+1}\right)$.

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.2.8 (*~15 min*)

Solve the IVP

$$y'' + y = \delta(t), \quad y(0) = 1, y'(0) = 2.$$

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Module D Section 3

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Observation D.3.1

To solve an IVP using Laplace transforms:

- 1) Apply \mathcal{L} to the ODE. Use the initial condition(s) in computing $\mathcal{L}(y')$, $\mathcal{L}(y'')$, etc.
- 2) Solve for $\mathcal{L}(y)$.
- 3) Take the inverse transform (using a table) to find the solution y .

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.3.2 (~ 5 min)Compute $\mathcal{L}^{-1}\left(\frac{e^{-10s}}{s}\right)$.

(a) $u(t - 10)$

(b) $\delta(t - 10)$

(c) $u(t - 10)e^{-t}$

(d) $\delta(t - 10)e^{-t}$

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.3.3 (~ 5 min)Compute $\mathcal{L}^{-1}\left(\frac{1}{s+5}\right)$.

(a) $u(t-5)$

(b) $\delta(t-5)$

(c) e^{5t}

(d) e^{-5t}

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.3.4 (~ 5 min)Compute $\mathcal{L}^{-1}\left(\frac{2e^{-2s}}{s^2+4}\right)$.

- (a) $u(t) \sin(2t)$
- (b) $u(t-2) \sin(2t)$
- (c) $u(t-2) \sin(2t-2)$
- (d) $u(t-2) \sin(2t-4)$

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.3.5 (~ 5 min)Compute $\mathcal{L}^{-1}\left(\frac{e^{-100s}}{s^2}\right)$.

(a) $u(t)t$

(b) $u(t)(t - 100)$

(c) $u(t - 100)t$

(d) $u(t - 100)(t - 100)$

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.3.6 (*~15 min*)

Solve the IVP

$$500v' = 20(u(t) - u(t - 100)), \quad v(0) = 50.$$

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.3.7 (*~15 min*)

Solve the IVP

$$y'' + 4y = \delta(t - 2),$$

$$y(0) = 0, \quad y'(0) = 1$$

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Module D Section 4

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.4.1 (~ 30 min)

A spacecraft weighing 500 kg is travelling 50 m/s. At time $t = 0$, its thrusters (which provide 20 N of force) are turned on and burn for 100 s.

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.4.1 (~ 30 min)

A spacecraft weighing 500 kg is travelling 50 m/s. At time $t = 0$, its thrusters (which provide 20 N of force) are turned on and burn for 100 s.

Part 1: Write down a function modelling the thrust force on the spacecraft.

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.4.1 (~ 30 min)

A spacecraft weighing 500 kg is travelling 50 m/s. At time $t = 0$, its thrusters (which provide 20 N of force) are turned on and burn for 100 s.

Part 1: Write down a function modelling the thrust force on the spacecraft.

Part 2: Write down an IVP modelling the velocity of the spacecraft.

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.4.1 (~ 30 min)

A spacecraft weighing 500 kg is travelling 50 m/s. At time $t = 0$, its thrusters (which provide 20 N of force) are turned on and burn for 100 s.

Part 1: Write down a function modelling the thrust force on the spacecraft.

Part 2: Write down an IVP modelling the velocity of the spacecraft.

Part 3: Solve the IVP:

$$500v' = 20(u(t) - u(t - 100)), \quad v(0) = 50.$$

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Activity D.4.1 (~ 30 min)

A spacecraft weighing 500 kg is travelling 50 m/s. At time $t = 0$, its thrusters (which provide 20 N of force) are turned on and burn for 100 s.

Part 1: Write down a function modelling the thrust force on the spacecraft.

Part 2: Write down an IVP modelling the velocity of the spacecraft.

Part 3: Solve the IVP:

$$500v' = 20(u(t) - u(t - 100)), \quad v(0) = 50.$$

Part 4: What is its velocity after 200 s?

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Module D Section 5

Module D

Section D.1

Section D.2

Section D.3

Section D.4

Section D.5

Section D.6

Module D Section 6