

Module C

Standard C1

C1. Find the general solution to

$$y' + 3y = 6t + 5.$$

C1. Find the general solution to

$$y' + 4y = 4.$$

C1. Find the general solution to

$$y' + 2y = 6t - 1.$$

C1. Find the general solution to

$$y' - y = e^t.$$

C1. Find the general solution to

$$y' + y = e^t.$$

C1. Find the general solution to

$$y' - y = e^{-t}.$$

C1. Find the general solution to

$$y' + y = e^{-t}.$$

C1. Find the general solution to

$$y' + 3y = 10e^{-3t} \sin(t).$$

C1. Find the general solution to

$$y' + 2y = 10e^{-2t} \sin(t).$$

C1. Find the general solution to

$$y' + 2y = 5e^{-2t} \sin(t).$$

C1. Find the general solution to

$$y' + 3y = 10e^{-3t} \cos(t).$$

C1. Find the general solution to

$$y' + 2y = 10e^{-2t} \cos(t).$$

C1. Find the general solution to

$$y' + 2y = 5e^{-2t} \cos(t).$$

C2. A water droplet with a radius of $100\ \mu\text{m}$ has a mass of about $4 \times 10^{-12}\text{kg}$ and a terminal velocity of $27\ \frac{\text{cm}}{\text{s}}$. Such a droplet is dropped from rest.

- (a) Write down an Initial Value Problem (IVP) modelling the velocity.
- (b) What is its velocity after $0.01\ \text{s}$?

C2. A water droplet with a radius of $50\ \mu\text{m}$ has a mass of about $5 \times 10^{-13}\text{kg}$ and a terminal velocity of $3.5\ \frac{\text{cm}}{\text{s}}$. Such a droplet is dropped from rest.

- (a) Write down an Initial Value Problem (IVP) modelling the velocity.
- (b) What is its velocity after $0.01\ \text{s}$?

C2. A water droplet with a radius of $10\ \mu\text{m}$ has a mass of about $4 \times 10^{-15}\text{kg}$ and a terminal velocity of $270\ \frac{\mu\text{m}}{\text{s}}$. Such a droplet is dropped from rest.

- (a) Write down an Initial Value Problem (IVP) modelling the velocity.
- (b) What is its velocity after $0.001\ \text{s}$?

C2. A water droplet with a radius of $5\ \mu\text{m}$ has a mass of about $5 \times 10^{-16}\text{kg}$ and a terminal velocity of $35\ \frac{\mu\text{m}}{\text{s}}$. Such a droplet is dropped from rest.

- (a) Write down an Initial Value Problem (IVP) modelling the velocity.
- (b) What is its velocity after $0.001\ \text{s}$?

C2. A single grain of corn pollen with a radius of $50\ \mu\text{m}$ and a mass of about $5 \times 10^{-13}\text{kg}$ has a terminal velocity of $27\ \frac{\text{cm}}{\text{s}}$. Such a pollen grain is dropped from rest.

- (a) Write down an Initial Value Problem (IVP) modelling the velocity.
- (b) What is its velocity after $0.01\ \text{s}$?

C2. A single grain of spruce pollen with a radius of $25\ \mu\text{m}$ and a mass of about $6 \times 10^{-14}\text{kg}$ has a terminal velocity of $3\ \frac{\text{cm}}{\text{s}}$. Such a pollen grain is dropped from rest.

- (a) Write down an Initial Value Problem (IVP) modelling the velocity.
- (b) What is its velocity after $0.01\ \text{s}$?

C3. Find the general solution to

$$y'' + 2y' + y = 0.$$

C3. Find the general solution to

$$y'' + 2y' - 8y = 0.$$

C3. Find the general solution to

$$y'' + 4y' + 3y = 0.$$

C3. Find the general solution to

$$y'' + 2y' - 3y = 0.$$

C3. Find the general solution to

$$y'' - 2y' - 3y = 0.$$

C3. Find the general solution to

$$y'' + 4y' + 4y = 0.$$

C3. Find the general solution to

$$y'' - 4y' + 4y = 0.$$

C3. Find the general solution to

$$y'' + 5y' + 6y = 0.$$

C3. Find the general solution to

$$y'' - 2y' + 2y = 0.$$

C3. Find the general solution to

$$y'' + 2y' + 2y = 0.$$

C3. Find the general solution to

$$y'' - 6y' + 10y = 0.$$

C3. Find the general solution to

$$y'' + 6y' + 10y = 0.$$

C3. Find the general solution to

$$y'' - 2y' + 5y = 0.$$

C3. Find the general solution to

$$y'' + 2y' + 5y = 0.$$

C3. Find the general solution to

$$y'' - 4y' + 5y = 0.$$

C3. Find the general solution to

$$y'' + 4y' + 5y = 0.$$

C4. Find the solution to

$$y'' + 2y' + y = 0$$

when $y(0) = 0$ and $y'(0) = 2$.

C4. Find the solution to

$$y'' + 2y' + y = 0$$

when $y(0) = 2$ and $y'(0) = 0$.

C4. Find the solution to

$$y'' + 2y' - 8y = 0$$

when $y(0) = 3$ and $y'(0) = -6$.

C4. Find the solution to

$$y'' + 4y' + 3y = 0$$

when $y(0) = 1$ and $y'(0) = 5$.

C4. Find the solution to

$$y'' + 2y' - 3y = 0$$

when $y(0) = 5$ and $y'(0) = 1$.

C4. Find the solution to

$$y'' + 2y' - 3y = 0$$

when $y(0) = 2$ and $y'(0) = 2$.

C4. Find the solution to

$$y'' - 2y' - 3y = 0$$

when $y(0) = 2$ and $y'(0) = 2$.

C4. Find the solution to

$$y'' + 4y' + 4y = 0$$

when $y(0) = 1$ and $y'(0) = 3$.

C4. Find the solution to

$$y'' - 4y' + 4y = 0$$

when $y(0) = 1$ and $y'(0) = 3$.

C4. Find the solution to

$$y'' + 4y' + 4y = 0$$

when $y(0) = 3$ and $y'(0) = 1$.

C4. Find the solution to

$$y'' - 4y' + 4y = 0$$

when $y(0) = 3$ and $y'(0) = 1$.

C4. Find the solution to

$$y'' + 5y' + 6y = 0$$

when $y(0) = 3$ and $y'(0) = 1$.

C4. Find the solution to

$$y'' + 5y' + 6y = 0$$

when $y(0) = 1$ and $y'(0) = 2$.

C5. Find a general solution to the given equation.

$$y'' + 2y' + y = 3x + 4$$

C5. Find a general solution to the given equation.

$$y'' + 4y' + 3y = 2 \sin(3x)$$

C5. Find a general solution to the given equation.

$$y'' - 2y' - 3y = 1 + xe^x$$

C5. Find a general solution to the given equation.

$$y'' - 4y' + 4y = e^{2x}$$

C5. Find a general solution to the given equation.

$$y'' + 4y' + 4y = e^{2x}$$

C5. Find a general solution to the given equation.

$$y'' + 4y = \cos(2x)$$

C5. Find a general solution to the given equation.

$$y'' - 4y = \cos(2x)$$

C5. Find a general solution to the given equation.

$$y'' + 9y = \sin(3x)$$

C5. Find a general solution to the given equation.

$$y'' - 9y = \sin(3x)$$

C5. Find a general solution to the given equation.

$$y'' - 2y' + 2y = \sin(x)$$

C5. Find a general solution to the given equation.

$$y'' - 2y' + 5y = 2x + 1$$

C6. Consider the following scenario: A 1kg mass is suspended by a spring (with spring constant 4kg/s^2). The mass is pulled down 1m from its equilibrium position and released from rest.

- (a) Write down an IVP modelling the position of the mass.
- (b) How long does it take for the mass to return to its equilibrium point?

C6. Consider the following scenario: A 1kg mass is suspended by a spring (with spring constant 4kg/s^2). The mass is pushed up 0.5m from its equilibrium position and released from rest.

- (a) Write down an IVP modelling the position of the mass.
- (b) How long does it take for the mass to return to its equilibrium point?

C6. Consider the following scenario: A 4kg mass is suspended by a spring (with spring constant 1kg/s^2). The mass is pulled down 1m from its equilibrium position and released from rest.

- (a) Write down an IVP modelling the position of the mass.
- (b) Where is the mass after 3s?

C6. Consider the following scenario: A 4kg mass is suspended by a spring (with spring constant 1kg/s^2). The mass is pushed up 0.5m from its equilibrium position and released from rest.

- (a) Write down an IVP modelling the position of the mass.
- (b) Where is the mass after 2s?

C6. Consider the following scenario: A 2kg mass is suspended by a spring (with spring constant 4kg/s^2). A linear damper is attached to the system (with constant 6kg/s). The mass is pulled down 1m from its equilibrium position and released from rest.

- (a) Write down an IVP modelling the position of the mass.
- (b) Where is the mass after 2s?

C6. Consider the following scenario: A 2kg mass is suspended by a spring (with spring constant 4kg/s^2). A linear damper is attached to the system (with constant 1kg/s). The mass is pulled down 1m from its equilibrium position and released from rest.

- (a) Write down an IVP modelling the position of the mass.
- (b) Where is the mass after 2s?

C6. Consider the following scenario: A 2kg mass is suspended by a spring (with spring constant 4kg/s^2). A linear damper is attached to the system (with constant 6kg/s). An external force is applied, modelled by the function $F(t) = \sin(t)$. The mass is pulled down 1m from its equilibrium position and released from rest.

- (a) Write down an IVP modelling the position of the mass.
- (b) Where is the mass after 2s?

C6. Consider the following scenario: A 2kg mass is suspended by a spring (with spring constant 4kg/s^2). A linear damper is attached to the system (with constant 6kg/s). An external force is applied, modelled by the function $F(t) = \cos(t)$. The mass is pulled down 1m from its equilibrium position and released from rest.

- (a) Write down an IVP modelling the position of the mass.
- (b) Where is the mass after 2s?

Module F

Standard F1


F1. Sketch a solution curve through each point marked in the slope field.




F1. Sketch a solution curve through each point marked in the slope field.



F1. Sketch a solution curve through each point marked in the slope field.




F1. Sketch a solution curve through each point marked in the slope field.



F1. Sketch a solution curve through each point marked in the slope field.



F1. Sketch a solution curve through each point marked in the slope field.



F1. Sketch a solution curve through each point marked in the slope field.




F1. Sketch a solution curve through each point marked in the slope field.



F1. Sketch a solution curve through each point marked in the slope field.

$$y' = x^2 - y$$



F2. Find the general solution to $\frac{dy}{dx} + 3xy = 0$.

F2. Find the general solution to $y' - y \sin(x) = 0$.

F2. Find the general solution to $y' - y^2 e^x = 0$.

F2. Find the general solution to $y' = \frac{x+2}{y^2}$.

F2. Find the general solution to $xy' = y$.

F2. Find the general solution to $y \frac{dy}{dx} = y^2 \cos(x)$.

F2. Find the general solution to $xy^2 \frac{dy}{dx} = 1$.

F2. Find the general solution to $x \cos(y) y' = 1$.

F3. A tennis ball has a mass of 0.055kg and a drag coefficient of 0.001kg/m. It leaves the server's racket travelling 60m/s.

- (a) Write down an IVP modelling the horizontal velocity of the ball (ignore any vertical movement in the ball).
- (b) How long does it take the ball to travel across a tennis court, which is 24m long?

F3. A tennis ball has a mass of 0.055kg and a drag coefficient of 0.001kg/m. It leaves the server's racket travelling 60m/s.

- (a) Write down an IVP modelling the horizontal velocity of the ball (ignore any vertical movement in the ball).
- (b) How far has the ball gone after 0.5s?

F3. A tennis ball has a mass of 0.055kg and a drag coefficient of 0.001kg/m. It leaves the server's racket travelling 50m/s.

- (a) Write down an IVP modelling the horizontal velocity of the ball (ignore any vertical movement in the ball).
- (b) How long does it take the ball to travel across a tennis court, which is 24m long?

F3. A tennis ball has a mass of 0.055kg and a drag coefficient of 0.001kg/m. It leaves the server's racket travelling 50m/s.

- (a) Write down an IVP modelling the horizontal velocity of the ball (ignore any vertical movement in the ball).
- (b) How far has the ball gone after 0.5s?

F3. A baseball has a mass of 0.145kg and a drag coefficient of 0.0009kg/m. A batter hits a line drive; the ball leaves the bat travelling 50m/s.

- (a) Write down an IVP modelling the horizontal velocity of the ball (ignore any vertical movement in the ball).
- (b) How long does it take for the ball to reach a defender standing 30m away?

F3. A baseball has a mass of 0.145kg and a drag coefficient of 0.0009kg/m. A batter hits a line drive; the ball leaves the bat travelling 50m/s.

- (a) Write down an IVP modelling the horizontal velocity of the ball (ignore any vertical movement in the ball).
- (b) How far has the ball gone after 1s?

F3. A baseball has a mass of 0.145kg and a drag coefficient of 0.0009kg/m. A batter hits a line drive; the ball leaves the bat travelling 45m/s.

- (a) Write down an IVP modelling the horizontal velocity of the ball (ignore any vertical movement in the ball).

(b) How long does it take for the ball to reach a defender standing 30m away?

F3. A baseball has a mass of 0.145kg and a drag coefficient of 0.0009kg/m. A batter hits a line drive; the ball leaves the bat travelling 45m/s.

(a) Write down an IVP modelling the horizontal velocity of the ball (ignore any vertical movement in the ball).

(b) How far has the ball gone after 1s?

F4. Consider the autonomous equation

$$\frac{dx}{dt} = x - 3.$$

- (a) Find and classify the critical points.
- (b) Describe the long term behavior of the solution passing through the point $x(0) = 4$.

F4. Consider the autonomous equation

$$\frac{dx}{dt} = 1 - x.$$

- (a) Find and classify the critical points.
- (b) Describe the long term behavior of the solution passing through the point $x(2) = 2$.

F4. Consider the autonomous equation

$$\frac{dx}{dt} = (x - 3)^2.$$

- (a) Find and classify the critical points.
- (b) Describe the long term behavior of the solution passing through the point $x(1) = 2$.

F4. Consider the autonomous equation

$$\frac{dx}{dt} = (x + 4)^2.$$

- (a) Find and classify the critical points.
- (b) Describe the long term behavior of the solution passing through the point $x(4) = 0$.

F4. Consider the autonomous equation

$$\frac{dx}{dt} = (4 - x)^3.$$

- (a) Find and classify the critical points.
- (b) Describe the long term behavior of the solution passing through the point $x(3) = 2$.

F4. Consider the autonomous equation

$$\frac{dx}{dt} = (5 - x)^3.$$

- (a) Find and classify the critical points.
- (b) Describe the long term behavior of the solution passing through the point $x(0) = 4$.

F4. Consider the autonomous equation

$$\frac{dx}{dt} = x^2 - 7x + 10.$$

- (a) Find and classify the critical points.
- (b) Describe the long term behavior of the solution passing through the point $x(0) = 3$.

F4. Consider the autonomous equation

$$\frac{dx}{dt} = x^2 - x - 6.$$

- (a) Find and classify the critical points.
- (b) Describe the long term behavior of the solution passing through the point $x(3) = 0$.

F4. Consider the autonomous equation

$$\frac{dx}{dt} = x^2(x^2 - x - 6).$$

- (a) Find and classify the critical points.
- (b) Describe the long term behavior of the solution passing through the point $x(5) = 1$.

F4. Consider the autonomous equation

$$\frac{dx}{dt} = x^2 - 4x + 3.$$

- (a) Find and classify the critical points.
- (b) Describe the long term behavior of the solution passing through the point $x(2) = 2$.

F4. Consider the autonomous equation

$$\frac{dx}{dt} = x(x^2 - 4x + 3).$$

- (a) Find and classify the critical points.
- (b) Describe the long term behavior of the solution passing through the point $x(2) = 2$.

F4. Consider the autonomous equation

$$\frac{dx}{dt} = x(x^2 - 9x + 20).$$

- (a) Find and classify the critical points.
- (b) Describe the long term behavior of the solution passing through the point $x(2) = 2$.

- F5.** Find the general solution to $xy' + 4y = 2x$.
- F5.** Find the general solution to $xy' + 2y = x^2$.
- F5.** Find the general solution to $xy' + 2y = 4x^2 - 3x$.
- F5.** Find the general solution to $xy' + 2y = x^2 - 3x$.
- F5.** Find the general solution to $\cos(x)y' + \sin(x)y = x + \sin(x)\cos(x)$.
- F5.** Find the general solution to $\cos(x)y' + \sin(x)y = x\cos^2(x)$.
- F5.** Find the general solution to $(x^2 + 1)y' - 2xy = 1$.
- F5.** Find the general solution to $(x^2 + 1)y' - 2xy = x + 1$.

F6. One of the two ODEs below is exact. Identify which one, and solve it.

$$\begin{aligned}(x + 2y)y' + y &= 2x \\ (x + 2y)y' - y &= -2x\end{aligned}$$

F6. One of the two ODEs below is exact. Identify which one, and solve it.

$$\begin{aligned}(3x + 2y)y' + 3y &= 2x \\ (3x + 2y)y' - 3y &= -2x\end{aligned}$$

F6. One of the two ODEs below is exact. Identify which one, and solve it.

$$\begin{aligned}(x^2 + 3y^2)y' - 2xy &= -3x^2 \\ (x^2 + 3y^2)y' + 2xy &= 3x^2\end{aligned}$$

F6. One of the two ODEs below is exact. Identify which one, and solve it.

$$\begin{aligned}(2xy + 3y^2)y' + y^2 &= 3x^2 \\ (2xy + 3y^2)y' - y^2 &= -3x^2\end{aligned}$$

F6. One of the two ODEs below is exact. Identify which one, and solve it.

$$\begin{aligned}\cos(x) \cos(y)y' &= \sin(x) \sin(y) \\ \cos(x) \cos(y)y' &= \sin(x) + \sin(y)\end{aligned}$$

F6. One of the two ODEs below is exact. Identify which one, and solve it.

$$\begin{aligned}\sin(x) \sin(y)y' &= \cos(x) + \cos(y) \\ \sin(x) \sin(y)y' &= \cos(x) \cos(y)\end{aligned}$$

F6. One of the two ODEs below is exact. Identify which one, and solve it.

$$\begin{aligned}(y^3 e^x + x e^x)y' + 3e^x y^2 &= 3x^2 \\ (2y e^x + e^y)y' + e^x y^2 &= 3x^2\end{aligned}$$

Module S

Standard S1

S1. Find the general solution of the system

$$\begin{aligned}x' &= x + y, \\y' &= 4x + y.\end{aligned}$$

S1. Find the general solution of the system

$$\begin{aligned}x' &= x + 2y, \\y' &= 3x + 2y.\end{aligned}$$

S1. Find the general solution of the system

$$\begin{aligned}x' &= 2x + y, \\y' &= x + 2y.\end{aligned}$$

S1. Find the general solution of the system

$$\begin{aligned}x' &= 2x + y, \\y' &= 2x + 3y.\end{aligned}$$

S1. Find the general solution of the system

$$\begin{aligned}x' &= 3x + y, \\y' &= x + 3y.\end{aligned}$$

S1. Find the general solution of the system

$$\begin{aligned}x' &= 3x + y, \\y' &= 2x + 2y.\end{aligned}$$

S1. Find the general solution of the system

$$\begin{aligned}x' &= 4x + y, \\y' &= 2x + 3y.\end{aligned}$$

S1. Find the general solution of the system

$$\begin{aligned}x' &= 4x + 3y, \\y' &= x + 2y.\end{aligned}$$

S2. Two populations of competing species of fish, bluegills and greenfish, are modelled by the system

$$\begin{aligned}\frac{dG}{dt} &= 0.1G - 0.002G^2 - 0.005BG \\ \frac{dB}{dt} &= 0.1B - 0.003B^2 - 0.005BG.\end{aligned}$$

- (a) Identify all equilibrium points for the system.
- (b) If a lake is stocked with , what will happen to the two populations in the long term?

S2. Two populations of competing species of fish, bluegills and greenfish, are modelled by the system

$$\begin{aligned}\frac{dG}{dt} &= 0.1G - 0.002G^2 - 0.005BG \\ \frac{dB}{dt} &= 0.1B - 0.004B^2 - 0.005BG.\end{aligned}$$

- (a) Identify all equilibrium points for the system.
- (b) If a lake is stocked with , what will happen to the two populations in the long term?

S2. Two populations of competing species of fish, bluegills and greenfish, are modelled by the system

$$\begin{aligned}\frac{dG}{dt} &= 0.1G - 0.002G^2 - 0.004BG \\ \frac{dB}{dt} &= 0.1B - 0.003B^2 - 0.005BG.\end{aligned}$$

- (a) Identify all equilibrium points for the system.
- (b) If a lake is stocked with , what will happen to the two populations in the long term?

S2. The populations of foxes and rabbits in an ecosystem are modelled by the system

$$\begin{aligned}\frac{dF}{dt} &= -0.1F - 0.001F^2 + 0.002FR \\ \frac{dR}{dt} &= 0.2R - 0.001R^2 - 0.005FR.\end{aligned}$$

- (a) Identify all equilibrium points for the system.
- (b) If there are presently 100 foxes and 200 rabbits , what will happen to the two populations in the long term?

S2. The populations of foxes and rabbits in an ecosystem are modelled by the system

$$\begin{aligned}\frac{dF}{dt} &= -0.1F - 0.001F^2 + 0.002FR \\ \frac{dR}{dt} &= 0.4R - 0.001R^2 - 0.005FR.\end{aligned}$$

- (a) Identify all equilibrium points for the system.
- (b) If there are presently 100 foxes and 200 rabbits , what will happen to the two populations in the long term?

S2. The populations of greenflies and ladybirds in an ecosystem are modelled by the system

$$\begin{aligned}\frac{dL}{dt} &= -0.1L - 0.001L^2 + 0.002LG \\ \frac{dG}{dt} &= 0.4G - 0.001G^2 - 0.001GL.\end{aligned}$$

- (a) Identify all equilibrium points for the system.
- (b) If there are presently 1000 ladybirds and 2000 greenflies, what will happen to the two populations in the long term?

S2. The populations of greenflies and ladybirds in an ecosystem are modelled by the system

$$\begin{aligned}\frac{dL}{dt} &= -0.1L - 0.001L^2 + 0.002LG \\ \frac{dG}{dt} &= 0.2G - 0.001G^2 - 0.001GL.\end{aligned}$$

- (a) Identify all equilibrium points for the system.
- (b) If there are presently 1000 ladybirds and 2000 greenflies, what will happen to the two populations in the long term?

S3.

Module N

Standard N1

N1. Determine whether existence of at least one solution of the given initial value problem is guaranteed and, if so, whether uniqueness of that solution is guaranteed.

$$y' = x^2y + xy^2; \quad y(1) = 3$$

N1. Determine whether existence of at least one solution of the given initial value problem is guaranteed and, if so, whether uniqueness of that solution is guaranteed.

$$y' = 2x^2 + xy + 3y^2; \quad y(1) = -1$$

N1. Determine whether existence of at least one solution of the given initial value problem is guaranteed and, if so, whether uniqueness of that solution is guaranteed.

$$y' = x + \ln(y); \quad y(1) = 2$$

N1. Determine whether existence of at least one solution of the given initial value problem is guaranteed and, if so, whether uniqueness of that solution is guaranteed.

$$y' = \sqrt{x+y}; \quad y(1) = 1$$

N1. Determine whether existence of at least one solution of the given initial value problem is guaranteed and, if so, whether uniqueness of that solution is guaranteed.

$$y' = \sqrt[3]{x-y}; \quad y(2) = 2$$

N1. Determine whether existence of at least one solution of the given initial value problem is guaranteed and, if so, whether uniqueness of that solution is guaranteed.

$$y' = \frac{y}{x}; \quad y(2) = 1$$

N2. Consider the differential equation

$$xy'' + y' = 0.$$

Determine all intervals on which a unique solution is guaranteed to exist.

N2. Consider the differential equation

$$xy'' - y' = 0.$$

Determine all intervals on which a unique solution is guaranteed to exist.

N2. Consider the differential equation

$$x^2y'' - 4xy' + 4y = 0.$$

Determine all intervals on which a unique solution is guaranteed to exist.

N2. Consider the differential equation

$$x^2y'' - xy' - 3y = 0.$$

Determine all intervals on which a unique solution is guaranteed to exist.

N2. Consider the differential equation

$$x^2y'' + xy' + 4y = 0.$$

Determine all intervals on which a unique solution is guaranteed to exist.

N2. Consider the differential equation

$$y'' - \frac{1}{1+x}y' + \frac{1}{(1+x)^2}y = 0.$$

Determine all intervals on which a unique solution is guaranteed to exist.

N2. Consider the differential equation

$$y'' + \frac{2}{x-2}y' - \frac{6}{(x-2)^2}y = 0.$$

Determine all intervals on which a unique solution is guaranteed to exist.

N2. Consider the differential equation

$$e^xy'' - 2y' + 4e^{4x}y = 0.$$

Determine all intervals on which a unique solution is guaranteed to exist.

N2. Consider the differential equation

$$y'' + y' - e^{-2x}y = 0.$$

Determine all intervals on which a unique solution is guaranteed to exist.

N3. Determine all intervals on which a unique solution is guaranteed to exist.

$$\begin{aligned}x' &= -\frac{3}{t}x + 2y, \\y' &= 2\ln(t)x + y + 1\end{aligned}$$

N3. Determine all intervals on which a unique solution is guaranteed to exist.

$$\begin{aligned}x' &= -\frac{2}{t}x + y, \\y' &= x + \ln(t)y + 2\end{aligned}$$

N3. Determine all intervals on which a unique solution is guaranteed to exist.

$$\begin{aligned}x' &= -x + \sqrt{t}, \\y' &= 2x + ty + \sqrt[3]{t}\end{aligned}$$

N3. Determine all intervals on which a unique solution is guaranteed to exist.

$$\begin{aligned}x' &= x + 2y + \sqrt{t}, \\y' &= x + y + \sqrt[3]{t}\end{aligned}$$

N3. Determine all intervals on which a unique solution is guaranteed to exist.

$$\begin{aligned}x' &= x + y + \sqrt[3]{t}, \\y' &= x + 2y + \sqrt{t}\end{aligned}$$

N3. Determine all intervals on which a unique solution is guaranteed to exist.

$$\begin{aligned}x' &= tx + 2y + \sqrt[3]{t}, \\y' &= -y + \sqrt{t}\end{aligned}$$

N3. Determine all intervals on which a unique solution is guaranteed to exist.

$$\begin{aligned}x' &= x + \ln(t)y + 2, \\y' &= -\frac{1}{t}y + 2t\end{aligned}$$

N3. Determine all intervals on which a unique solution is guaranteed to exist.

$$x' = 2 \ln(t)x + y + 1,$$

$$y' = -\frac{2}{t}x + y$$

N4.

N4.

Module D

Standard D1

D1. Demonstrate directly from the definition that

$$\mathcal{L}\{u(t+1)\}(s) = \frac{e^s}{s}.$$

D1. Demonstrate directly from the definition that

$$\mathcal{L}\{u(t-5)\}(s) = \frac{e^{-5s}}{s}.$$

D1. Demonstrate directly from the definition that

$$\mathcal{L}\{\delta(t+3)\}(s) = e^{3s}.$$

D1. Demonstrate directly from the definition that

$$\mathcal{L}\{\delta(t-2)\}(s) = e^{-2s}.$$

D1. Demonstrate directly from the definition that

$$\mathcal{L}\{e^{3t}\}(s) = \frac{1}{s-3}.$$

D1. Demonstrate directly from the definition that

$$\mathcal{L}\{e^{-2t}\}(s) = \frac{1}{s+2}.$$

D1. Demonstrate directly from the definition that

$$\mathcal{L}\{\delta(t+4) + e^t\}(s) = e^{4s} + \frac{1}{s-1}.$$

D1. Demonstrate directly from the definition that

$$\mathcal{L}\{\delta(t) + u(t-5)\}(s) = 1 + \frac{e^{-5s}}{s}.$$

D1. Demonstrate directly from the definition that

$$\mathcal{L}\{1 + e^t\}(s) = \frac{1}{s} + \frac{1}{s-1}.$$

D2.

D3.

D4.