

ICPSR 36498

Population Assessment of Tobacco and Health (PATH) Study [United States] Public-Use Files

United States Department of Health and Human Services. National Institutes of Health. National Institute on Drug Abuse

United States Department of Health and Human Services. Food and Drug Administration. Center for Tobacco Products

Wave 4: Adult - Wave 4 Cohort Cross-Sectional Weights

Inter-university Consortium for
Political and Social Research
P.O. Box 1248
Ann Arbor, Michigan 48106
www.icpsr.umich.edu

Terms of Use

The terms of use for this study can be found at:
<http://www.icpsr.umich.edu/web/ICPSR/studies/36498/terms>

Information about Copyrighted Content

Some instruments administered for studies archived with ICPSR may contain in whole or substantially in part contents from copyrighted instruments. Reproductions of the instruments are provided as documentation for the analysis of the data associated with this collection. Restrictions on "fair use" apply to all copyrighted content. More information about the reproduction of copyrighted works by educators and librarians is available from the United States Copyright Office.

NOTICE

WARNING CONCERNING COPYRIGHT RESTRICTIONS

The copyright law of the United States (Title 17, United States Code) governs the making of photocopies or other reproductions of copyrighted material. Under certain conditions specified in the law, libraries and archives are authorized to furnish a photocopy or other reproduction. One of these specified conditions is that the photocopy or reproduction is not to be "used for any purpose other than private study, scholarship, or research." If a user makes a request for, or later uses, a photocopy or reproduction for purposes in excess of "fair use," that user may be liable for copyright infringement.

NOTE

This document does not currently comply with 508 standards. If you need an alternative means of access to any information, please contact NAHDAP at nahdap@icpsr.umich.edu. Let us know the nature of your accessibility problem, the Web address of the required information, and your contact information.

CODEBOOK DISPLAY NOTES

PATH STUDY

WEIGHTS FILE ICPSR

#36498

1. ICPSR customized the display of variables in this codebook. There are no variables that contain a frequency table displaying value labels and unweighted counts. This is because all of the variables in this file are identification, sample design, or sampling weight variables for which univariate statistics are not meaningful.
2. Each variable contains the following statement:

“This variable has XXXXX valid cases out of XXXXX total cases.”

This statement describes the number of valid cases for a variable that would be used in the calculation of percentages in the frequency tables.

ICPSR 36498

Population Assessment of Tobacco and Health (PATH) Study [United States] Public-Use Files

Variable Description and Frequencies

Note: Frequencies displayed for the variables are not weighted. They are purely descriptive and may not be representative of the study population. Please review any sampling or weighting information available with the study.

Summary statistics (minimum, maximum, mean, median, and standard deviation) may not be available for every variable in the codebook. Conversely, a listing of frequencies in table format may not be present for every variable in the codebook either. However, all variables in the dataset are present and display sufficient information about each variable. These decisions are made intentionally and are at the discretion of the archive producing this codebook.

Wave 4: Adult - Wave 4 Cohort Cross-Sectional Weights

CASEID: Case Identification Number

Case Identification Number

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1-5 (width: 5; decimal: 0)

Variable Type: numeric

PERSONID: Participant ID

Participant ID Number.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 6-15 (width: 10; decimal: 0)

Variable Type: character

VARPSU: PSU Indicator for Variance Estimation

PSU indicator for variance estimation using the Taylor series approximation method.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 18-18 (width: 1; decimal: 0)

Variable Type: numeric

VARSTRAT: Stratum Indicator for Variance Estimation

Stratum indicator for variance estimation using the Taylor series approximation method.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 16-17 (width: 2; decimal: 0)

Variable Type: numeric

R04_A_C04WGT: Wave 4 Adult Cross-sectional Weight for the Wave 4 Cohort

Wave 4 adult cross-sectional full-sample weight for the Wave 4 cohort.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 19-33 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT1: Wave 4 Adult Cross-sectional Replicate Weight 1 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 1 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 34-48 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT2: Wave 4 Adult Cross-sectional Replicate Weight 2 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 2 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 49-63 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT3: Wave 4 Adult Cross-sectional Replicate Weight 3 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 3 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 64-78 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT4: Wave 4 Adult Cross-sectional Replicate Weight 4 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 4 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 79-93 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT5: Wave 4 Adult Cross-sectional Replicate Weight 5 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 5 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 94-108 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT6: Wave 4 Adult Cross-sectional Replicate Weight 6 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 6 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 109-123 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT7: Wave 4 Adult Cross-sectional Replicate Weight 7 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 7 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 124-138 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT8: Wave 4 Adult Cross-sectional Replicate Weight 8 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 8 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 139-153 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT9: Wave 4 Adult Cross-sectional Replicate Weight 9 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 9 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 154-168 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT10: Wave 4 Adult Cross-sectional Replicate Weight 10 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 10 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 169-183 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT11: Wave 4 Adult Cross-sectional Replicate Weight 11 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 11 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 184-198 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT12: Wave 4 Adult Cross-sectional Replicate Weight 12 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 12 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 199-213 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT13: Wave 4 Adult Cross-sectional Replicate Weight 13 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 13 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 214-228 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT14: Wave 4 Adult Cross-sectional Replicate Weight 14 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 14 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 229-243 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT15: Wave 4 Adult Cross-sectional Replicate Weight 15 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 15 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 244-258 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT16: Wave 4 Adult Cross-sectional Replicate Weight 16 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 16 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 259-273 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT17: Wave 4 Adult Cross-sectional Replicate Weight 17 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 17 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 274-288 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT18: Wave 4 Adult Cross-sectional Replicate Weight 18 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 18 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 289-303 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT19: Wave 4 Adult Cross-sectional Replicate Weight 19 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 19 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 304-318 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT20: Wave 4 Adult Cross-sectional Replicate Weight 20 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 20 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 319-333 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT21: Wave 4 Adult Cross-sectional Replicate Weight 21 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 21 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 334-348 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT22: Wave 4 Adult Cross-sectional Replicate Weight 22 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 22 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 349-363 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT23: Wave 4 Adult Cross-sectional Replicate Weight 23 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 23 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 364-378 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT24: Wave 4 Adult Cross-sectional Replicate Weight 24 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 24 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 379-393 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT25: Wave 4 Adult Cross-sectional Replicate Weight 25 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 25 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 394-408 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT26: Wave 4 Adult Cross-sectional Replicate Weight 26 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 26 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 409-423 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT27: Wave 4 Adult Cross-sectional Replicate Weight 27 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 27 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 424-438 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT28: Wave 4 Adult Cross-sectional Replicate Weight 28 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 28 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 439-453 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT29: Wave 4 Adult Cross-sectional Replicate Weight 29 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 29 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 454-468 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT30: Wave 4 Adult Cross-sectional Replicate Weight 30 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 30 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 469-483 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT31: Wave 4 Adult Cross-sectional Replicate Weight 31 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 31 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 484-498 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT32: Wave 4 Adult Cross-sectional Replicate Weight 32 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 32 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 499-513 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT33: Wave 4 Adult Cross-sectional Replicate Weight 33 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 33 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 514-528 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT34: Wave 4 Adult Cross-sectional Replicate Weight 34 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 34 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 529-543 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT35: Wave 4 Adult Cross-sectional Replicate Weight 35 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 35 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 544-558 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT36: Wave 4 Adult Cross-sectional Replicate Weight 36 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 36 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 559-573 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT37: Wave 4 Adult Cross-sectional Replicate Weight 37 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 37 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 574-588 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT38: Wave 4 Adult Cross-sectional Replicate Weight 38 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 38 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 589-603 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT39: Wave 4 Adult Cross-sectional Replicate Weight 39 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 39 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 604-618 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT40: Wave 4 Adult Cross-sectional Replicate Weight 40 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 40 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 619-633 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT41: Wave 4 Adult Cross-sectional Replicate Weight 41 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 41 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 634-648 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT42: Wave 4 Adult Cross-sectional Replicate Weight 42 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 42 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 649-663 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT43: Wave 4 Adult Cross-sectional Replicate Weight 43 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 43 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 664-678 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT44: Wave 4 Adult Cross-sectional Replicate Weight 44 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 44 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 679-693 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT45: Wave 4 Adult Cross-sectional Replicate Weight 45 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 45 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 694-708 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT46: Wave 4 Adult Cross-sectional Replicate Weight 46 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 46 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 709-723 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT47: Wave 4 Adult Cross-sectional Replicate Weight 47 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 47 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 724-738 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT48: Wave 4 Adult Cross-sectional Replicate Weight 48 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 48 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 739-753 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT49: Wave 4 Adult Cross-sectional Replicate Weight 49 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 49 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 754-768 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT50: Wave 4 Adult Cross-sectional Replicate Weight 50 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 50 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 769-783 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT51: Wave 4 Adult Cross-sectional Replicate Weight 51 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 51 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 784-798 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT52: Wave 4 Adult Cross-sectional Replicate Weight 52 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 52 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 799-813 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT53: Wave 4 Adult Cross-sectional Replicate Weight 53 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 53 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 814-828 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT54: Wave 4 Adult Cross-sectional Replicate Weight 54 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 54 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 829-843 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT55: Wave 4 Adult Cross-sectional Replicate Weight 55 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 55 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 844-858 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT56: Wave 4 Adult Cross-sectional Replicate Weight 56 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 56 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 859-873 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT57: Wave 4 Adult Cross-sectional Replicate Weight 57 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 57 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 874-888 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT58: Wave 4 Adult Cross-sectional Replicate Weight 58 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 58 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 889-903 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT59: Wave 4 Adult Cross-sectional Replicate Weight 59 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 59 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 904-918 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT60: Wave 4 Adult Cross-sectional Replicate Weight 60 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 60 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 919-933 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT61: Wave 4 Adult Cross-sectional Replicate Weight 61 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 61 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 934-948 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT62: Wave 4 Adult Cross-sectional Replicate Weight 62 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 62 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 949-963 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT63: Wave 4 Adult Cross-sectional Replicate Weight 63 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 63 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 964-978 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT64: Wave 4 Adult Cross-sectional Replicate Weight 64 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 64 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 979-993 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT65: Wave 4 Adult Cross-sectional Replicate Weight 65 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 65 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 994-1008 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT66: Wave 4 Adult Cross-sectional Replicate Weight 66 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 66 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1009-1023 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT67: Wave 4 Adult Cross-sectional Replicate Weight 67 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 67 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1024-1038 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT68: Wave 4 Adult Cross-sectional Replicate Weight 68 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 68 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1039-1053 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT69: Wave 4 Adult Cross-sectional Replicate Weight 69 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 69 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1054-1068 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT70: Wave 4 Adult Cross-sectional Replicate Weight 70 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 70 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1069-1083 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT71: Wave 4 Adult Cross-sectional Replicate Weight 71 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 71 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1084-1098 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT72: Wave 4 Adult Cross-sectional Replicate Weight 72 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 72 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1099-1113 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT73: Wave 4 Adult Cross-sectional Replicate Weight 73 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 73 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1114-1128 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT74: Wave 4 Adult Cross-sectional Replicate Weight 74 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 74 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1129-1143 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT75: Wave 4 Adult Cross-sectional Replicate Weight 75 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 75 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1144-1158 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT76: Wave 4 Adult Cross-sectional Replicate Weight 76 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 76 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1159-1173 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT77: Wave 4 Adult Cross-sectional Replicate Weight 77 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 77 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1174-1188 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT78: Wave 4 Adult Cross-sectional Replicate Weight 78 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 78 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1189-1203 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT79: Wave 4 Adult Cross-sectional Replicate Weight 79 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 79 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1204-1218 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT80: Wave 4 Adult Cross-sectional Replicate Weight 80 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 80 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1219-1233 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT81: Wave 4 Adult Cross-sectional Replicate Weight 81 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 81 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1234-1248 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT82: Wave 4 Adult Cross-sectional Replicate Weight 82 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 82 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1249-1263 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT83: Wave 4 Adult Cross-sectional Replicate Weight 83 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 83 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1264-1278 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT84: Wave 4 Adult Cross-sectional Replicate Weight 84 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 84 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1279-1293 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT85: Wave 4 Adult Cross-sectional Replicate Weight 85 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 85 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1294-1308 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT86: Wave 4 Adult Cross-sectional Replicate Weight 86 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 86 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1309-1323 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT87: Wave 4 Adult Cross-sectional Replicate Weight 87 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 87 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1324-1338 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT88: Wave 4 Adult Cross-sectional Replicate Weight 88 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 88 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1339-1353 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT89: Wave 4 Adult Cross-sectional Replicate Weight 89 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 89 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1354-1368 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT90: Wave 4 Adult Cross-sectional Replicate Weight 90 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 90 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1369-1383 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT91: Wave 4 Adult Cross-sectional Replicate Weight 91 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 91 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1384-1398 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT92: Wave 4 Adult Cross-sectional Replicate Weight 92 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 92 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1399-1413 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT93: Wave 4 Adult Cross-sectional Replicate Weight 93 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 93 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1414-1428 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT94: Wave 4 Adult Cross-sectional Replicate Weight 94 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 94 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1429-1443 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT95: Wave 4 Adult Cross-sectional Replicate Weight 95 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 95 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1444-1458 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT96: Wave 4 Adult Cross-sectional Replicate Weight 96 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 96 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1459-1473 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT97: Wave 4 Adult Cross-sectional Replicate Weight 97 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 97 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1474-1488 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT98: Wave 4 Adult Cross-sectional Replicate Weight 98 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 98 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1489-1503 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT99: Wave 4 Adult Cross-sectional Replicate Weight 99 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 99 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1504-1518 (width: 15; decimal: 8)

Variable Type: numeric

R04_A_C04WGT100: Wave 4 Adult Cross-sectional Replicate Weight 100 for the Wave 4 Cohort

Wave 4 adult cross-sectional replicate weight 100 for variance estimation for the Wave 4 cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 33,644 valid cases out of 33,644 total cases.

Location: 1519-1533 (width: 15; decimal: 8)

Variable Type: numeric