

ICPSR 36498

Population Assessment of Tobacco and Health (PATH) Study [United States] Public-Use Files

United States Department of Health and Human Services. National Institutes of Health. National Institute on Drug Abuse

United States Department of Health and Human Services. Food and Drug Administration. Center for Tobacco Products

ICPSR Codebook for Wave 7: Adult - Wave 7
Cohort Cross-Sectional Weights

Inter-university Consortium for
Political and Social Research
P.O. Box 1248
Ann Arbor, Michigan 48106
www.icpsr.umich.edu

Terms of Use

The terms of use for this study can be found at:
<http://www.icpsr.umich.edu/icpsrweb/ICPSR/studies/36498/terms>

Information about Copyrighted Content

Some instruments administered as part of this study may contain in whole or substantially in part contents from copyrighted instruments. Reproductions of the instruments are provided as documentation for the analysis of the data associated with this collection. Restrictions on "fair use" apply to all copyrighted content. More information about the reproduction of copyrighted works by educators and librarians is available from the United States Copyright Office.

NOTICE

WARNING CONCERNING COPYRIGHT RESTRICTIONS

The copyright law of the United States (Title 17, United States Code) governs the making of photocopies or other reproductions of copyrighted material. Under certain conditions specified in the law, libraries and archives are authorized to furnish a photocopy or other reproduction. One of these specified conditions is that the photocopy or reproduction is not to be "used for any purpose other than private study, scholarship, or research." If a user makes a request for, or later uses, a photocopy or reproduction for purposes in excess of "fair use," that user may be liable for copyright infringement.

NOTE

This document does not currently comply with 508 standards. If you need an alternative means of access to any information, please contact NAHDAP at nahdap@icpsr.umich.edu. Let us know the nature of your accessibility problem, the Web address of the required information, and your contact information.

CODEBOOK DISPLAY NOTES

PATH STUDY

WEIGHTS FILE

ICPSR #36498-4331

1. ICPSR customized the display of variables in this codebook. There are no variables that contain a frequency table displaying value labels and unweighted counts. This is because all of the variables in this file are identification, sample design, or sampling weight variables for which univariate statistics are not meaningful.
2. Each variable contains the following statement:

“This variable has XXXXX valid cases out of XXXXX total cases.”

This statement describes the number of valid cases for a variable that would be used in the calculation of percentages in the frequency tables.

ICPSR 36498

Population Assessment of Tobacco and Health (PATH) Study [United States] Public-Use Files

Variable Description and Frequencies

Note: Frequencies displayed for the variables are not weighted. They are purely descriptive and may not be representative of the study population. Please review any sampling or weighting information available with the study.

Summary statistics (minimum, maximum, mean, median, and standard deviation) may not be available for every variable in the codebook. Conversely, a listing of frequencies in table format may not be present for every variable in the codebook either. However, all variables in the dataset are present and display sufficient information about each variable. These decisions are made intentionally and are at the discretion of the archive producing this codebook.

Wave 7: Adult - Wave 7 Cohort Cross-Sectional Weights

CASEID: Case Identification Number

Case Identification Number

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1-5 (width: 5; decimal: 0)

Variable Type: numeric

PERSONID: Participant ID

Participant ID Number.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 6-15 (width: 10; decimal: 0)

Variable Type: character

VARSTRAT: Stratum Indicator for Variance Estimation

Stratum indicator for variance estimation using the Taylor series approximation method.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 16-17 (width: 2; decimal: 0)

Variable Type: numeric

VARPSU: PSU Indicator for Variance Estimation

PSU indicator for variance estimation using the Taylor series approximation method.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 18-18 (width: 1; decimal: 0)

Variable Type: numeric

R07_A_C07WGT: Wave 7 Adult Cross-sectional Weight for the Wave 7 Cohort

Wave 7 adult cross-sectional full-sample weight for the Wave 7 Cohort.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 19-33 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT1: Wave 7 Adult Cross-sectional Replicate Weight 1 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 1 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 34-48 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT2: Wave 7 Adult Cross-sectional Replicate Weight 2 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 2 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 49-63 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT3: Wave 7 Adult Cross-sectional Replicate Weight 3 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 3 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 64-78 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT4: Wave 7 Adult Cross-sectional Replicate Weight 4 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 4 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 79-93 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT5: Wave 7 Adult Cross-sectional Replicate Weight 5 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 5 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 94-108 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT6: Wave 7 Adult Cross-sectional Replicate Weight 6 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 6 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 109-123 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT7: Wave 7 Adult Cross-sectional Replicate Weight 7 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 7 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 124-138 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT8: Wave 7 Adult Cross-sectional Replicate Weight 8 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 8 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 139-153 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT9: Wave 7 Adult Cross-sectional Replicate Weight 9 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 9 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 154-168 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT10: Wave 7 Adult Cross-sectional Replicate Weight 10 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 10 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 169-183 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT11: Wave 7 Adult Cross-sectional Replicate Weight 11 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 11 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 184-198 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT12: Wave 7 Adult Cross-sectional Replicate Weight 12 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 12 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 199-213 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT13: Wave 7 Adult Cross-sectional Replicate Weight 13 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 13 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 214-228 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT14: Wave 7 Adult Cross-sectional Replicate Weight 14 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 14 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 229-243 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT15: Wave 7 Adult Cross-sectional Replicate Weight 15 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 15 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 244-258 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT16: Wave 7 Adult Cross-sectional Replicate Weight 16 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 16 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 259-273 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT17: Wave 7 Adult Cross-sectional Replicate Weight 17 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 17 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 274-288 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT18: Wave 7 Adult Cross-sectional Replicate Weight 18 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 18 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 289-303 (width: 15; decimal: 8)
Variable Type: numeric

R07_A_C07WGT19: Wave 7 Adult Cross-sectional Replicate Weight 19 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 19 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 304-318 (width: 15; decimal: 8)
Variable Type: numeric

R07_A_C07WGT20: Wave 7 Adult Cross-sectional Replicate Weight 20 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 20 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 319-333 (width: 15; decimal: 8)
Variable Type: numeric

R07_A_C07WGT21: Wave 7 Adult Cross-sectional Replicate Weight 21 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 21 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 334-348 (width: 15; decimal: 8)
Variable Type: numeric

R07_A_C07WGT22: Wave 7 Adult Cross-sectional Replicate Weight 22 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 22 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 349-363 (width: 15; decimal: 8)
Variable Type: numeric

R07_A_C07WGT23: Wave 7 Adult Cross-sectional Replicate Weight 23 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 23 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 364-378 (width: 15; decimal: 8)
Variable Type: numeric

R07_A_C07WGT24: Wave 7 Adult Cross-sectional Replicate Weight 24 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 24 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 379-393 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT25: Wave 7 Adult Cross-sectional Replicate Weight 25 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 25 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 394-408 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT26: Wave 7 Adult Cross-sectional Replicate Weight 26 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 26 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 409-423 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT27: Wave 7 Adult Cross-sectional Replicate Weight 27 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 27 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 424-438 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT28: Wave 7 Adult Cross-sectional Replicate Weight 28 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 28 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 439-453 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT29: Wave 7 Adult Cross-sectional Replicate Weight 29 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 29 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 454-468 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT30: Wave 7 Adult Cross-sectional Replicate Weight 30 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 30 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 469-483 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT31: Wave 7 Adult Cross-sectional Replicate Weight 31 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 31 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 484-498 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT32: Wave 7 Adult Cross-sectional Replicate Weight 32 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 32 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 499-513 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT33: Wave 7 Adult Cross-sectional Replicate Weight 33 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 33 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 514-528 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT34: Wave 7 Adult Cross-sectional Replicate Weight 34 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 34 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 529-543 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT35: Wave 7 Adult Cross-sectional Replicate Weight 35 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 35 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 544-558 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT36: Wave 7 Adult Cross-sectional Replicate Weight 36 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 36 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 559-573 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT37: Wave 7 Adult Cross-sectional Replicate Weight 37 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 37 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 574-588 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT38: Wave 7 Adult Cross-sectional Replicate Weight 38 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 38 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 589-603 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT39: Wave 7 Adult Cross-sectional Replicate Weight 39 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 39 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 604-618 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT40: Wave 7 Adult Cross-sectional Replicate Weight 40 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 40 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 619-633 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT41: Wave 7 Adult Cross-sectional Replicate Weight 41 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 41 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 634-648 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT42: Wave 7 Adult Cross-sectional Replicate Weight 42 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 42 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 649-663 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT43: Wave 7 Adult Cross-sectional Replicate Weight 43 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 43 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 664-678 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT44: Wave 7 Adult Cross-sectional Replicate Weight 44 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 44 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 679-693 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT45: Wave 7 Adult Cross-sectional Replicate Weight 45 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 45 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 694-708 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT46: Wave 7 Adult Cross-sectional Replicate Weight 46 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 46 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 709-723 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT47: Wave 7 Adult Cross-sectional Replicate Weight 47 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 47 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 724-738 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT48: Wave 7 Adult Cross-sectional Replicate Weight 48 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 48 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 739-753 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT49: Wave 7 Adult Cross-sectional Replicate Weight 49 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 49 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 754-768 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT50: Wave 7 Adult Cross-sectional Replicate Weight 50 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 50 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 769-783 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT51: Wave 7 Adult Cross-sectional Replicate Weight 51 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 51 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 784-798 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT52: Wave 7 Adult Cross-sectional Replicate Weight 52 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 52 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 799-813 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT53: Wave 7 Adult Cross-sectional Replicate Weight 53 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 53 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 814-828 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT54: Wave 7 Adult Cross-sectional Replicate Weight 54 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 54 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 829-843 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT55: Wave 7 Adult Cross-sectional Replicate Weight 55 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 55 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 844-858 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT56: Wave 7 Adult Cross-sectional Replicate Weight 56 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 56 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 859-873 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT57: Wave 7 Adult Cross-sectional Replicate Weight 57 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 57 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 874-888 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT58: Wave 7 Adult Cross-sectional Replicate Weight 58 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 58 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 889-903 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT59: Wave 7 Adult Cross-sectional Replicate Weight 59 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 59 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 904-918 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT60: Wave 7 Adult Cross-sectional Replicate Weight 60 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 60 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 919-933 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT61: Wave 7 Adult Cross-sectional Replicate Weight 61 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 61 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 934-948 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT62: Wave 7 Adult Cross-sectional Replicate Weight 62 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 62 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 949-963 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT63: Wave 7 Adult Cross-sectional Replicate Weight 63 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 63 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 964-978 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT64: Wave 7 Adult Cross-sectional Replicate Weight 64 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 64 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 979-993 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT65: Wave 7 Adult Cross-sectional Replicate Weight 65 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 65 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 994-1008 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT66: Wave 7 Adult Cross-sectional Replicate Weight 66 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 66 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1009-1023 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT67: Wave 7 Adult Cross-sectional Replicate Weight 67 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 67 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1024-1038 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT68: Wave 7 Adult Cross-sectional Replicate Weight 68 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 68 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1039-1053 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT69: Wave 7 Adult Cross-sectional Replicate Weight 69 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 69 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1054-1068 (width: 15; decimal: 8)
Variable Type: numeric

R07_A_C07WGT70: Wave 7 Adult Cross-sectional Replicate Weight 70 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 70 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1069-1083 (width: 15; decimal: 8)
Variable Type: numeric

R07_A_C07WGT71: Wave 7 Adult Cross-sectional Replicate Weight 71 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 71 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1084-1098 (width: 15; decimal: 8)
Variable Type: numeric

R07_A_C07WGT72: Wave 7 Adult Cross-sectional Replicate Weight 72 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 72 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1099-1113 (width: 15; decimal: 8)
Variable Type: numeric

R07_A_C07WGT73: Wave 7 Adult Cross-sectional Replicate Weight 73 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 73 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1114-1128 (width: 15; decimal: 8)
Variable Type: numeric

R07_A_C07WGT74: Wave 7 Adult Cross-sectional Replicate Weight 74 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 74 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1129-1143 (width: 15; decimal: 8)
Variable Type: numeric

R07_A_C07WGT75: Wave 7 Adult Cross-sectional Replicate Weight 75 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 75 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1144-1158 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT76: Wave 7 Adult Cross-sectional Replicate Weight 76 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 76 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1159-1173 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT77: Wave 7 Adult Cross-sectional Replicate Weight 77 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 77 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1174-1188 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT78: Wave 7 Adult Cross-sectional Replicate Weight 78 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 78 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1189-1203 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT79: Wave 7 Adult Cross-sectional Replicate Weight 79 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 79 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1204-1218 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT80: Wave 7 Adult Cross-sectional Replicate Weight 80 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 80 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1219-1233 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT81: Wave 7 Adult Cross-sectional Replicate Weight 81 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 81 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1234-1248 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT82: Wave 7 Adult Cross-sectional Replicate Weight 82 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 82 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1249-1263 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT83: Wave 7 Adult Cross-sectional Replicate Weight 83 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 83 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1264-1278 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT84: Wave 7 Adult Cross-sectional Replicate Weight 84 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 84 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1279-1293 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT85: Wave 7 Adult Cross-sectional Replicate Weight 85 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 85 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1294-1308 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT86: Wave 7 Adult Cross-sectional Replicate Weight 86 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 86 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1309-1323 (width: 15; decimal: 8)
Variable Type: numeric

R07_A_C07WGT87: Wave 7 Adult Cross-sectional Replicate Weight 87 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 87 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1324-1338 (width: 15; decimal: 8)
Variable Type: numeric

R07_A_C07WGT88: Wave 7 Adult Cross-sectional Replicate Weight 88 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 88 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1339-1353 (width: 15; decimal: 8)
Variable Type: numeric

R07_A_C07WGT89: Wave 7 Adult Cross-sectional Replicate Weight 89 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 89 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1354-1368 (width: 15; decimal: 8)
Variable Type: numeric

R07_A_C07WGT90: Wave 7 Adult Cross-sectional Replicate Weight 90 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 90 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1369-1383 (width: 15; decimal: 8)
Variable Type: numeric

R07_A_C07WGT91: Wave 7 Adult Cross-sectional Replicate Weight 91 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 91 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1384-1398 (width: 15; decimal: 8)
Variable Type: numeric

R07_A_C07WGT92: Wave 7 Adult Cross-sectional Replicate Weight 92 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 92 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1399-1413 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT93: Wave 7 Adult Cross-sectional Replicate Weight 93 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 93 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1414-1428 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT94: Wave 7 Adult Cross-sectional Replicate Weight 94 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 94 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1429-1443 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT95: Wave 7 Adult Cross-sectional Replicate Weight 95 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 95 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1444-1458 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT96: Wave 7 Adult Cross-sectional Replicate Weight 96 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 96 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1459-1473 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT97: Wave 7 Adult Cross-sectional Replicate Weight 97 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 97 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1474-1488 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT98: Wave 7 Adult Cross-sectional Replicate Weight 98 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 98 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1489-1503 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT99: Wave 7 Adult Cross-sectional Replicate Weight 99 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 99 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1504-1518 (width: 15; decimal: 8)

Variable Type: numeric

R07_A_C07WGT100: Wave 7 Adult Cross-sectional Replicate Weight 100 for the Wave 7 Cohort

Wave 7 adult cross-sectional replicate weight 100 for variance estimation for the Wave 7 Cohort created using balanced repeated replication with a Fay's adjustment factor of 0.3.

This variable has 29,780 valid cases out of 29,780 total cases.

Location: 1519-1533 (width: 15; decimal: 8)

Variable Type: numeric