Cloning of Human TFEB and Analysis of Its Nutrient-Dependent Subcellular Localization in HeLa Cells

Siyuan Jiang Ziyan Song
Molecular Biology Laboratory Course

Version: Working draft
Date: October 23, 2026

Abstract

Transcription factor EB (TFEB) is a nutrient-responsive transcription factor that coordinates autophagy and lysosomal biogenesis. This proposal aims to construct wild-type human TFEB isoform 1 fused to C-terminal EGFP in pEGFP-N1 and use it to visualize TFEB localization in HeLa cells under nutrient-replete, starvation, and mTOR-inhibition conditions. The central hypothesis is that nutrient deprivation or pharmacological inhibition of mTOR will increase nuclear accumulation of the reporter. The construct will be generated by PCR amplification of a defined human TFEB coding sequence, restriction-enzyme cloning into a mammalian expression vector, bacterial screening, and Sanger sequencing. Transfected cells will be evaluated by fluorescence microscopy and immunoblotting. All experimental outcomes described here are expected results of a proposed experiment.

Keywords: TFEB; autophagy; lysosome; mTORC1; EGFP fusion protein; subcellular localization

1 Rationale and Research Basis

TFEB is a member of the MiT/TFE transcription-factor family and a major transcriptional regulator of the autophagy–lysosome pathway. Under nutrient-rich conditions, mTORC1-dependent phosphorylation favors cytoplasmic retention of TFEB. During nutrient starvation or mTOR inhibition, TFEB accumulates in the nucleus and activates genes involved in lysosomal biogenesis and cellular clearance [5, 6]. Nutrient-dependent control of nuclear export also contributes to TFEB distribution [2].

This project connects molecular cloning with a visible cell-biological phenotype. Its primary readout is the nuclear-to-cytoplasmic distribution of a fluorescent TFEB fusion. The study will test whether the sequence-verified reporter exhibits the expected nutrient-dependent redistribution in HeLa cells.

1.1 Research Objectives and Hypothesis

The first objective is to construct and sequence-verify an EGFP-tagged TFEB mammalian expression plasmid. The second is to compare its subcellular localization under normal culture, starvation, and mTOR-inhibitor conditions. The hypothesis is that starvation and mTOR inhibition increase the nuclear-to-cytoplasmic fluorescence ratio relative to the vehicle-treated control.

2 Experimental Design

2.1 Overall Workflow

The study proceeds from reference-sequence retrieval and primer design to PCR amplification, plasmid construction, clone verification, transient transfection, and localization analysis. Sequence verification is the gate between molecular cloning and cell-based experiments.

2.2 Construct Strategy and Materials

The selected reporter is wild-type human TFEB isoform 1 fused to C-terminal EGFP (TFEB–EGFP), expressed from pEGFP-N1 under the CMV promoter. This architecture follows the HeLa localization study of Roczniak-Ferguson et al. and its deposited construct, pEGFP-N1-TFEB (Addgene #38119) [1, 5]. It replaces the initial outline's provisional pEGFP-C1 strategy. The choice follows a directly relevant experimental precedent; it does not establish that fluorescent tagging leaves endogenous TFEB regulation unchanged.

The TFEB reference is RefSeq NM_007162.3, encoding NP_009093.1 (476 amino acids). Its CDS occupies transcript positions 303–1733 (1-based, inclusive): 1,431 nucleotides including the terminal TGA, or 1,428 nucleotides without that stop codon [4]. Computational comparison confirmed that its CDS and protein are identical to historical NM_007162.2, the accession listed for Addgene #38119. The older transcript has ten additional terminal adenines outside the CDS. The independently retrieved NP_009093.1 protein matches the translated CDS.

The empty-vector sequence is fixed to GenBank U55762.1, a 4,733-bp circular pEGFP-N1 reference submitted by Clontech [3]. The published plasmid is a construct precedent, whereas this assignment retains PCR and cloning into the empty backbone. We have not obtained or verified the complete sequence of Addgene #38119 or a physical vector stock. Kanamycin is the bacterial selection marker; neomycin/G418 is the mammalian selection marker. Stable mammalian selection is not required for the planned transient-expression assay.

For the design, a synthetic cDNA template clone matching the native NM_007162.3 CDS, including its stop codon and without codon optimization, is proposed. No material has been purchased or experimentally verified. The fusion insert will retain the TFEB initiation ATG but omit its terminal TGA. The complete EGFP coding region is intended to follow in frame and retain its terminal TAA; its ATG will encode an internal methionine. The initiation context and junction sequence still require explicit design and validation. The 1,428-nt TFEB coding portion is not the final tailed amplicon or complete fusion length.

The remaining materials are high-fidelity DNA polymerase, two compatible restriction enzymes, DNA ligase, competent *Escherichia coli* DH5α, and kanamycin-containing selection medium. A final plasmid map will be generated after the junctions and all relevant vector features have been verified.

2.3 Sequence Retrieval and Primer Design

The versioned TFEB and empty-vector records, extracted CDS and protein, input hashes, and reference-verification script are archived with the project. HindIII/KpnI is the published cloning pair and will be evaluated first [1, 5]; it is not yet a validated pair for the designed junctions. Each cloning primer will be reported in the $5'$ to $3'$ direction and annotated by protective bases, restriction site, any frame-adjustment or linker bases, and template-annealing sequence. The selected pair must cut the

intended backbone without cutting the TFEB insert internally. Primer sequences, annealing-region melting temperatures, final junctions, and the complete fusion translation remain to be determined.

2.4 PCR Amplification and Restriction Cloning

The TFEB insert will be amplified with a high-fidelity polymerase. A no-template control and an appropriate amplification control will accompany the reaction. The intended amplicon will be purified, and both insert and vector will be digested and purified before ligation. An initial insert-to-vector molar ratio of approximately 3:1 is proposed. The ligation will be transformed into competent DH5α cells, followed by selection using the backbone's confirmed antibiotic marker. The reaction composition, thermal program, digestion conditions, and transformation details remain to be specified for the selected reagents.

2.5 Clone Screening and Sequence Verification

Candidate colonies will be screened by colony PCR and diagnostic restriction digestion. Plasmid DNA from positive clones will be purified for Sanger sequencing with vector-flanking and, where necessary, internal primers. A construct will be accepted only when the complete TFEB insert, both junctions, fusion reading frame, and terminal stop codon match the intended sequence without unintended substitutions, insertions, or deletions. An expected-size band alone will not establish sequence identity.

2.6 Cell Culture, Transfection, and Treatment

After sequence verification, the pEGFP-N1-based TFEB–EGFP plasmid will be transiently transfected into HeLa cells. The proposed comparison includes normal culture with vehicle, starvation medium, and Torin1 treatment. The final methods will identify the culture medium, transfection reagent, cell density, DNA amount, and treatment concentration and duration. Empty pEGFP-N1 expressing unfused EGFP and mock-transfected cells will provide expression and background controls. A matching vehicle control will accompany Torin1 treatment.

2.7 Imaging and Localization Analysis

After treatment, cells will be fixed and counterstained with DAPI. Fluorescence images will be collected using consistent acquisition settings. Nuclear and cytoplasmic regions will be defined for each eligible transfected cell, and background-corrected mean fluorescence will be used to calculate the nuclear-to-cytoplasmic ratio:

$$R_{\mathrm{N/C}} = \frac{I_{\mathrm{nucleus}} - I_{\mathrm{background}}}{I_{\mathrm{cytoplasm}} - I_{\mathrm{background}}}. \tag{1}$$

At least three independent experiments are planned. The number of cells per replicate, eligibility criteria, and handling of background-level cytoplasmic signals will be specified before analysis. Cells within one experiment will not be treated as independent biological replicates. The final statistical comparison will account for the experimental unit and the planned control-versus-treatment comparisons.

2.8 Expression Validation

An anti-GFP Western blot will assess fusion-protein expression and possible cleavage products. The expected molecular mass will be calculated from the finalized construct. The final protocol will specify protein extraction, loading control, antibody information, and interpretation of additional bands. Expression validation will be considered alongside the localization data.

3 Expected Results and Criteria for Success

3.1 Molecular Cloning

PCR is expected to produce an amplicon matching the finalized TFEB insert design, with no corresponding product in the no-template control. Colony PCR and diagnostic digestion should yield the predicted fragment pattern. Exact amplicon and restriction-fragment lengths will be recorded with the completed sequence design.

The decisive criterion is Sanger confirmation of the complete intended fusion construct. Successful cloning requires correct insert identity and orientation, both junctions, the continuous open reading frame, and the intended termination codon. The final report will include a plasmid map, a cloning workflow, and explicitly labeled schematic predictions for the gel patterns.

3.2 Expression and Nutrient-Dependent Localization

The reporter is expected to show a higher nuclear-to-cytoplasmic fluorescence ratio after starvation or Torin1 treatment than in the vehicle-treated condition. An anti-GFP-reactive band consistent with the finalized fusion-protein size would support expression of the intended product.

Evidence for nutrient responsiveness will require a reproducible treatment effect across independent experiments, supported by expression and imaging controls. The magnitude and uncertainty of the localization change will be reported alongside the statistical comparison. A failure to observe redistribution would prompt checks of sequence, fusion orientation, expression level, treatment response, and imaging conditions.

4 Discussion and Limitations

The proposed design links molecular construction to a testable localization phenotype. Its main limitation is that overexpression and fluorescent tagging can influence TFEB regulation. Fusion orientation therefore requires explicit justification, and a reporter response should not be equated with a complete description of endogenous TFEB behavior.

The localization assay examines redistribution rather than autophagic flux or lysosomal function directly. Changes in nuclear localization alone will not establish increased autophagic degradation. Follow-up studies could compare reporter localization with endogenous TFEB or test selected downstream targets. These extensions are outside the initial cloning-and-localization objective.

Author Contributions

Siyuan Jiang will lead TFEB sequence retrieval, vector selection, cloning and primer design, molecular methods, plasmid and workflow diagrams, clone-verification criteria, and the related literature review. Ziyan Song will lead the abstract, biological rationale, HeLa-cell experiments, localization and expression analysis, cell-related expected results, discussion, and the related literature review. Both authors will review the construct-to-cell-assay interface and the integrated report.

AI Use Declaration

AI assistance was used to organize the supplied outline, review literature and database records, propose the TFEB reference, vector and tag orientation, write and review a reference-sequence verification script, and prepare the LaTeX manuscript. The authors will review the final scientific content, sequence design, experimental parameters, references, and contribution statement before submission.

References

[1] Addgene. *pEGFP-N1-TFEB, plasmid #38119*. URL: https://www.addgene.org/38119/ (visited on 10/23/2026).

[2] Gennaro Napolitano et al. "mTOR-dependent phosphorylation controls TFEB nuclear export". In: *Nature Communications* 9, 3312 (2018). DOI: 10.1038/s41467-018-05862-6.

[3] NCBI GenBank. *Cloning vector pEGFP-N1, complete sequence: U55762.1*. URL: https://www.ncbi.nlm.nih.gov/nuccore/U55762.1 (visited on 10/23/2026).

[4] NCBI RefSeq. *Homo sapiens transcription factor EB: NM_007162.3 and NP_009093.1*. URL: https://www.ncbi.nlm.nih.gov/nuccore/NM_007162.3 (visited on 10/23/2026).

[5] Agnes Roczniak-Ferguson et al. "The transcription factor TFEB links mTORC1 signaling to transcriptional control of lysosome homeostasis". In: *Science Signaling* 5.228 (2012), ra42. DOI: 10.1126/scisignal.2002790.

[6] Carmine Settembre et al. "A lysosome-to-nucleus signalling mechanism senses and regulates the lysosome via mTOR and TFEB". In: *The EMBO Journal* 31.5 (2012), pp. 1095–1108. DOI: 10.1038/emboj.2012.32.