Trend entired Briaminger

Endineering Balanings

The graphical suppresentation of an object containing details like size, shape etc. on a 2D plane is called Engineering Drawing."

Conic Section

Focus: The fixed point es called focus.

Dinectorin: The fixed storaget line is called directoria

Lester:- The point at which conic section intersects the axis is called restex. all with a let me

Eccent Micity:

e = Distance of the point from the focus VF

VA

Distance of the point from the directrix VA

If e=1 for parabola ex1 for ellipse.

e>1 for hypeopla

parabola : Ellipse : Hyperbola

HB = Harner & Boldness & rounder to sin Barrer.

4 =

24 =

O. Constance a pariabola, when the distance of the focus strom the distance of the tangent and normal to me course at a point Ho room the districtality of the formypier

Sol : Porocedwie :- 1 m of again, of the sale

Co Daraw a hosizontal Alne MB. Messay

On Doraw the resitted line CD Pie, distectsion.

(M) Mark Locus (B) at a distance of 50mm from The drojectoiry.

(Go) Mark vertex (O) in the middle of AF!

(a) From V draw a veritical line, se VE=VF)

CVB John NE and entind. From V, devide the equal numbers of parts say each 10mm.

(M) From 11213,..., draw the verifical lines to . Introsect at AE line and mark or 1',2',3'.

(Child take 11 as stadius and F as centure to cut 11' line either side of AB line.


(PX) 221, 331, ... as madiup and Filhas centre to cut 221, 331, lines:

(x) Join all arcs along with 1, we get parallel


To draw tangent and normal:

(8) Mark P on the curve of a distance of 40min from the directals. Join PF.


(10) From F, draw a peorpenderular line to PF. Potensect at CD line, mark as t. (in) Join TP and entend, r.c, tangent TT!




D. Constanct a panabola, when the distance of the focus from the dissectable is 40. Also draw a tangent and normal to the curve at a distance of from the dissectable.



3. Construct a hyperbola, when the distance of the focus from the distance of the focus of the eccentricity of 3. Also draw a tangent and norm at to the curve at a distance of 40 from the directaria.




(4) Construct a hyperbola when the dirtance of The focus from the disrectorin is 70. and eccentrations 5 4. Also draw a tangent and normal to the curve at a distance of 40 from the dispectally B A 40 D




6. Enstruct an ellipse when the distance blu the tou and dispectary is 50. and eccentricity 3. Also draw a normal and tangent to the curve at a point yourn from the disectain. Anectary. UF LUA 1 A 40 50 D


a construct an ellepse when the distance blw the focus and the despectater is 35mm and eccentaricity is 3. Also draw the targent and normal at any point p' on the curve using divectoria. 14 30 35 p



Rectareular Hyperbola = e= 2.

3. Contriuct a nectargular hypeoloolq when a point ip, and it is at a distance of 18mm and 34mm from two asymptotes. Also draw a tangent to the curve at a point 20mm from an asymptote.





Involuter:


Involute es a single curved line toraced out by an end of a staring when unwound it self from a Stagget line (a) circle (on polygon, the staling being always kept tout.

m. Draw the involute of a sle of 25mm side.


(13. Draw the envolute of a square of side 20mm. 50% Tap = 360 = 120°, Square = 360 = 90° | Hepta, octor

pent a = 360 = 72° , Hexa = 310 = 60°

hone, dra-



(B) Draw the involute of a pentagon of 20mm side.



(14). Praw the involute of a hexagon of 20mm side. toler of yet to the till a st holy probe other Almes since is to consist 83 the string I point out of surely in it In fact you me allow white of the forbit spins Francis and de conto the (et) 7 '012 1111KC . 21 W. 40 0 10 10000 Partin 11. 11 the 15 to all Ps I men brown of a file winds ignity faith of our many 19 20 quibile growal graham or or of the and reform has a first dim as a second out what it was a work of our fine The filter of the man 100 H 10 4 12 La state from manyor and the

Sycloid:

Cycloid is a curve traced by a point on the circum. serence of a circle which rolls along a straight line without slipping.

Epicy chili-

Epicycloid is a curve traced by a point on the circum ference of a circle which rolls on the outside of another circle without slipping.

HABOCA Cloud:-

Hypocycloss is a curve traced by a point on the circumsterience of a circle which rolls in the inside of another circle without slipping.

(B). A cincle of yomm diameter nolls on a line for one nevolution clockwise. Doyaw the locus of a point or the cincle which is incontact with the line. Also draw the taugent and normal to the curve at a point 35mm from the directing line.

Sol= Polocedure:

(9) Draw a clade of 20 mm radiup.

(%) Devide the circle in to 12 equal parts. i.e, $\frac{360}{12} = 30^{\circ}$ each part, and marker the points names as $1,2,3,\dots, 1,p$.

(FM). From 'p', draw a horizontal line with the length of Td. i.e, T.x40=125.6 = 126mm.


(9v) From , A' draw a vertical line of 4 omm.

(4) From 7,8,9,10,11 draw the horizontal lines to AB line.

some positives to a mind simile

(3) Device PA into 12 equal parts.


9.e, 126 = 10.5mm each part.



Epicy closed : 1 colome of explications of O Dolaw an epicyclord of a cescle of Homm drameter which sollo outside on another circle of Izarmy

digmeter og no revolution dockwise. Dergu d tangent and nonmal to let at any point on the curvo

(Epicyclopa).





2). Construct an epicydoid generated by a rolling of of 150mm. Daaw the tangent and nogmal at any point on the epicycloid.

nt on the epicyclopd. R = 750, R = 750


O Dogaw a hypocyclopd of Homm diameter which anolls inside of another ciarde of 100mm for one revolution dos one counter clockwise. Deraw a tayen and normal to it at any point on the curve. D = 160, R = 80 $\theta = 360 \times \frac{8}{R} = 360 \times \frac{20}{80} = 90^{\circ} \left| \frac{\theta}{12} \right| = 1.$

d=40, 8=20






 $\theta = 360 \times \frac{7}{R} = 360 \times \frac{30}{190}$ d = 60 R = 100 $\frac{108}{12} = \frac{108}{12}$



Sil. Hypotycloid when, radius of the directing of rule is twice the radius of the generating ciocle. Radius of the generating ciocle.

$$\theta = 360 \times \frac{8}{R} = 360 \times \frac{25}{70_2} = 180^{\circ}$$
 $\frac{\Theta}{12} = \frac{180}{12} = 15^{\circ}$



Concentric Ciorder Method:

A. A Howey bed in a botanical gooden is elliptical in shape. Major and minor axis are 120mm and 90mm respectively. Draw the profile of the Howey bed.

Maros axis = 120 mm R = 60 mm Minos axis = 90 mm 8=45 mm at the second of the second of the second A 11 Henrolli Elmis Dietaria

Sellicht of town the coperting the profess of ellicht of the profess of ellicht of town the town the profile parties the store the profile

MANUAL TO ALL THAT AND THE PARTY OF THE PART