

## Some visually interesting non-standard quaternion fractal sets

S. Halayka \*

Brook Bay Regina, SK, Canada S4N 1M7

### ARTICLE INFO

#### Article history:

Accepted 21 October 2008

### ABSTRACT

The quaternion Julia set's standard iterative function is replaced with several others, most of which seem to be previously undocumented. The results are reproduced here mainly for the sake of visual interest.

© 2008 Elsevier Ltd. All rights reserved.

### 1. Introduction

The quaternion Julia set [1] can be produced by marching along a finite lattice of regularly spaced points in  $\mathbf{R}^3$ , testing at each one to see if the iterative function

$$Z = Z^2 + C \quad (1)$$

causes the magnitude  $|Z| = \sqrt{Z_x^2 + Z_y^2 + Z_z^2 + Z_w^2}$  to exceed a threshold value while undergoing a limited number of iterations. Only when the magnitude remains below or equal to the threshold is a point within the set. For all sets given here, the magnitude threshold value is 4, the iteration limit is 8, and the lattice extends from  $-1.5$  to  $1.5$  along all three spatial dimensions.

For each lattice point  $x, y, z$ , the initial value of  $Z$  is:

$$Z_x = x, \quad (2)$$

$$Z_y = y, \quad (3)$$

$$Z_z = z, \quad (4)$$

and  $Z_w$  is some arbitrary value. For all sets given here, initially  $Z_w = 0$ . All four components of the quaternion  $C$  are also arbitrary.

Using the standard product operation described in [2,3], the squaring of  $Z$  produces a quaternion in the form of a scalar and a 3-vector:

$$t = t' = Z_x, \quad (5)$$

$$\vec{V} = \vec{V}' = Z_y, Z_z, Z_w, \quad (6)$$

$$Z \cdot Z = tt' - \vec{V} \cdot \vec{V}', \quad t\vec{V}' + t'\vec{V} + \vec{V} \times \vec{V}', \quad (7)$$

$$(Z \cdot Z)_x = Z_x Z_x - (Z_y Z_y + Z_z Z_z + Z_w Z_w), \quad (8)$$

$$(Z \cdot Z)_y = Z_x Z_y + Z_x Z_y + (Z_z Z_w - Z_w Z_z), \quad (9)$$

$$(Z \cdot Z)_z = Z_x Z_z + Z_x Z_z + (Z_w Z_y - Z_y Z_w), \quad (10)$$

$$(Z \cdot Z)_w = Z_x Z_w + Z_x Z_w + (Z_y Z_z - Z_z Z_y). \quad (11)$$

The non-commutativity of the product operation is important in the case of inequal operands, ex: if  $A \neq B$ , then  $A \cdot B \neq B \cdot A$ .

Quaternion addition is performed piecewise, ex:  $Z^2 + C = (Z \cdot Z)_x + C_x, \dots$

\* Tel.: +1 (306) 535 4192.

E-mail address: [shalayka@gmail.com](mailto:shalayka@gmail.com).

## 2. Standard and non-standard sets visualized

Where  $Z = Z^2 + C$ , and  $C_{xyzw} = 0.3, 0.5, 0.4, 0.2$ , all points within the standard quaternion Julia set are visualized in Fig. 1 using a crude surface approximation.

As mentioned in [4], altering the standard iterative function by increasing the exponent beyond 2 provides interesting results. See Fig. 2 for  $Z = Z^4 + C$ , and  $C_{xyzw} = 0.3, 0.5, 0.4, 0.2$ .

Visually interesting sets can also be produced through the use of functions such as  $\sin Z$  [2]:

$$t = Z_x, \quad (12)$$

$$\vec{V} = Z_y, \quad Z_z, \quad Z_w, \quad (13)$$

$$|\vec{V}| = \sqrt{Z_y^2 + Z_z^2 + Z_w^2}, \quad (14)$$

$$\text{norm}(Z) = |Z|^2 = t^2 + \vec{V} \cdot \vec{V}, \quad (15)$$

$$\text{inv}(Z) = t/\text{norm}(Z), \quad -\vec{V}/\text{norm}(Z), \quad (16)$$

$$\sin Z = \sin t \cosh |\vec{V}|, \quad \cos t \sinh |\vec{V}| \vec{V}/|\vec{V}|, \quad (17)$$

$$\cos Z = \cos t \cosh |\vec{V}|, \quad -\sin t \sinh |\vec{V}| \vec{V}/|\vec{V}|, \quad (18)$$

$$\exp Z = \exp t \cos |\vec{V}|, \quad \exp t \sin |\vec{V}| \vec{V}/|\vec{V}|, \quad (19)$$

$$\sinh Z = \sinh t \cos |\vec{V}|, \quad \cosh t \sin |\vec{V}| \vec{V}/|\vec{V}|, \quad (20)$$

$$\cosh Z = \cosh t \cos |\vec{V}|, \quad \sinh t \sin |\vec{V}| \vec{V}/|\vec{V}|. \quad (21)$$




Fig. 1.  $Z = Z^2 + C$ , and  $C_{xyzw} = 0.3, 0.5, 0.4, 0.2$ .




Fig. 2.  $Z = Z^4 + C$ , and  $C_{xyzw} = 0.3, 0.5, 0.4, 0.2$ .

See Fig. 3 for  $Z = \sin Z + C \cdot \sin Z$ , and  $C_{xyzw} = 0.3, 0.5, 0.4, 0.2$ .

See Fig. 4 for  $Z = \sin Z + C \cdot \sin Z$ , and  $C_{xyzw} = -0.3, 0.5, 0.7, 0.2$ .

See Fig. 5 for  $Z = \cos Z + C \cdot \cos Z$ , and  $C_{xyzw} = 0.3, 0.5, 0.4, 0.2$ .

See Fig. 6 for  $Z = \exp(Z^2) + C$ , and  $C_{xyzw} = 0.3, 0.5, 0.4, 0.2$ , where the complement of the set is used for visualization.

See Fig. 7 for  $Z = \text{inv}(\sinh Z) + C \cdot \text{inv}(\sinh Z)$ , and  $C_{xyzw} = 1, 1, 1, 1$ .

See Fig. 8 for  $Z = C \cdot (\text{inv}(\sinh Z) \cdot \cosh Z)$ , and  $C_{xyzw} = 1, 1, 1, 1$ , where the complement of the set is used for visualization.

### 3. Comments

Without straying too far from the present subject of fractal sets, it should be noted that the quaternions also have a wide range of practical applications.

In terms of group theory, the unit quaternions represent the special unitary Lie group  $SU(2)$  [5], which is an integral part of the standard model of particle physics [6,7]. As well, the algebras related to the exceptional Lie groups  $E_7$  are constructed using the quaternions (and the 8-component octonions) [5], which have been used by El Naschie to investigate  $E^{(\infty)}$  theory [8,9].

The unit quaternion representation of rotation in 3-dimensional space is also commonly used in computer science [10], including virtual camera control [11], inverse kinematics [12], and even encryption [13].




Fig. 3.  $Z = \sin Z + C \cdot \sin Z$ , and  $C_{xyzw} = 0.3, 0.5, 0.4, 0.2$ .







Fig. 4.  $Z = \sin Z + C \cdot \sin Z$ , and  $C_{xyzw} = -0.3, 0.5, 0.7, 0.2$ .




**Fig. 5.**  $Z = \cos Z + C \cdot \cos Z$ , and  $C_{xyzw} = 0.3, 0.5, 0.4, 0.2$ .



**Fig. 6.** The complement of  $Z = \exp(Z^2) + C$ , and  $C_{xyzw} = 0.3, 0.5, 0.4, 0.2$ .



**Fig. 7.**  $Z = \text{inv}(\sinh Z) + C \cdot \text{inv}(\sinh Z)$ , and  $C_{xyzw} = 1, 1, 1, 1$ .



**Fig. 8.** The complement of  $Z = C \cdot (\text{inv}(\sinh Z) \cdot \cosh Z)$ , and  $C_{xyzw} = 1, 1, 1, 1$ .

The work of Griffin and Joshi [14,15] is recommended to those wishing to explore the octonion Julia set.

Thank you to Douglas Sweetser and Paul Bourke for their encouragement and clarification on quaternion algebra, fractals, and visualization in general.

Thank you to James DeRosa and LazART, Inc. for generously providing a sculpture of the set  $Z = \sin Z + C \cdot \sin Z$ .

Special thanks to Godwin Vickers. *Avec tous mes remerciements.*

## References

- [1] Norton A. Generation and display of geometric fractals in 3-D. In: SIGGRAPH'82: proceedings of the 9th annual conference on computer graphics and interactive techniques; 1982. ISBN: 0-89791-076-1.
- [2] Sweetser D. A quaternion algebra tool set; 1997. Available from: <http://world.std.com/sweetser/quaternions/intro/tools/tools.html>.
- [3] Crane K. Ray tracing quaternion Julia sets on the GPU; 2005. Available from: [http://graphics.cs.uiuc.edu/svn/kcrane/web/project\\_qjulia.html](http://graphics.cs.uiuc.edu/svn/kcrane/web/project_qjulia.html).
- [4] Bourke P. Quaternion Julia fractals; 2001. Available from: <http://local.wasp.uwa.edu.au/pbourke/fractals/quatjulia/>.
- [5] Baez JC. The octonions. Bull Am Math Soc 2002;39.
- [6] Anderson R, Joshi GC. Interpreting mathematics in physics: charting the applications of SU(2) in 20th century physics. Chaos, Solitons & Fractals 2008;36(2).
- [7] Pokorski S. Gauge field theories; 2000. ISBN: 978-0521478168.
- [8] El Naschie MS. Freudenthal magic square and its dimensional implication for  $\tilde{\alpha}_0 \simeq 137$  and high energy physics. Chaos, Solitons & Fractals 2008;36(3).
- [9] El Naschie MS. Deriving the largest expected number of elementary particles in the standard model from the maximal compact subgroup  $H$  of the exceptional Lie group  $E_{7(-5)}$ . Chaos, Solitons & Fractals 2008;38(4).
- [10] Dunn F, Parberry I. 3D math primer for graphics and game development; 2002. ISBN: 1-55622-911-9.
- [11] Shoemaker K. ARCBALL: a user interface for specifying three-dimensional orientation using a mouse. In: Proceedings of the conference on graphics interface'92; 1992. ISBN: 0-9695338-1-0.
- [12] Aydin Y, Kucuk S. Quaternion based inverse kinematics for industrial robot manipulators with Euler Wrist. In: 2006 IEEE international conference on mechatronics; 2006. doi:10.1109/ICMECH.2006.252591.
- [13] Nagase T, Komata M, Araki T. Secure signals transmission based on quaternion encryption scheme. In: 8th international conference on advanced information networking and applications (AINA'04), vol. 2; 2004. doi:10.1109/AINA.2004.1283751.
- [14] Griffin CJ, Joshi GC. Octonionic Julia sets. Chaos, Solitons & Fractals 1992;2(1).
- [15] Griffin CJ, Joshi GC. Transition points in octonionic Julia sets. Chaos, Solitons & Fractals 1993;3(1).