

On the Monte Carlo simulation of anisotropic Newtonian gravitation

S. Halayka*

Monday 11th August, 2025 15:09

Abstract

This paper contains a short introduction to anisotropic Newtonian gravitation. The main focus is on some C++ code.

1 Introduction

$$\alpha = \frac{\beta(R + \epsilon) - \beta(R)}{\epsilon}. \quad (1)$$

$$g = \frac{-\alpha}{r^2}. \quad (2)$$

$$a_N = \sqrt{\frac{nG\hbar \log 2}{4k\pi R^4}}, \quad (3)$$

$$v_N = \sqrt{a_N R}. \quad (4)$$

$$v_{\text{flat}} = x v_N \quad (5)$$

where $x = 2$ for example.

$$a_{\text{flat}} = \frac{v_{\text{flat}}^2}{R} = \frac{g R \hbar \log 2}{k 2 \pi M}. \quad (6)$$

$$g_N = \frac{a_N k 2 \pi M}{R \hbar \log 2}. \quad (7)$$

$$a_{\text{flat}} \propto g. \quad (8)$$

$$a_{\text{ratio}} = \frac{a_{\text{flat}}}{a_N}. \quad (9)$$

$$g_{\text{ratio}} = \frac{g}{g_N}. \quad (10)$$

D is where $g_{\text{ratio}} \geq a_{\text{ratio}}$.

*sjhalayka@gmail.com




Figure 1: Where $D = 3$, as viewed from the side. The field lines are isotropic, spherical.




Figure 2: Where $D = 2.1$, as viewed from the side. The field lines are increasingly anisotropic.



Figure 3: Where $D = 2.001$, as viewed from the side. The field lines are anisotropic, disk-like.




Figure 4: $R = 100$, $r = 1$, $n = 10^8$, $\epsilon = 1$. The analytical plot is generated by the formula $y = n/(2R^D)$.

References

- [1] Misner et al. Gravitation. (1970)
- [2] 't Hooft. Dimensional reduction in quantum gravity. (1993)
- [3] Susskind. The World as a Hologram. (1994)




Figure 5: Visualization of the relativistic Doppler effect for 10^6 stars. The Newtonian orbit is on the left, and the flat rotation curve orbit is on the right. The redshift of the wavelength indicates stars moving away from the camera, and blueshift of the wavelength indicates stars moving toward the camera. Thus, the stars in the galaxy are orbiting counterclockwise. On the left, the wavelength is dependent on angle and distance from the galactic centre. On the right, the wavelength is dependent only on angle, which means that there is a constant orbit speed that is independent of the distance from the galactic centre. This is exactly what Vera Rubin discovered in the disks of the galaxies that she observed, and so dark matter was posited.