

Can HIV epidemics among men who have sex with men be eliminated through participation to PrEP rollouts?

Sofia JIJÓN¹, Jean-Michel MOLINA², Dominique COSTAGLIOLA¹, Virginie SUPERVIE¹ and Romulus BREBAN^{3,§}

1 Sorbonne Université, INSERM, Institut Pierre Louis d'épidémiologie et de Santé Publique (IPLESP UMR-S 1136), 75012 Paris, France

2 Département de Maladies Infectieuses, APHP-Hôpital Saint Louis, UMR 941 Inserm et Sorbonne Paris Cité, Paris, France

3 Institut Pasteur, Unité d'Épidémiologie des Maladies Émergentes, 75015 Paris, France

§Corresponding author: Romulus Breban

Institut Pasteur, Unité d'Épidémiologie des Maladies Émergentes

25 rue du Docteur Roux

75015 Paris, France

+33 (0)1 40 61 39 65

romulus.breban@pasteur.fr

E-mail addresses of authors:

SJ: sofia.jijon@iplesp.upmc.fr

J-MM: jean-michel.molina@aphp.fr

DC: dominique.costagliola@iplesp.upmc.fr

VS: virginie.supervie@inserm.fr

RB: romulus.breban@pasteur.fr

Word count:

Abstract: 243/250

Main text: 3471/3500

Abstract

Objectives: To study the conditions under which PrEP coverage can eliminate HIV among men who have sex with men (MSM) in the Paris region.

Design: Mathematical modeling.

Methods: We propose an innovative approach, combining a transmission model with a game-theoretic model, for decision-making about PrEP use. Individuals at high risk of HIV infection decide to use PrEP, depending on their perceived risk of infection and the relative cost of using PrEP versus antiretroviral treatment (ART), which includes monetary and/or non-monetary aspects, such as price and access model of PrEP, consequences of being infected and lifelong ART.

Results: If individuals perceive fairly their infection risk, and the cost of using PrEP is sufficiently low, the PrEP coverage can lead to elimination. Specifically, assuming 86% PrEP effectiveness, as observed in two clinical trials, a minimum PrEP coverage of 55% (95% CI:43%–64%) among high-risk MSM would achieve elimination in the Paris region. A complete condom drop by PrEP users slightly increases the minimum PrEP coverage by ~1%, while underestimation of their own HIV infection risk would demand PrEP programs to reduce the cost of using PrEP by a factor ~2 to achieve elimination.

Conclusions: Elimination conditions are not yet met in the Paris region, where at most 47% of high-risk MSM were using PrEP as of mid-2019. Further lowering the cost of

24 PrEP and promoting a fair perception of HIV risk are required and should be maintained
25 in the long run, to maintain elimination status.

26

27

28 **Keywords:** Pre-exposure prophylaxis; HIV; men who have sex with men; behavioral
29 epidemiology; game theory; prevention coverage.

30 Introduction

31 In many settings, men who have sex with men (MSM) are most affected by HIV [1]. Pre-
32 exposure prophylaxis (PrEP) is a highly effective prevention method recommended by
33 the WHO for individuals at high risk of infection with HIV [2]. Both IPERGAY and
34 PROUD clinical trials showed that PrEP can reduce HIV incidence among MSM by 86%
35 [3,4]. Modeling studies, elaborating on these results, suggested that PrEP has the potential
36 to curtail, and even eliminate HIV epidemics, notably among MSM [5–8]. For instance,
37 in the Netherlands, elimination would require 82% PrEP coverage in the highest-risk
38 group [6].

39
40 The question of whether it is possible to achieve a certain PrEP coverage in a population
41 has not been addressed; modeling studies only assume that the coverage reaches certain
42 values, which may not be granted in public health practice. It is therefore unclear whether,
43 and under what conditions, target PrEP coverage levels, required to eliminate HIV
44 epidemics, can be reached voluntarily and maintained in the long run. Currently, PrEP
45 remains underutilized in many settings [12]. For instance, in the United States, 220,000–
46 225,000 individuals were on PrEP as of April 2020 [9], still short of the CDC estimate
47 that 1.2 million persons have indications for considering PrEP use [10]. Furthermore, a
48 recent study shows that only two in five individuals keep using PrEP for >2 years [11].

49
50 Mathematical tools for modeling individual-level decision-making are offered by game
51 theory [13–15]. We propose an innovative approach, combining an epidemic model at
52 the population level, and a game-theoretic model for decision-making about PrEP at the
53 individual level. We model PrEP adoption in a population at high risk of HIV infection

54 to determine whether and under what conditions certain PrEP coverage levels can be
55 reached voluntarily. Particularly, we study the potential impact of PrEP among MSM in
56 the Paris region of France, where universal antiretroviral treatment (ART) is in place, and
57 PrEP is available for eligible individuals.

Methods

We built an HIV epidemic model (Figures S1 and S2) to describe the epidemiological context of an MSM community where eligible individuals make informed decisions about adopting PrEP. The decision making is modeled as a non-cooperative game, where individuals act to maximize the utility of adopting PrEP, or, in other words, minimize the cost of using PrEP to avoid acquiring HIV and lifelong ART. Individual's decision is, however, indirectly influenced by that of others. The sum of all individuals' decisions determines the PrEP coverage, which, consequently, affects epidemic progression and the risk of acquiring HIV. The decision-making game model is thus intertwined with the epidemic model. Below, we describe the main features of our two-component model; see the Supplementary Material (SM) for further details.

The epidemic model

The epidemic model stratifies the MSM population into two risk groups (low and high), to account for heterogeneity in the infection risk. The majority of partnerships occur within the same risk group and individuals at high risk of infection drive the epidemic. The model also stratifies over HIV status, disease progression, diagnosis and the use of PrEP or ART. Once diagnosed, individuals immediately begin ART, no longer transmitting HIV. We varied PrEP effectiveness, denoted ε , from 0 to 100% to study sub-optimal PrEP adherence. The PrEP coverage, p , was not fixed; rather, it was obtained through the decision-making game model (see below).

We computed the effective reproduction number for the epidemic model, R , defined as the expected number of secondary cases caused by one infected individual, during his

entire infectious period, in an uninfected population subject to control interventions [20,21]. PrEP use may change individuals' preference for other prevention tools, turning R into a function of PrEP parameters. $R(p, \varepsilon) > 1$ indicates epidemic persistence, meaning that an endemic state will be reached. Elimination requires $R(p, \varepsilon) < 1$, such that the disease-free state will be reached. Elimination implies that incidence is reduced to zero in the studied population, but HIV can re-emerge in absence of control interventions, as it does not imply eradication. We say the epidemic is controlled using PrEP if $R(p, \varepsilon)$ decreases with the PrEP parameters, although the decrease is not below 1. Our model shows that epidemic control and elimination can occur through PrEP, provided that two thresholds in PrEP effectiveness are exceeded; $\varepsilon \geq \varepsilon_C$ is required for epidemic control and $\varepsilon \geq \varepsilon_E$ for epidemic elimination (see SM section 1.2.3). These thresholds are called the epidemic control and the epidemic elimination thresholds, respectively.

The decision-making game model

During an epidemic, individuals may adopt PrEP according to their perceived HIV risk [18], price [19] and access model of PrEP [20], adverse effects [21], social stigma [22], consequences of being infected, lifelong ART, and other pros and cons. These factors, summarizing monetary and/or non-monetary aspects, are expressed in our decision-making model as costs perceived by the individual.

We assume that individuals choose between two mutually-exclusive strategies. If an individual decides not to use PrEP, then in the case of acquiring HIV he will start ART upon positive HIV diagnosis, and pay the cost of ART for the rest of his life; we use the

notation $C_{\text{No-PrEP}}$ for the lifetime cost of this strategy. Otherwise, the individual decides to adopt PrEP prevention. Thus, he takes and pays the cost of PrEP and, in the case of acquiring HIV despite PrEP uptake, being diagnosed and starting ART, pays the cost of ART for the rest of his life. We use the notation C_{PrEP} for the lifetime cost of the second strategy. The total cost explicitly depends on the yearly costs of ART and PrEP, the PrEP parameters, and, implicitly, the yearly risk of acquiring HIV. Introducing r , the relative cost of PrEP versus ART, the balance of cost, when the probability to adopt PrEP is p , becomes

$$C(p, \varepsilon, r) = pC_{\text{PrEP}}(p, \varepsilon, r) + (1 - p)C_{\text{No-PrEP}}(p, \varepsilon),$$

where all functions and parameters, other than p , are given in our mathematical modeling. The value of p that minimizes $C(p, \varepsilon, r)$, denoted $\hat{p}(\varepsilon, r)$, estimates the probability that a typical high-risk individual adopts PrEP, and also represents the voluntary PrEP coverage among high-risk MSM. The solution of the game represents a endemic state where individuals make decisions to adopt PrEP in stationary epidemiological context. We thus assumed that, for the long run, individuals stand by their decisions about adopting PrEP.

Application to the HIV epidemic among MSM in the Paris region

We calibrated the epidemic model to represent the epidemiological context before the introduction of PrEP [23,24], and obtained many HIV parameter sets, to reveal uncertainty in the model output (SM section 2 and Tables S1-S4). In our baseline scenario, we assumed that MSM on PrEP get tested for HIV quarterly, according to the French recommendations [25]. The testing frequency on PrEP was thus much higher than that observed off PrEP, as data shows 3.1 years for the median time from HIV infection

130 to diagnosis among MSM before the introduction of PrEP (personal communication with
131 VS). We further assumed that individuals have a fair perception of their infection risk
132 when making decisions about PrEP use; the infection risk was determined by the force of
133 HIV infection of the epidemic model. Furthermore, MSM were assumed to drop condom
134 use from 30% to 20% when adopting PrEP [3], and the condom effectiveness was 58%–
135 80% [26]. Sensitivity scenarios were explored assuming that i) MSM misperceived their
136 risk of acquiring HIV, ii) MSM adopting PrEP completely dropped condom use [27], or
137 iii) MSM did not change their HIV testing behavior upon adopting PrEP (SM Section 3).

Results

About 500 parameter sets calibrated our epidemic model to the HIV epidemiology among MSM in the Paris region, before the introduction of PrEP: total yearly mean incidence was 1.3%, prevalence was 17%, and 17% of MSM living with HIV were undiagnosed (Table S3). The mean number of MSM was $\sim 111,000$, of which 13% (i.e., $\sim 14,200$) were at high risk of infection and eligible for PrEP. Yearly incidence for high-risk MSM was 7%. The model parameters implied that the PrEP rollout had two effects: first, it offered the prevention benefits of the regimen, and, second, it behaved as a test-and-treat strategy [28,29], imposing a major change in HIV testing practice (SM Section 3.1 and Figure S3).

The voluntary PrEP coverage if individuals perceived correctly HIV infection risk

We first investigated a typical parameter set calibrating our model; Table S2. The PrEP coverage starts at zero, before introducing PrEP, and then reaches an equilibrium value where the expected cost of adopting PrEP is minimum. The final value reached depends on HIV parameters of the epidemic before the introduction of PrEP, the PrEP effectiveness, ϵ , and the perceived relative cost of PrEP versus ART, r . Figure 1A shows the voluntary PrEP coverage reached among high-risk MSM, $\hat{p}(\epsilon, r)$. Figure 1B shows the corresponding relative reduction in HIV incidence in the MSM community. Each of these two figures shows three regions:

- Region III, where no high-risk MSM adopts PrEP, because the perceived relative cost of PrEP versus ART is too high. Therefore, HIV remains endemic, unaffected by the introduction of PrEP (i.e., no reduction in incidence);

- Region II, where some, but not enough, high-risk MSM adopt PrEP, since the relative cost remains high. The epidemic is controlled and incidence decreases, but not enough for elimination (i. e., $R(\hat{p}, \varepsilon) > 1$);
- Region I, where PrEP is offered at low relative cost. This allows reaching high levels of PrEP coverage (~54–75%) and the epidemic can be eliminated; for Region I, $R(\hat{p}, \varepsilon) < 1$. HIV elimination for low PrEP effectiveness (bottom part of Figure 1A) occurs as a consequence of the test-and-treat effect of the PrEP rollout; consequently, $\varepsilon_C = \varepsilon_E = 0\%$. In this case, MSM taking PrEP are poorly protected against HIV. However, they are diagnosed and treated very early in the course of infection, because they get tested for HIV every three months. Early diagnosis and treatment prevent further HIV transmission. In contrast, when PrEP effectiveness is high (top part of Figure 1A), most on-PrEP MSM do not acquire HIV, so the test-and-treat benefit of the PrEP rollout is marginal. It is PrEP, particularly its high effectiveness, that contributes decisively to epidemic elimination. If $\varepsilon = 86\%$, as observed in the IPERGAY and PROUD trials, a minimum PrEP coverage of 56% should be reached among high-risk MSM, to eliminate HIV; Figure S4.

It is important to note that elimination is temporary, as the disease-free state is unstable.

Indeed, once the epidemic is eliminated, individuals perceive HIV risk as being low and may reevaluate PrEP-induced advantages and disadvantages; this may severely increase the relative cost of PrEP versus ART, since the epidemic is considered to be eliminated.

As fewer individuals consider PrEP use, the PrEP coverage decreases and the HIV

epidemic dynamics in Region I can enter Region II, where the epidemic reemerges and becomes again of public health concern.

We generated the outputs in Figure 1 using each of the ~ 500 parameter sets obtained through calibration, to estimate uncertainty intervals for our results (SM Section 2). Figure 2A shows the probability that HIV is eliminated, as a function of ε and r . The probability is high on the left, where Region I is found, and declines severely toward Region II. In Figure 2B, we illustrate the boundaries between Regions I and II (continuous line), and between Regions II and III (dashed line); the three-region structure appears robust to parameter uncertainties. Additionally, when $\varepsilon = 86\%$, we found that the 95% confidence interval (CI) of the minimum PrEP coverage needed for elimination (i.e., 55%) is 43%–64%.

Sensitivity scenarios

We assumed that individuals could misperceive their HIV risk when deciding to adopt PrEP, and repeated our analyses. Specifically, rather than having a fair sense of HIV risk, based on the force of infection, high-risk MSM could get a sense of HIV risk from, for instance, the proportion of their high-risk MSM peers being diagnosed each year with HIV (SM Section 3.3.1). The voluntary PrEP coverage computed for this scenario is illustrated in Figure 3A and reveals a qualitatively similar structure to that in Figure 1. However, when high-risk MSM misinterpret and underestimate their HIV risk, Region I is smaller, implying that the relative cost of PrEP versus ART must be lower to achieve epidemic elimination. In particular, when $\varepsilon = 86\%$, the relative cost needed for epidemic

elimination decreases by a factor of ~ 2 , making Region I harder to reach in practice of public health.

We performed two other sensitivity analyses. First, we analyzed PrEP-driven condom drop. In our baseline scenario, MSM dropped condom use from 30% to 20% when adopting PrEP. Similar results were obtained assuming that PrEP users stopped using condoms completely (SM Section 3.3.2 and Figure S5). We thus concluded that condom drop is not a major factor against HIV elimination when PrEP effectiveness is high. Specifically, epidemic elimination where $\varepsilon = 86\%$ requires a coverage of $>57\%$, rather than $>56\%$ in the baseline scenario.

Second, we analyzed a scenario where PrEP uptake does not require higher rates of HIV testing and MSM do not change their HIV testing behavior when adopting PrEP; Figure 3B and SM Section 3.3.3. In this case, HIV elimination is entirely due to PrEP and can only be reached if $\varepsilon \geq \varepsilon_E = 58\%$. For high PrEP effectiveness, the results are very close to those of the baseline scenario, as very few MSM fail PrEP and the testing frequency on PrEP does not impact many MSM. For instance, epidemic elimination when $\varepsilon = 86\%$ requires a coverage of $>58\%$, rather than $>56\%$ in the baseline scenario. For low PrEP effectiveness, Figure 3B shows a fourth region, where low relative cost encourages all high-risk MSM to adopt PrEP ($\hat{p} = 100\%$) and PrEP effectiveness is above the epidemic control threshold ($\varepsilon_C = 8\%$), but below the epidemic elimination threshold ($\varepsilon_E = 58\%$). Therefore, the epidemic is controlled, but not eliminated, and a new HIV endemic state is reached; $R(\hat{p}, \varepsilon) > 1$.

Perspectives on the PrEP rollout in the Paris region

In 2016, a PrEP rollout started in the Paris region, offering fully subsidized PrEP to eligible individuals. As aforementioned, under the baseline scenario, for 86% PrEP effectiveness, we found that at least 55% (95%CI: 43%–64%) of the high-risk MSM would need to take PrEP for the HIV epidemic be eliminated. Since, according to our calibration, the estimated number of PrEP-eligible MSM in the Paris region is 14,200 (95%CI: 9,200–23,000), this means that 7,700 (95%CI: 5,800–10,100) high-risk MSM should be on PrEP. This is an objective to be reached. As of mid-2019, ~6,700 men were on PrEP in the Paris region [30], with a marked growing trend; the 30-month dropout rate was ~32% [31]. The PrEP coverage among high-risk MSM was then estimated to be at most 47% (95%CI: 30%–73%), assuming that all men on PrEP were indeed high-risk MSM. If all these MSM maintained using PrEP in the long term, our model predicted epidemic control (i.e., Region II), with a reduction of 90% (95%CI: 81%–100%) in HIV incidence at the new endemic state.

Discussion

We addressed the role of individual-level decision-making in the potential impact of PrEP on the HIV epidemic, identified the conditions for epidemic control or elimination, and estimated PrEP coverage levels which can be reached voluntarily. We obtained four major findings for PrEP rollouts. First, HIV epidemics can be eliminated provided that the relative cost of using PrEP versus ART is sufficiently low. Second, frequent HIV testing while taking PrEP can compensate for poor PrEP adherence and act as a test-and-treat intervention. Third, HIV risk perception may play a major role for elimination, while drop in condom use among PrEP users may not. Fourth, epidemic elimination may be only temporary.

We applied our model to the Paris region. Assuming a PrEP effectiveness of 86%, as reported in two major clinical trials, we found that at least 55% (95%CI: 43%–64%) of the high-risk MSM would need to be on PrEP to achieve HIV elimination. As of mid-2019, at most 47% high-risk MSM were on PrEP in the Paris region, meaning that the PrEP-rollout protocol did not reduce enough the cost of PrEP for epidemic elimination, so far. Still, a recent update on new HIV diagnoses in Paris [32] shows that the numbers among French-born MSM decreased by 28%, between 2015 and 2018, with no significant decrease for other MSM. This decrease could be partly due to the PrEP rollout starting in 2016, and, according to our modeling, should continue in the near future. In two other settings, a moderate-high PrEP coverage has been quickly reached. The region of New South Wales witnessed a rapid PrEP rollout (~9,000 MSM on PrEP within 2 years) during an implementation study providing PrEP for free at several sites, including public HIV and sexual health services, and private general practices with expertise in ART

prescription [33]. About 41% of the high-risk MSM in Australia were on PrEP in 2017 [34]. Since April 2018, PrEP is subsidized by the Australian government and can be prescribed by any practitioner [35]. In San Francisco, a citywide-coordinated PrEP rollout, within the Getting to Zero program, strongly promoted PrEP use and allowed many people to access PrEP for free or at low monetary cost, through insurance benefits or patient assistance programs. Close to 50% of the eligible MSM were on PrEP in 2017 in San Francisco [36]. Although these levels of PrEP coverage contributed to decreasing HIV transmission [33,36,37], HIV elimination has not been reported.

Moving toward epidemic elimination will require further decreasing the cost of PrEP, which may involve reducing monetary and non-monetary barriers to PrEP uptake, such as difficulties in accessing PrEP, pill burden, tolerability of the molecules, social stigma and discrimination, and the acquisition of other sexually transmitted infections in case of dropping condom use [20–22]. Online tools [38], home-based programs [39], long-lasting injectable versions of PrEP [40], rather than daily or on-demand pills, allowing trained general practitioners to prescribe PrEP and interventions that increase awareness, motivations and behavioral skills about risk reduction [41] may also help reduce the perceived cost of PrEP and decrease the drop-out rate. It is important to note that, in practice, estimating the cost of PrEP relative to that of ART can be complex, but not strictly needed, as reducing this cost may be intuitive and places the PrEP rollout in the right direction. Instead, the PrEP coverage and HIV incidence can serve as indicators for how far the PrEP rollout is from achieving elimination.

Moving toward epidemic elimination will also require reaching MSM who may not perceive themselves at high risk, and thus require a lower cost for adopting PrEP, in order to join the prevention effort. Recent studies found that high-risk individuals can underestimate their HIV risk [42] and there are many missed opportunities for PrEP uptake [43]. Specifically, in France, >90% of the recently infected individuals were eligible for PrEP [43]. Therefore, assessing and communicating individual-level risk for acquiring HIV remains a key objective for achieving elimination. Promoting a fair perception of HIV risk can be achieved through, not only advertising and marketing PrEP [44], but also through using electronic health records for identifying high-risk MSM [45].

Importantly, if HIV is eliminated, interventions will be needed so individuals keep perceiving a low cost for PrEP and fair perception of HIV risk, to maintain a high PrEP coverage. Otherwise, HIV can reemerge and reach again an endemic state of concern for public health. The situation is similar to that of vaccination prevention, which requires continuous vaccine coverage even though the disease is nearly eliminated [46].

Our study has some limitations. First, we assumed that individuals act out of self-interest and do not cooperate to avoid getting infected by HIV. Modeling PrEP adoption through other theories of health behavior, considering for instance interactions between individuals [15], remains a subject to be studied in further work. Second, we assumed that MSM are homogeneous regarding risk perception. In reality, the MSM population is certainly heterogeneous, fair perception co-existing with misperception. Nevertheless, our baseline and alternative scenarios can be regarded as optimistic and pessimistic scenarios, respectively. Third, we did not account for migration or travel [47], nor for condom drop

among non-PrEP users [48], which could influence elimination efforts. Fourth, our estimates of the number of high-risk individuals, who should be on PrEP for HIV elimination, depend on the size of the MSM community, which is a metric difficult to estimate. Also, the number of high-risk MSM on PrEP currently reported, and hence the PrEP coverage, may represent an overestimate because establishing PrEP eligibility relies on self-reported behavior, which is difficult to appraise by practitioners.

Conclusion

Perception of the cost of PrEP and of HIV risk are two important levers to increase voluntary use of PrEP, reach coverage levels necessary to eliminate HIV, and maintain elimination in the context of less epidemic adversity. Current PrEP rollouts should aim at lowering the perceived cost of using PrEP and promoting a fair perception of the risk of acquiring HIV, to realize the full potential of PrEP prevention.

Authors' contributions

SJ, VS and RB conceived the model. SJ conducted the numerical simulations. All authors participated to the writing of the manuscript, analysis and interpretation of the results. All authors read and approved the final manuscript.

Acknowledgements

SJ was supported by an ANRS fellowship and a PhD fellowship from the French Ministry of Higher Education and Research, obtained via the Public Health Doctoral Network coordinated by the EHESP. The sponsors had no role in the study.

List of abbreviations

ART: Antiretroviral treatment; MSM: Men who have sex with men; PrEP: Pre-exposure prophylaxis; CI: confidence interval.

345 **References**

- 346 1. Beyrer C, Baral SD, Collins C, et al. The global response to HIV in men who
347 have sex with men. *Lancet* (London, England) **2016**; 388:198–206.
- 348 2. World Health Organization. Policy brief: WHO expands recommendation on oral
349 pre-exposure prophylaxis of HIV infection (PrEP). World Heal. Organ. 2015.
350 Available at: <https://apps.who.int/iris/handle/10665/197906>. [Website accessed
351 on June 20, 2020].
- 352 3. Molina J-M, Capitant C, Spire B, et al. On-Demand Preexposure Prophylaxis in
353 Men at High Risk for HIV-1 Infection. *N Engl J Med* **2015**; 373:2237–2246.
- 354 4. McCormack S, Dunn DT, Desai M, et al. Pre-exposure prophylaxis to prevent the
355 acquisition of HIV-1 infection (PROUD): Effectiveness results from the pilot
356 phase of a pragmatic open-label randomised trial. *Lancet* **2016**; 387:53–60.
- 357 5. Jenness SM, Goodreau SM, Rosenberg E, et al. Impact of the Centers for Disease
358 Control’s HIV Preexposure Prophylaxis Guidelines for Men Who Have Sex With
359 Men in the United States. *J Infect Dis* **2016**; 214 :1800–1807..
- 360 6. Rozhnova G, Heijne J, Bezemer D, et al. Elimination prospects of the Dutch HIV
361 epidemic among men who have sex with men in the era of preexposure
362 prophylaxis. *Aids* **2018**; 32:2615–2623.
- 363 7. Scott N, Stoové M, Kelly SL, Wilson DP, Hellard ME. Achieving 90-90-90
364 Human Immunodeficiency Virus (HIV) Targets Will Not Be Enough to Achieve
365 the HIV Incidence Reduction Target in Australia. *Clin Infect Dis* **2018**; 66:1019–
366 1023.
- 367 8. Hansson D, Stromdahl S, Leung KY, Britton T. Introducing pre-exposure
368 prophylaxis to prevent HIV acquisition among men who have sex with men in

- Sweden: insights from a mathematical pair formation model. *BMJ Open* **2020**; 10:e033852.
9. PrEP Watch. Global PrEP Tracker. 2010. Available at: <https://www.prepwatch.org/resource/global-prep-tracker/>. [Website accessed on June 20, 2020].
10. Smith DK, Van Handel M, Wolitski RJ, et al. Vital signs: Estimated percentages and numbers of adults with indications for preexposure prophylaxis to prevent HIV acquisition - United States, 2015. *Morb Mortal Wkly Rep* **2015**; 64:1291–1295.
11. Coy KC, Hazen RJ, Kirkham HS, Delpino A, Siegler AJ. Persistence on HIV preexposure prophylaxis medication over a 2-year period among a national sample of 7148 PrEP users, United States, 2015 to 2017. *J Int AIDS Soc* **2019**; 22:e25252.
12. Cohen J. Concern as HIV prevention strategy languishes. *Science* (80-) **2018**; 359:1205–1205.
13. Verelst F, Willem L, Beutels P. Behavioural change models for infectious disease transmission: a systematic review (2010–2015). *J R Soc Interface* **2016**; 13:20160820.
14. Manfredi P, D’Onofrio A, editors. *Modeling the Interplay Between Human Behavior and the Spread of Infectious Diseases*. New York, NY: Springer New York, 2013.
15. Chang SL, Piraveenan M, Pattison P, Prokopenko M. Game theoretic modelling of infectious disease dynamics and intervention methods: a review. *J Biol Dyn* **2020**; 14:57–89.

- 393 16. Jacquez JA, Simon CP, Koopman J, Sattenspiel L, Perry T. Modeling and
 394 analyzing HIV transmission: the effect of contact patterns. *Math Biosci* **1988**;
 395 92:119–199.
- 396 17. van den Driessche P, Watmough J. Reproduction numbers and sub-threshold
 397 endemic equilibria for compartmental models of disease transmission. *Math*
 398 *Biosci* **2002**; 180:29–48.
- 399 18. Bull L, Dimitrijevic P, Beverley S, et al. Perceived need of, and interest in, HIV
 400 pre-exposure prophylaxis amongst men who have sex with men attending three
 401 sexual health clinics in London, UK. *Int J STD AIDS* **2018**; 29:435–442.
- 402 19. van Dijk M, de Wit JBF, Guadamuz TE, Martinez JE, Jonas KJ. Slow uptake of
 403 PrEP: Behavioral predictors and the influence of price on PrEP uptake among
 404 MSM with a high interest in PrEP. *OSF Preprints* **2020**;
- 405 20. Gilson RI, Clutterbuck DJ, Chen ZE. Demand for pre-exposure prophylaxis for
 406 HIV and the impact on clinical services: Scottish men who have sex with men
 407 perspectives. *Int J STD AIDS* **2018**; 29:273–277.
- 408 21. Thomann M, Grosso A, Zapata R, Chiasson MA. ‘WTF is PrEP?’: attitudes
 409 towards pre-exposure prophylaxis among men who have sex with men and
 410 transgender women in New York City. *Cult Health Sex* **2017**; 20:772–786.
- 411 22. Brooks RA, Nieto O, Landrian A, Donohoe TJ. Persistent stigmatizing and
 412 negative perceptions of pre-exposure prophylaxis (PrEP) users: implications for
 413 PrEP adoption among Latino men who have sex with men. *AIDS Care* **2019**;
 414 31:427–435.
- 415 23. Marty L, Cazein F, Panjo H, Pillonel J, Costagliola D, Supervie V. Revealing
 416 geographical and population heterogeneity in HIV incidence, undiagnosed HIV

- 417 prevalence and time to diagnosis to improve prevention and care: estimates for
 418 France. *J Int AIDS Soc* **2018**; 21:e25100.
- 419 24. Santé Publique France. Surveillance de l'infection à VIH (dépistage et
 420 déclaration obligatoire), 2010-2017. Available at:
 421 [https://www.santepubliquefrance.fr/maladies-et-traumatismes/infections-](https://www.santepubliquefrance.fr/maladies-et-traumatismes/infections-sexuellement-transmissibles/vih-sida/documents/bulletin-national/bulletin-de-sante-publique-infection-a-vih.-mars-2019)
 422 [sexuellement-transmissibles/vih-sida/documents/bulletin-national/bulletin-de-](https://www.santepubliquefrance.fr/maladies-et-traumatismes/infections-sexuellement-transmissibles/vih-sida/documents/bulletin-national/bulletin-de-sante-publique-infection-a-vih.-mars-2019)
 423 [sante-publique-infection-a-vih.-mars-2019](https://www.santepubliquefrance.fr/maladies-et-traumatismes/infections-sexuellement-transmissibles/vih-sida/documents/bulletin-national/bulletin-de-sante-publique-infection-a-vih.-mars-2019) [Website accessed on June 20, 2020].
- 424 25. CNS, ANRS. Prise en charge médicale des personnes vivant avec le VIH. 2018;
 425 Available at: [https://cns.sante.fr/actualites/prise-en-charge-du-vih-](https://cns.sante.fr/actualites/prise-en-charge-du-vih-recommandations-du-groupe-dexperts/)
 426 [recommandations-du-groupe-dexperts/](https://cns.sante.fr/actualites/prise-en-charge-du-vih-recommandations-du-groupe-dexperts/) [Website accessed on June 20, 2020].
- 427 26. Smith DK, Herbst JH, Zhang X, Rose CE. Condom Effectiveness for HIV
 428 Prevention by Consistency of Use Among Men Who Have Sex With Men in the
 429 United States. *J Acquir Immune Defic Syndr* **2015**; 68:337–344.
- 430 27. Holt M, Lea T, Mao L, et al. Community-level changes in condom use and
 431 uptake of HIV pre-exposure prophylaxis by gay and bisexual men in Melbourne
 432 and Sydney, Australia: results of repeated behavioural surveillance in 2013-17.
 433 *Lancet HIV* **2018**; 3018:1–9.
- 434 28. Kretzschmar ME, van der Loeff MF, Birrell PJ, De Angelis D, Coutinho RA.
 435 Prospects of elimination of HIV with test-and-treat strategy. *Proc Natl Acad Sci*
 436 *U S A* **2013**; 110:15538–15543.
- 437 29. WHO. Prevent HIV, test and treat all. 2016; Available at: [https://www.who.int](https://www.who.int/hiv/pub/progressreports/2016-progress-report/en/)
 438 [/hiv/pub/progressreports/2016-progress-report/en/](https://www.who.int/hiv/pub/progressreports/2016-progress-report/en/) [Website accessed on June 20,
 439 2020].
- 440 30. Gage SB De, Tri TLE, Spira RD. Suivi de l'utilisation de Truvada ou génériques

pour une prophylaxie pré-exposition (PrEP) au VIH à partir des données du
Système National des Données de Santé (SNDS). Actualisation des
données jusqu'au 30 juin 2019. :1–37. Available at:
https://www.ansm.sante.fr/var/ansm_site/storage/original/application/0511b009d265687a2ff8d2fa266085ce.pdf. [Website accessed on June 20, 2020].

31. Costagliola D, Ghosn J, Spire B, et al. PrEP persistence and associated factors: an analysis from the ANRS Prevenir study. *HIV Med* **2019**; 20:33.
32. Santé Publique France. Dépistage du VIH et découvertes de séropositivité VIH à Paris, données 2018. Available at: <https://www.santepubliquefrance.fr/les-actualites/2019/depistage-du-vih-et-decouvertes-de-seropositivite-vih-a-paris-donnees-2018> [Website accessed on June 20, 2020].
33. Grulich AE, Guy R, Amin J, et al. Population-level effectiveness of rapid, targeted, high-coverage roll-out of HIV pre-exposure prophylaxis in men who have sex with men: the EPIC-NSW prospective cohort study. *Lancet HIV* **2018**; 3018:1–9.
34. Kirby Institute. HIV diagnoses in Australia drop to lowest number in 18 years. Available at: <https://kirby.unsw.edu.au/news/hiv-diagnoses-australia-drop-lowest-number-18-years> [Website accessed on June 20, 2020].
35. Australasian Society for HIV VH and SHM (ASHM). HIV PrEP available on PBS in Australia from 1 April. 2018; Available at <https://www.ashm.org.au/news/hiv-PrEP-available-announced-on-pbs/> [Website accessed on June 20, 2020].
36. San Francisco Department of Public Health. HIV Epidemiology Annual Report 2017. 2017. Available at:

- 465 <https://www.sfdph.org/dph/comupg/oprograms/HIVepiSec/HIVepiSecReports.as>
 466 p.
- 467 37. Smith DK, Sullivan PS, Cadwell B, et al. Evidence of an Association of
 468 Increases in Pre-exposure Prophylaxis Coverage With Decreases in Human
 469 Immunodeficiency Virus Diagnosis Rates in the United States, 2012--2016. *Clin*
 470 *Infect Dis* **2020**; Online ahead of print.
- 471 38. Getting to zero San Francisco. PrEP Committee. Available at:
 472 <http://www.gettingtozerosf.org/prep-committee/> [Website accessed on June 20,
 473 2020].
- 474 39. Siegler AJ, Mayer KH, Liu AY, et al. Developing and Assessing the Feasibility
 475 of a Home-based Preexposure Prophylaxis Monitoring and Support Program.
 476 *Clin Infect Dis* **2019**; 68:501–504.
- 477 40. Marshall BDL, Goedel WC, King MRF, et al. Potential effectiveness of long-
 478 acting injectable pre-exposure prophylaxis for HIV prevention in men who have
 479 sex with men: a modelling study. *Lancet HIV* **2018**; 3018:1–8.
- 480 41. Dubov A, Altice FL, Fraenkel L. An Information–Motivation–Behavioral Skills
 481 Model of PrEP Uptake. *AIDS Behav* **2018**; 22:3603–3616.
- 482 42. Blumenthal J, Jain S, Mulvihill E, et al. Perceived Versus Calculated HIV Risk. *J*
 483 *Acquir Immune Defic Syndr* **2019**; 80:e23–e29.
- 484 43. Lions C, Cabras O, Cotte L, et al. Missed opportunities of HIV pre-exposure
 485 prophylaxis in France: a retrospective analysis in the French DAT’AIDS cohort.
 486 *BMC Infect Dis* **2019**; 19:278.
- 487 44. Amico KR, Bekker L. Global PrEP roll-out : recommendations for programmatic
 488 success. *Lancet HIV* **2019**; 6:e137–e140.

- 489 45. Marcus JL, Hurley LB, Krakower DS, Alexeeff S, Silverberg MJ, Volk JE. Use
490 of electronic health record data and machine learning to identify candidates for
491 HIV pre-exposure prophylaxis: a modelling study. *Lancet HIV* **2019**; 3018:1–8.
- 492 46. Jijón S, Supervie V, Breban R. Prevention of treatable infectious diseases: A
493 game-theoretic approach. *Vaccine* **2017**; 35:5339–5345.
- 494 47. Palk L, Gerstoft J, Obel N, Blower S. A modeling study of the Danish HIV
495 epidemic in men who have sex with men: travel, pre-exposure prophylaxis and
496 elimination. *Sci Rep* **2018**; 8:16003.
- 497 48. Phanuphak N, Phanuphak P. Time to focus more on condomless anal sex in non-
498 PrEP users. *Lancet HIV* **2018**; 3018:17–18.
- 499
- 500

Figure legends

Figure 1. The voluntary PrEP coverage and its impact on HIV incidence, assuming fair risk perception


Color maps of **(A)** the voluntary PrEP coverage among high-risk men who have sex with men (MSM), \hat{p} , and **(B)** the corresponding reduction in the overall endemic HIV incidence rate, as functions of ε and r , assuming that individuals have a fair perception of HIV risk. The model outputs were obtained for one typical parameter set calibrating our model. Three regions were identified, depending on \hat{p} : Region III, where r is high and no MSM uses PrEP ($\hat{p} = 0\%$), so HIV incidence is not reduced; Region II, where some, but not enough MSM use PrEP, since r remains high, and thus the epidemic is controlled; and Region I (marked by blue stripes), where epidemic elimination is possible.

Figure 2. The probability of HIV elimination and boundary uncertainty for the three-region structure

(A) The probability of HIV epidemic elimination due to voluntary PrEP coverage, obtained from the ~500 calibrated parameter sets. **(B)** The boundaries (the mean is represented as a line and the 95% confidence interval as grey area) between Regions I and II (continuous line), and between Regions II and III (dashed line).


Figure 3. Sensitivity analyses for the baseline scenario

(A) Decision-making based on misperceived risk of acquiring HIV can significantly reduce the size of Region I, where epidemic elimination is possible (blue stripes), despite high levels of PrEP effectiveness. **(B)** By assuming fair perception of HIV risk but no change in HIV testing behavior among PrEP users (i.e., the time from infection to diagnosis remains at 3.1 years), the PrEP effectiveness thresholds required to reach epidemic control and epidemic elimination become $\varepsilon_C = 8\%$ and $\varepsilon_E = 58\%$, respectively. In turn, this yields a fourth region, denoted Region IV, where all individuals adopt PrEP but $\varepsilon_C \leq \varepsilon < \varepsilon_E$, so the epidemic is controlled, yet not eliminated. Regions II and III in both panels depict some and no PrEP adoption, respectively, similarly to Figure 1A.


537 **Figures**538 **Figure 1**

539

540

541 **Figure 2**

545 **Figure 3**



546

547

548 **Supplemental Digital Content**

549 Supplementary Material (pdf document containing the model description, model
550 calibration and supplementary results).

551