Isometry Pursuit: Convex Recovery of Orthonormal Column Submatrices

Anonymous authors
Paper under double-blind review

Abstract

Isometry pursuit is a convex algorithm for identifying orthonormal column-submatrices of wide matrices. It consists of a vector normalization followed by multitask basis pursuit. Applied to Jacobians of putative coordinate functions, it helps identity locally isometric embeddings from within interpretable dictionaries. We provide theoretical and experimental results justifying this method, including a proof with realistic assumptions that such isometric submatrices, should they exist, are contained within the obtained support. For problems involving coordinate selection and diversification, it offers a synergistic alternative to greedy and brute force search.

1 Background

Our algorithm is motivated by a classic question in numerical linear algebra.

1.1 Problem

Our goal is, given a matrix $X \in \mathbb{R}^{D\times P}$, to select a subset $S \subset [P]$ with $|S| = D$ such that $X_{.S}$ is as orthonormal as possible in a computationally efficient way. To this end, we define a ground truth loss function that measures orthonormalness, and then introduce a surrogate loss function that convexifies the problem so that it may be efficiently solved. Figure **??** and **??** provide a visual depiction of this problem when $D = 2$ and $P = 30$.

1.2 Isometry

The mathematical construct central to our estimation problem is the isometric linear map.

Definition 1 *A matrix $X \in \mathbb{R}^{D\times P}$ is an **isometry** or, equivalently, **unitary** if*

$$X^T X = I_D. \tag{1}$$

Note that necessarily, $P \geq D$.

Definition 1 has several equivalent formulations.

Proposition 1 *The singular values $\sigma_1(X) \dots \sigma_D(X)$ are equal to 1 if and only if X is an isometry.*

On the other hand, if $P = D$, X

Proposition 2 *The component vectors $X_{.}1 \dots X_{.D} \in \mathbb{R}^D$ form an orthonormal matrix if and only if, for all $d_1, d_2 \in [D], \langle X_{.d_1}, X_{.d_2}\rangle = \begin{cases} 1 & d_1 = d_2 \\ 0 & d_1 \neq d_2 \end{cases}$.*

That is, X is an isometry if its columns are orthonormal.

The applications of isometry are manifold. Overall, is a classically interesting problem. It is naturally a type of diversification. Embedding algorithms like Local Tangent Space

Alignment (**?**), Multidimensional Scaling (**?**) and Isomap (**?**) non-parametrically estimate embeddings that are as isometric as possible. Note that an isometric embedding is a map whose differential induces isometry between tangent spaces. The inspiration for this paper comes from an approach that stitches together isometries selected from a dictionary to form global embeddings (Kohli et al., 2021). This method constructs such isometries through greedy search, with putative dictionary features added one at a time.

1.3 Functions minimized by isometries

The distortion loss function used the evaluate isometricness in (Kohli et al., 2021) reduces in the linear tangent space limit to the condition number. The *condition number* (**?**Boyd and Vandenberghe, 2004) $\frac{\sigma_1(X)}{\sigma_D(X)}$ measures the ratio of the largest and smallest singular values of a matrix A. While it is minimized by an isometric function, it is also minimized by any conformal (i.e. low distortion) linear map.

A class of functions more suitable for measuring isometry are S-divergences (**?**). S-divergences define a metric $D(P, P')$ on positive definite matrices P, P'. If those matrices are $X^T X$ and I_D, the S-divergence distance is

$$D(X^T X, I_D) = (\log\det(\frac{X^T X + I_D}{2}) - \frac{1}{2}\log\det(X^T X))^{\frac{1}{2}}. \tag{2}$$

This is a log singular value penalty. This is because, first of all,

$$\frac{1}{2}\log\det(X^T X) = \frac{1}{2}\log\prod_{d=1}^{D}\sigma_d(X)^2 \tag{3}$$

$$= \log\prod_{d=1}^{D}\sigma_d(X) \tag{4}$$

$$= \sum_{d=1}^{D}\log\sigma_d(X). \tag{5}$$

Secondly, I_D and $X^T X$ are diagonalizable in the same basis, so

$$\log\det(\frac{X^T X + I_D}{2}) = \log\prod_{d=1}^{D}\frac{\sigma_d(X)^2 + 1}{2} \tag{6}$$

$$= \sum_{d=1}^{D}\log\frac{\sigma_d(X)^2 + 1}{2}. \tag{7}$$

Then,

$$D(X^T X, I_D)^2 = \sum_{d=1}^{D}(\log\frac{\sigma_d(X)^2 + 1}{2} - \log\sigma_d(X)) \tag{8}$$

$$= \sum_{d=1}^{D}\log\frac{\sigma_d(X)^2 + 1}{2\sigma_d(X)} \tag{9}$$

$$= \sum_{d=1}^{D}\log\frac{\sigma_d(X) + \sigma_d(X)^{-1}}{2}. \tag{10}$$

By Proposition 1, this function is uniquely minimized by isometric matrices. TODO: explain how works for PSD not just PD.

1.4 Best subset selection

Given a matrix $X \in \mathbb{R}^{D\times P}$, we compare algorithmic paradigms for solving problems of the form

$$\arg\min_{S\in\binom{[P]}{D}} \ell(X_{\cdot S}) \tag{11}$$

where $\binom{[P]}{D} = \{A \subseteq [P] : |A| = D\}$. Brute force algorithms consider all possible solutions. These algorithms are conceptually simple, but have the often prohibitive time complexity $O(C_\ell P^D)$ where C_ℓ is the cost of evaluating ℓ. Greedy algorithms consist of iteratively adding one element at a time to S. This algorithms have time complexity $O(C_\ell PD)$ and so are computationally more efficient than brute force algorithms, but can get stuck in local minima. Formal definitions are given in Section 2.1.

Sometimes, it is possible to introduce an objective which convexifies problems of the above form (**???**). Solutions

$$\arg\min f(\beta) : Y = X\beta \tag{12}$$

to the overcomplete regression problem $Y = X\beta$ are a classic example (**?**). When $f(\beta) = \|\beta\|_0$, this problem is non-convex, and is thus suitable for greedy or brute algorithms, but when $f(\beta) = \|\beta\|_1$, the problem is convex, and may be solved efficiently via interior-point methods. When the equality constraint is relaxed, Lagrangian duality may be used to reformulate as a so-called Lasso problem, which leads to an even richer set of optimization algorithms (**?**).

The form of basis pursuit that we apply is inspired by the group basis pursuit (Koelle et al., 2022; **?**). Multitask basis pursuit is a special form of group basis pursuit which we call differently since grouping is dependent only on the structure of matrix-valued response variable y. The objective $f(\beta) = \|\beta\|_{1,2} := \sum_{p=1}^{P} \|\beta_{p\cdot}\|_2$ creates joint sparsity across entire rows $\beta_{p\cdot}$ (**???**). It was used in Koelle et al. (2022) to select independent subsets among interpretable features.

References

Samson J Koelle, Hanyu Zhang, Marina Meila, and Yu-Chia Chen. Manifold coordinates with physical meaning. *J. Mach. Learn. Res.*, 23(133):1–57, 2022.

Samson J Koelle, Hanyu Zhang, Octavian-Vlad Murad, and Marina Meila. Consistency of dictionary-based manifold learning. In Sanjoy Dasgupta, Stephan Mandt, and Yingzhen Li, editors, *Proceedings of The 27th International Conference on Artificial Intelligence and Statistics*, volume 238 of *Proceedings of Machine Learning Research*, pages 4348–4356. PMLR, 2024.

Dhruv Kohli, Alexander Cloninger, and Gal Mishne. LDLE: Low distortion local eigenmaps. *J. Mach. Learn. Res.*, 22, 2021.

Stephen P Boyd and Lieven Vandenberghe. *Convex Optimization*. Cambridge University Press, March 2004.

Ulrike Schneider and Patrick Tardivel. The geometry of uniqueness, sparsity and clustering in penalized estimation. *J. Mach. Learn. Res.*, 23(331):1–36, 2022.

Ryan J Tibshirani. The lasso problem and uniqueness. *arXiv [math.ST]*, June 2012.

Aaron Mishkin and Mert Pilanci. The solution path of the group lasso. 2022.

Brendan O'Donoghue, Eric Chu, Neal Parikh, and Stephen Boyd. Conic optimization via operator splitting and homogeneous self-dual embedding. *arXiv [math.OC]*, December 2013.

T Hastie, R Tibshirani, and M Wainwright. Statistical learning with sparsity: The lasso and generalizations. May 2015.

Nicolai Meinshausen. Relaxed lasso. *Comput. Stat. Data Anal.*, 52(1):374–393, September 2007.

Tim Hesterberg, Nam Hee Choi, Lukas Meier, and Chris Fraley. Least angle and ℓ_1 penalized regression: A review. February 2008.

Faming Liang, Runmin Shi, and Qianxing Mo. A split-and-merge approach for singular value decomposition of large-scale matrices. *Stat. Interface*, 9(4):453–459, September 2016.

Brendan O'Donoghue, Eric Chu, Neal Parikh, and Stephen Boyd. Conic optimization via operator splitting and homogeneous self-dual embedding. *Journal of Optimization Theory and Applications*, 169(3):1042–1068, June 2016. URL `http://stanford.edu/~boyd/papers/scs.html`.

Steven Diamond and Stephen Boyd. CVXPY: A Python-embedded modeling language for convex optimization. *Journal of Machine Learning Research*, 17(83):1–5, 2016.

Akshay Agrawal, Robin Verschueren, Steven Diamond, and Stephen Boyd. A rewriting system for convex optimization problems. *Journal of Control and Decision*, 5(1):42–60, 2018.

CVXPY Developers. Sparse solution with cvxpy. `https://www.cvxpy.org/examples/applications/sparse_solution.html`. Accessed: 2024-07-11.

Bingsheng He and Xiaoming Yuan. On the $o(1/n)$ convergence rate of the douglas–rachford alternating direction method. *SIAM J. Numer. Anal.*, 50(2):700–709, January 2012.

Thomas H. Cormen, Charles E. Leiserson, Ronald L. Rivest, and Clifford Stein. *Introduction to Algorithms, Third Edition*. The MIT Press, 3rd edition, 2009. ISBN 0262033844.

Stuart Russell and Peter Norvig. *Artificial Intelligence: A Modern Approach*. Prentice Hall Press, USA, 3rd edition, 2009. ISBN 0136042597.

Travis Dick, Eric Wong, and Christoph Dann. How many random restarts are enough. 2014. URL https://api.semanticscholar.org/CorpusID:9473630.

Stephen Boyd. Distributed optimization and statistical learning via the alternating direction method of multipliers. *Found. Trends® Mach. Learn.*, 3(1):1–122, 2010.

2 Supplement

This section contains algorithms, proofs, and experiments in support of the main text.

2.1 Algorithms

We give definitions of the brute and greedy algorithms for the combinatorial problem studied in this paper. The brute force algorithm is computationally intractable for all but the smallest problems, but always finds the global minima.

BruteSearch(Matrix $X \in \mathbb{R}^{D \times P}$, objective f)

1: **for** each combination $S \subseteq \{1, 2, \ldots, P\}$ with $|S| = D$ **do**
2: Evaluate $f(X_{.S})$
3: **end for**
4: **Output** the combination S^* that minimizes $f(X_{.S})$

Greedy algorithms are computationally expedient but can get stuck in local optima (Cormen et al., 2009; Russell and Norvig, 2009), even with randomized restarts (Dick et al., 2014).

GreedySearch(Matrix $X \in \mathbb{R}^{D \times P}$, objective f, selected set $S = \emptyset$, current size $d = 0$)

1: **if** $d = D$ **then**
2: **Return** S
3: **else**
4: **Initialize** $S_{\text{best}} = S$
5: **Initialize** $f_{\text{best}} = \infty$
6: **for** each $p \in \{1, 2, \ldots, P\} \setminus S$ **do**
7: **Evaluate** $f(X_{.(S \cup \{p\})})$
8: **if** $f(X_{.(S \cup \{p\})}) < f_{\text{best}}$ **then**
9: **Update** $S_{\text{best}} = S \cup \{p\}$
10: **Update** $f_{\text{best}} = f(\mathcal{X}_{.(S \cup \{p\})})$
11: **end if**
12: **end for**
13: **Return** GreedySearch(X, f, S_{best}, $d + 1$)
14: **end if**

2.2 Proofs

2.2.1 Proof of Proposition ??

In this proof we first show that the penalty $\|\beta\|_{1,2}$ is unchanged by unitary transformation of β.

Proposition 3 *Let $U \in \mathbb{R}^{D \times D}$ be unitary. Then $\|\beta\|_{1,2} = \|\beta U\|$.*

Proof:

$$\|\beta U\|_{1,2} = \sum_{p=1}^{P} \|\beta_{p.} U\| \tag{13}$$

$$= \sum_{p=1}^{P} \|\beta_{p.}\| \tag{14}$$

$$= \|\beta\|_{1,2} \tag{15}$$

□

We then show that this implies that the resultant loss is unchanged by unitary transformation of X.

Proposition 4 *Let $U \in \mathbb{R}^{D\times D}$ be unitary. Then $\widehat{\beta}(UX) = \widehat{\beta}(X)U$.*

Proof:

$$\widehat{\beta}(UX) = \arg\min_{\beta\in\mathbb{R}^{P\times D}} \|\beta\|_{1,2} \ : \ I_D = UX\beta \tag{16}$$

$$= \arg\min_{\beta\in\mathbb{R}^{P\times D}} \|\beta\|_{1,2} \ : \ U^{-1}U = U^{-1}UX\beta U \tag{17}$$

$$= \arg\min_{\beta\in\mathbb{R}^{P\times D}} \|\beta\|_{1,2} \ : \ I_D = X\beta U \tag{18}$$

$$= \arg\min_{\beta\in\mathbb{R}^{P\times D}} \|\beta U\|_{1,2} \ : \ I_D = X\beta U \tag{19}$$

$$= \arg\min_{\beta\in\mathbb{R}^{P\times D}} \|\beta\|_{1,2} \ : \ I_D = X\beta. \tag{20}$$

□

2.2.2 Proof of Proposition ??

In this proof we first show that the orthonormal submatrix $X_{.S}$ is contained within the set $\widehat{\beta}_{MBP}(w(X), I_D)$.

Proposition 5 *Let $X_{.S}$ be a rank D orthonormal submatrix of X. Then $S \in S(\beta) : \beta \in \widehat{\beta}_{MBP}(w(X), I_D))$.*

Proof: By Proposition **??** without loss of generality, let $X_{.S} = I_D$. Also, without loss of generality let $S = \{1 \dots D\}$. We show that $\beta = [I_D \ 0]$ satisfies the KKT conditions for $X = [I_D \ X_{-S}]$.

Recall that the KKT conditions for a convex optimization problem are a set of conditions on an element of the domain of the optimization algorithm that, if satisfied, certify that the element is a solution. To write out the KKT conditions, we define the Lagrangian

$$\mathcal{L}(X, \beta, \nu) = \|\beta\|_{1,2} + \nu^T(I_D - w(X)\beta) \tag{21}$$

where $\nu \in \mathbb{R}^{D\times D}$ is the dual variable.

The KKT conditions are then

- Primal feasibility: $w(X)\beta = I_D$
- Stationarity: there exists a dual variable ν such that $0 \in \partial\|\beta\|_{1,2} - w(X)^T\nu$ where ∂ is the subdifferential operator.

where dual feasibility and complementary slackness are ignorable by virtue of the absence of inequality constraints.

First note that in our case, primal feasibility is satisfied by X being rank D and w not effecting the rank of the transformed matrix since it only rescales length and not direction.

For stationarity, recall that

$$\|\beta\|_{1,2} = \sum_{p=1}^{P} \|\beta_{p.}\|_2. \tag{22}$$

Then, by the definition of subdifferential

$$\partial\|\beta\|_{1,2} = \begin{bmatrix} I_D \\ v_{d+1} \\ \dots \\ v_P \end{bmatrix} \tag{23}$$

where $v_p \in \mathbb{R}^D$ can be any vector satisfying $\|v_p\|_2 \leq 1$.

We therefore must show that there exists a ν that satisfies

$$\begin{bmatrix} I_D \\ w(X_{-S})^T \end{bmatrix} \nu = \begin{bmatrix} I_D \\ V_{-S} \end{bmatrix}. \tag{24}$$

At this point we can see that in fact I_D is an appropriate choice of ν since normalization by w leads to vectors satisfying $\|w(X)\|_2 \leq 1$.

Since for this β, $S(\beta) = S$, we have shown that S is the support of a solution to $\beta_{MBP}(w(X), I_D)$. □

The next part of the proof handles the presence of non-unique solutions.

Proposition 6 $S(\widehat{\beta}(X)) = \cup\, S(\beta) \;:\; \beta \in \beta_{MBP}(w(X), I_D)$

Proof: The proposition follows from geometric properties of the solution set. We ignore the case for singleton solutions, for which the proposition is self-evident. Recall that $\widehat{\beta}(X) := \widehat{\beta}^{\ell_2}_{MBP}(w(X), I_D) = \arg\min_{\beta \in \mathbb{R}^{P\times D}} \|\beta\|_F : \beta \in \widehat{\beta}_{MBP}(X, I_D)$. By definition of $\widehat{\beta}_{MBP}(X, I_D)$, $\|\beta_{p\cdot}\|_2 : \beta \in \widehat{\beta}_{MBP}(X, I_D)$ is an affine linear subspace of $\mathbb{R}^P$. The projection of the vector of minimum ℓ_2 distance from the origin to a affine space intersects that space perpendicularly. However, for all $\|\beta_{\cdot p}\|_2$ such that β is on the intersection of this affine space with the coordinate hyperplanes formed by fixing any choices of $\|\beta_{\cdot p}\|_2 = 0$, this vector doesn't intersect this space perpendicularly. Thus, $\widehat{\beta}(X)$ is not on the intersection of the solution polytope $\widehat{\beta}_{MBP}(w(X), I_D)$ with the coordinate hyperplanes. It is therefore maximally non-sparse with respect to any of the minimizing solutions. □

2.2.3 Proof of Proposition **??**

X_S admits a QR decomposition $X_S = QR$ where Q is orthogonal and R is upper triangular. By Proposition **??**, we can assume without loss of generality that $Q = I_D$. That is, X_S is upper triangular. Thus, $\beta = X_S^{-1}$ is upper triangular with diagonal elements $\beta_{dd} = R^{-1}_{dd}$. $\|\beta_{d\cdot}\|_2 \geq \beta_{dd}$, with equality only occurring when the off-diagonal elements are 0. Therefore, $\|\beta_{1,2}\| \geq D$ with equality occurring only when $\beta = I_D$.

2.2.4 Proof of Proposition **??**

Proof: The augmented split Lagrangian minimized by ADMM is

$$\mathcal{L}_{\rho,\gamma,\nu}(X, \beta, z) = \|z\|_{1,2} + \mathrm{tr}(\nu^T(I_D - w(X)\beta)) + \rho\|I_D - w(X)\beta\|_F^2 + \gamma\|\beta - z\|_F^2 \tag{25}$$

This minimization problem is

$$\arg \min_{\beta, z \in P\times D} \mathcal{L}_{\rho,\gamma,\nu}(X, \beta, z) \tag{26}$$

For a more detailed explanation of the reasons to prefer optimizing the split augmented Lagrangian over direct optimization of Equation **??** see Boyd (2010).

The three main subroutines (30 - 34) called by ADMM determine the per iteration complexity. The update rules of the ADMM algorithm used to optimize the augmented split Lagrangian are

$$\beta^{k+1} = \arg \min_{\beta \in P\times D} \mathcal{L}_{\rho,\gamma,\nu}(X, \beta, z^k) \tag{27}$$

$$z^{k+1} = \arg \min_{z \in P\times D} \mathcal{L}_{\rho,\gamma,\nu}(X, \beta^{k+1}, z) \tag{28}$$

$$\nu^{k+1} = \nu^k + \rho(I_D - w(X)\beta^{k+1}) \tag{29}$$

We will analyze the computational complexity of each of these steps by rewriting the update rules according to the specifics of the isometry pursuit problem.

The beta update Equation 30 projects the solution onto the feasible set.

$$\beta^{k+1} = \arg\min_{\beta \in P \times D} \operatorname{tr}({\nu^k}^T (I_D - w(X)\beta)) + \rho \|I_D - w(X)\beta\|_2^2 + \gamma \|\beta - z^k\|_2^2 \tag{30}$$

$$= (\rho\, w(X)^T w(X) + \gamma I_D)^{-1} (\rho\, w(X)^T I_D + \gamma z^k + w(X)^T \nu^k) \tag{31}$$

The computational complexity of this computation is dominated by the inverse - that is, $O(D^2 P)$.

Subsequently, the z update Equation 32 is soft-thresholding operation onto a contracted level set $\|\|_{1,2}$ ball.

$$z^{k+1} = \arg\min_{z \in P \times D} \|z\|_{1,2} + \gamma \|z - \beta^{k+1}\|_2^2 \tag{32}$$

$$= \begin{cases} (1 - \frac{1}{\gamma \|\beta_{j\cdot}^{k+1}\|_2})(\beta_{j\cdot}^{k+1}) \text{ if } \|\beta_{j\cdot}^{k+1}\|_2 > \frac{1}{\gamma} \\ 0 \text{ otherwise} \end{cases} \tag{33}$$

The computational complexity of this operation is $O(PD)$.

Finally, the ν update Equation 34 increases the squeezing of the z and β variables towards to each other.

$$\nu^{k+1} = \nu^k + \rho(I_D - w(X)\beta^{k+1}) \tag{34}$$

In this step, the dual variable ν that controls the penalty accrued by the splitting is updated in the direction of the constraint violation. The computational complexity of this operation is dominated by that of the matrix multiplication $w(X)\beta^{k+1}$. That is, it is $O(D^2 P)$. □

2.3 Partial derivatives of candidate losses

We summarize the column-wise derivatives of the losses in Table **??**. Throughout this section let $X = [x_1, \ldots, x_D] \in \mathbb{R}^{D \times D}$ be full rank. For a scalar loss $L(X)$, write

$$\nabla_X L = \left[\frac{\partial L}{\partial x_1}, \ldots, \frac{\partial L}{\partial x_D} \right]. \tag{35}$$

All formulas are stated on smooth strata: rank deficient matrices, repeated singular values, and zero rows of pseudoinverse factors should be handled using subgradients.

We will use two common identities. First, if $X = U \operatorname{diag}(\sigma) V^\top$ is a singular value decomposition and $L(X) = \sum_{d=1}^{D} \phi(\sigma_d(X))$, then

$$\nabla_X L = U \operatorname{diag}(\phi'(\sigma_1), \ldots, \phi'(\sigma_D)) V^\top. \tag{36}$$

Second, for normalized losses let $N = n_q^{\times D}(X)$. If $M = \nabla_N L$, then the column-wise chain rule is

$$\frac{\partial L}{\partial x_p} = \mathrm{D} n_q(x_p)^\top M_{\cdot p}. \tag{37}$$

For $r = \|x\|_2$, write $n_q(x) = s(r)x$ with $s(r) = q(r)/r$. Then

$$\mathrm{D} n_q(x) = s(r) I_D + \frac{s'(r)}{r} x x^\top, \tag{38}$$

$$s'(r) = \frac{q'(r)}{r} - \frac{q(r)}{r^2}, \tag{39}$$

$$q'(r) = -\frac{2e(e^r - r^{-2} e^{1/r})}{(e^r + e^{1/r})^2}. \tag{40}$$

Since $\mathrm{D} n_q(x)$ is symmetric, the transpose in this chain rule may be omitted.

Distortion. Let $\sigma_1 = \sigma_{\max}(X)$ and $\sigma_D = \sigma_{\min}(X)$ with corresponding singular vectors u_1, v_1 and u_D, v_D. For

$$L_{\text{dist}}(X) = \frac{\sigma_1}{\sigma_D} - 1, \tag{41}$$

The spectral gradient identity gives

$$\nabla_X L_{\text{dist}} = \frac{u_1 v_1^\top}{\sigma_D} - \frac{\sigma_1}{\sigma_D^2} u_D v_D^\top. \tag{42}$$

The derivative is insensitive to uniform rescaling, which is the sense in which this loss does not prefer unit length.

Isometry loss. For

$$L_{\text{iso}}(X) = \sum_{d=1}^{D} g(\sigma_d(X)) - D, \tag{43}$$

we have

$$g'(t) = \frac{e^t - t^{-2} e^{1/t}}{2e}. \tag{44}$$

Thus

$$\nabla_X L_{\text{iso}} = U \operatorname{diag}(g'(\sigma_1), \ldots, g'(\sigma_D)) V^\top. \tag{45}$$

S-divergence to identity. Let $G = X^\top X$ and

$$L_{\text{sdiv}}(X) = \log\det\left(\frac{G + I_D}{2}\right) - \frac{1}{2}\log\det(G). \tag{46}$$

The constant $\log\det(1/2)$ does not contribute to the derivative. Using $d\log\det A = \operatorname{tr}(A^{-1}dA)$ and $dG = dX^\top X + X^\top dX$,

$$\nabla_X L_{\text{sdiv}} = 2X(G + I_D)^{-1} - XG^{-1}. \tag{47}$$

Normalized log-det volume. Let $N = n_q^{\times D}(X)$ and

$$L_{\text{nld}}(X) = -\sum_{d=1}^{D} \log \sigma_d(N). \tag{48}$$

For square full-rank N, this is $-\log|\det N|$, so

$$\nabla_N L_{\text{nld}} = -N^{-\top}. \tag{49}$$

Therefore, by the column-wise chain rule,

$$\frac{\partial L_{\text{nld}}}{\partial x_p} = -\mathrm{D}n_q(x_p)^\top (N^{-\top})_{\cdot p}. \tag{50}$$

Frobenius pseudoinverse. Let $N = n_q^{\times D}(X)$ and $B = N^{-1}$. For

$$L_{\text{frob}}(X) = \|N^{-1}\|_F - \sqrt{D}, \tag{51}$$

the differential identity $dB = -B(dN)B$ gives

$$\nabla_N L_{\text{frob}} = -\frac{N^{-\top} N^{-1} N^{-\top}}{\|N^{-1}\|_F}. \tag{52}$$

Thus

$$\frac{\partial L_{\text{frob}}}{\partial x_p} = \mathrm{D}n_q(x_p)^\top \left(-\frac{N^{-\top} N^{-1} N^{-\top}}{\|N^{-1}\|_F}\right)_{\cdot p}. \tag{53}$$

Isometry pursuit proxy. Let $N = n_q^{\times D}(X)$ and $B = N^{-1}$. The differentiable proxy for the IsometryPursuit table row is

$$L_{\mathrm{ip}}(X) = \|N^{-1}\|_{1,2} - D. \tag{54}$$

Define $R \in \mathbb{R}^{D\times D}$ by


$$R_{p\cdot} = \frac{B_{p\cdot}}{\|B_{p\cdot}\|_2}. \tag{55}$$

Then $R \in \partial\|B\|_{1,2}$ on the smooth stratum where every row of B is nonzero, and

$$\nabla_N L_{\mathrm{ip}} = -B^\top R B^\top. \tag{56}$$

The column-wise gradient is therefore

$$\frac{\partial L_{\mathrm{ip}}}{\partial x_p} = \mathrm{D}n_q(x_p)^\top (-B^\top R B^\top)_{\cdot p}. \tag{57}$$



(a) Wine (b) Iris (c) Ethanol

Figure 1: Support cardinalities for each dataset after IsometryPursuit.

2.4 Support cardinalities

Figure 1 plots the distribution of $|\widehat{S}_{IP}|$ from Table **??** in order to contextualize the reported means. While typically $|\widehat{S}_{IP}| << P$, there are cases for Ethanol where this is not the case that drive up the means. Further investigation into the extreme failure of isometry pursuit to sparsify in the the minor chunk of the bimodal Ethanol support cardinality distribution is warranted.

2.5 Timing

While wall-time of algorithms is a non-theoretical quantity that depends on implementation details, it provides valuable context for practitioners. We therefore report the following runtimes on a 2021 Macbook Pro. The particularly high variance for brute force search in the second step of TwoStageIsometryPursuit is likely due to the large cardinalities reported in Figure 1.

Name	IP	2nd stage	Greedy
Iris	1.24 ± 0.02	0.00 ± 0.00	0.02 ± 0.00
Wine	2.32 ± 0.17	0.13 ± 0.12	0.03 ± 0.00
Ethanol	8.38 ± 0.57	0.55 ± 1.08	0.07 ± 0.01

Table 1: Algorithm runtimes in seconds across replicates.

Example 1

$$X_S = \begin{bmatrix} \frac{1}{2} & \frac{-1}{2} & \frac{1}{2} & \frac{-1}{2} & \frac{1}{2} & \frac{-1}{2} \\ \frac{\sqrt{3}}{2} & \frac{\sqrt{3}}{2} & 0 & 0 & & \\ 0 & 0 & \frac{\sqrt{3}}{2} & \frac{\sqrt{3}}{2} & 0 & 0 \\ 0 & 00 & 0 & \frac{\sqrt{3}}{2} & \frac{\sqrt{3}}{2} & \end{bmatrix}. \tag{58}$$

Proof: Let

By Given the symmetry of the second and third columns around the y axis, the loss will not change regardless of whether we upweight one or downweight the other, these changes necessarily being in conjunction. We therefore assume that their loadings with respect to the second column of the target matrix (the D dimensional identity) are the same. Clearly, they will both then have value of $\frac{1}{\sqrt{3}}$. The question then becomes, what should the values of the first column in the loading matrix β be?

We can answer this question with a simple differential equation. Let b denote the constraint violation $1 - X_{1.}\beta_{.1}$.

Then

$$\frac{\partial b}{\partial \beta_{11}} = \frac{\partial b}{\partial \beta_{21}} \tag{59}$$

$$= \frac{1}{2} \tag{60}$$

On the other hand, due to the value of $\frac{1}{\sqrt{3}}$ already assumed for the second loading of the second and third features,

$$\frac{\partial l}{\partial \beta_{11}} = \frac{\partial l}{\partial \beta_{21}} \tag{61}$$

$$= \frac{\partial}{\partial x}\sqrt{\frac{1}{3} + x^2} \tag{62}$$

$$= \frac{1}{2}(\frac{1}{3} + x^2)^{-\frac{1}{2}}(2x) \tag{63}$$

$$= \frac{x}{\sqrt{\frac{1}{3} + x^2}}. \tag{64}$$

This means that the tradeoff

$$\frac{\partial l}{\partial b} = \frac{1}{2x}\sqrt{\frac{1}{3} + x^2}. \tag{65}$$

In contrast, $\frac{\partial l}{\partial \beta_{11}} = 1$ and similarly for its constraint violation. Solving for the equilibrium, we get

$$\frac{1}{2x}\sqrt{\frac{1}{3} + x^2} = 1 \tag{66}$$

$$\frac{1}{3} + x^2 = 4x^2 \tag{67}$$

$$\frac{1}{3} = 3x^2 \tag{68}$$

$$x = \frac{1}{\sqrt{3}}. \tag{69}$$

This leads to a value for l of

$$\sqrt{\frac{1}{3} + \frac{1}{9}} + \sqrt{\frac{1}{3} + \frac{1}{9}} + \frac{2}{3} \tag{70}$$

$$= \sqrt{\frac{4}{9}} + \sqrt{\frac{4}{9}} + \frac{2}{3} \tag{71}$$

$$= \frac{2}{3} + \frac{2}{3} + \frac{2}{3} \tag{72}$$

□