

# Tutte's Flow Conjectures



Candidate Number: 579167

Part C Dissertation submitted for the degree of  
*Master of Mathematics and Philosophy*

Hilary 2016

## Abstract

For  $G = (V, E)$  a directed multigraph and  $H$  an abelian group, a map  $f : E \rightarrow H$  is an  $H$ -flow if  $f(e) \neq 0$  for all  $e \in E$  and  $f$  obeys Kirchhoff's law: for all  $v \in V$

$$\sum_{e \in \delta^+(v)} f(e) = \sum_{e \in \delta^-(v)} f(e)$$

If  $H = \mathbb{Z}$ , the group of integers under addition, and  $k$  is a positive integer such that  $-k < f(e) < k$  for every edge  $e$ , we say that  $f$  is a  $k$ -flow.

W. T. Tutte [5, 6, 7] made a series of conjectures asserting certain weak conditions under which a multigraph  $G$  has a 3-, 4- or 5-flow. We will present the statements of these alongside a number of results about group-valued and  $k$ -flows including sufficient conditions for low  $k$ -flows and the duality between flow and colouring problems. As an example of this we shall demonstrate that the theorem of Grötzsch [2] that *every triangle free planar graph is 3-colourable* is a dual to the 3-flow conjecture with an added assumption of planarity. Finally we will give an exposition of Thomassen's [4] recent proof of a weakening of the 3-flow conjecture and discuss subsequent work.

## Acknowledgments

This essay was written in  $\text{\LaTeX}$  and the source file based upon a dissertation template by [Federico Maggi](#). The template was released under the Creative Commons [BY-NC-SA 3.0](#) license, in accordance with which the source files for this essay are available, under the same license, at my [GitHub repository](#).

# Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Seymour's 6-Flow Theorem</b>	<b>2</b>
	<b>Bibliography</b>	<b>3</b>



# Seymour's 6-Flow Theorem

2

**Lemma 2.1.** *For  $k > 2$ , let  $G = (V, E)$  be a bridgeless graph with no nowhere-zero  $k$ -flow such that  $G$  has  $|V| + |E|$  minimal. Then  $G$  is simple, cubic and 3-connected.*

Suppose there were a loop  $e$  in  $G$ , let  $G' = G - e$ . Any bridge in  $G'$  would also be a bridge in  $G$ , so  $G'$  must be bridgeless. Therefore, by the minimality of  $G$ , there must be a nowhere-zero  $k$ -flow  $f$  on  $G'$ . But then we can extend  $f$  to a nowhere-zero  $k$ -flow on  $G$  by setting  $f(e) = l$  for some  $0 < l < k$ , contradicting that  $G$  has no such flow. Thus  $G$  contains no loops. Now suppose there are vertices  $v, w \in V$  with multiple edges  $e_1, e_2, \dots, e_n$  between them and let  $G'' = G/e_1$ . Again any bridge in  $G''$  would be a bridge in  $G$  so that  $G''$  is bridgeless and by the minimality of  $G$  has a nowhere-zero  $k$ -flow  $f'$ . We will construct a flow  $f$  on  $G$  from  $f'$ . For all  $e \in E \setminus \{e_1, e_2\}$ , define  $f(e) = f'(e)$ . Since  $k > 2$ , we can either increase or decrease  $f'(e_2)$  by one such that  $|f'(e_2)|$  remains strictly between 0 and  $k$ . If we may increase  $f'(e_2)$  by one then define  $f(e_2) = f'(e_2) + 1$  and

$$f(e_1) = \begin{cases} 1 & \text{if } e_1 \text{ is oriented in the opposite direction to } e_2 \\ -1 & \text{if } e_1 \text{ is oriented in the same direction as } e_2 \end{cases}$$

Otherwise, define  $f(e_2) = f'(e_2) - 1$  and set

$$f(e_1) = \begin{cases} 1 & \text{if } e_1 \text{ is oriented in the opposite direction to } e_2 \\ -1 & \text{if } e_1 \text{ is oriented in the same direction as } e_2 \end{cases}$$

Thus (with respect to  $f'$ )  $f$  fixes the flow through all vertices except  $v$  and  $w$  and our construction ensures that at  $v$  and  $w$  any changes to the flow into the vertex is balanced by an equal and opposite change to the flow out of the vertex and vice versa. This ensures that  $f$  obeys Kirchhoff's Law, as  $f'$  does. In addition,  $0 < |f(e)| < k$  for all  $e \in E$ , so  $f$  is a nowhere-zero  $k$ -flow on  $G$ . This contradicts that  $G$  has no such flow and so we have shown that  $G$  must be simple.

# Bibliography

---

- [1] Reinhard Diestel. *Graph Theory*. Number 173 in Graduate Texts in Mathematics. Springer, third edition, 1997. [Chapter 6 in particular].
- [2] Herbert Grötzsch. Zur theorie der diskreten gebilde. vii. ein dreifarbensatz für dreikreisfreie netze auf der kugel. *Wiss. Z. Martin-Luther Univ. Halle-Wittenberg*, 8:109–120, 1958/1959.
- [3] László Miklós Lovász, Carsten Thomassen, Yezhou Wu, and Cun-Quan Zhang. Nowhere-zero 3-flows and modulo  $k$ -orientations. *Journal of Combinatorial Theory, Series B*, 103(5):587 – 598, 2013.
- [4] Carsten Thomassen. The weak 3-flow conjecture and the weak circular flow conjecture. *Journal of Combinatorial Theory, Series B*, 102(2):521 – 529, 2012.
- [5] William Thomas Tutte. A contribution to the theory of chromatic polynomials. *Canadian Journal of Mathematics*, 6:80–91, 1954.
- [6] William Thomas Tutte. On the algebraic theory of graph colourings. *Journal of Combinatorial Theory*, 1:15–50, 1966.
- [7] William Thomas Tutte. 3-flow conjecture. Unpublished, 1972.