

The Rössler System

Steven Rosendahl

Introduction

- Created by Otto Rössler in 1976

Introduction

- Created by Otto Rössler in 1976
- System constructed to display simplest possible *Strange Attractor*

Introduction

- Created by Otto Rössler in 1976
- System constructed to display simplest possible *Strange Attractor*

Definition

An **attractor** is a set of values towards which a system moves when initial conditions are *near* the attractor.

Introduction

- Created by Otto Rössler in 1976
- System constructed to display simplest possible *Strange Attractor*

Definition

An **attractor** is a set of values towards which a system moves when initial conditions are *near* the attractor. An attractor is called **strange** if it exhibits fractal behavior.

Introduction

- Created by Otto Rössler in 1976
- System constructed to display simplest possible *Strange Attractor*

Definition

An **attractor** is a set of values towards which a system moves when initial conditions are *near* the attractor. An attractor is called **strange** if it exhibits fractal behavior.

- Strange attractors are often associated with chaotic systems.

System

$$\dot{x} = -y - z$$

$$\dot{y} = x + ay$$

$$\dot{z} = b + z(x - c)$$

System

$$\dot{x} = -y - z$$

$$\dot{y} = x + ay$$

$$\dot{z} = b + z(x - c)$$

- a, b, c are responsible for attractive behavior

System

$$\dot{x} = -y - z$$

$$\dot{y} = x + ay$$

$$\dot{z} = b + z(x - c)$$

- a, b, c are responsible for attractive behavior
- Want to analyze the effect small a and b have on system

System

$$\dot{x} = -y - z$$

$$\dot{y} = x + ay$$

$$\dot{z} = b + z(x - c)$$

- a, b, c are responsible for attractive behavior
- Want to analyze the effect small a and b have on system
 - Is the system globally stable? Locally?

System

$$\dot{x} = -y - z$$

$$\dot{y} = x + ay$$

$$\dot{z} = b + z(x - c)$$

- a, b, c are responsible for attractive behavior
- Want to analyze the effect small a and b have on system
 - Is the system globally stable? Locally?
 - When does chaotic behavior occur?

First Glance

- Finding fixed points

First Glance

■ Finding fixed points

```
Solve[f[x, y, z] == 0 && g[x, y, z] == 0 && h[x, y, z] == 0, {x, y, z}]
```

First Glance

■ Finding fixed points

`Solve[f[x, y, z] == 0 && g[x, y, z] == 0 && h[x, y, z] == 0, {x, y, z}]`

$$\begin{cases} x = \frac{1}{2} \left(c - \sqrt{c^2 - 4ab} \right) \\ y = \frac{1}{2} \left(\frac{\sqrt{c^2 - 4ab}}{a} - \frac{c}{a} \right) \\ z = \frac{c - \sqrt{c^2 - 4ab}}{2a} \end{cases}$$

$$\begin{cases} x = \frac{1}{2} \left(c + \sqrt{c^2 - 4ab} \right) \\ y = \frac{1}{2} \left(-\frac{\sqrt{c^2 - 4ab}}{a} - \frac{c}{a} \right) \\ z = \frac{c + \sqrt{c^2 - 4ab}}{2a} \end{cases}$$

Simplification

- Let a, b tend to zero

Simplification

- Let a, b tend to zero
- Cannot actually be zero (division by 0 at fixed points)

Simplification

- Let a, b tend to zero
- Cannot actually be zero (division by 0 at fixed points)

$$\{x = 0 \quad y = 0 \quad z = 0\}$$

Simplification

- Let a, b tend to zero
- Cannot actually be zero (division by 0 at fixed points)

$$\{x = 0 \quad y = 0 \quad z = 0\}$$

$$\left\{x = c \quad y = -\frac{c}{a} \quad z = \frac{c}{a}\right\}$$

Simplification

- Let a, b tend to zero
- Cannot actually be zero (division by 0 at fixed points)

$$\{x = 0 \quad y = 0 \quad z = 0\}$$

$$\left\{x = c \quad y = -\frac{c}{a} \quad z = \frac{c}{a}\right\}$$

- Much easier to analyze stability of system

Linearization

- Can take Jacobian of system to analyze stability

Linearization

- Can take Jacobian of system to analyze stability
- Which stabilities are hyperbolic in a 3D system?

Linearization

- Can take Jacobian of system to analyze stability
- Which stabilities are hyperbolic in a 3D system?

Definition

When $\lambda_1 \in \mathbb{R}$ and λ_2, λ_3 are complex conjugates *and* the real part of λ_1, λ_2 , and λ_3 all have the same sign, the fixed point is called a **Focus Node**

Linearization

- Can take Jacobian of system to analyze stability
- Which stabilities are hyperbolic in a 3D system?

Definition

When $\lambda_1 \in \mathbb{R}$ and λ_2, λ_3 are complex conjugates *and* the real part of λ_1, λ_2 , and λ_3 all have the same sign, the fixed point is called a **Focus Node**

Definition

When $\lambda_1 \in \mathbb{R}$ and λ_2, λ_3 are complex conjugates, then if the sign of λ_1 is the opposite of the sign of the real part of λ_2 and λ_3 , the fixed point is called a **Saddle Focus**

Linearization

- Can take Jacobian of system to analyze stability
- Which stabilities are hyperbolic in a 3D system?

Definition

When $\lambda_1 \in \mathbb{R}$ and λ_2, λ_3 are complex conjugates *and* the real part of λ_1, λ_2 , and λ_3 all have the same sign, the fixed point is called a **Focus Node**

Definition

When $\lambda_1 \in \mathbb{R}$ and λ_2, λ_3 are complex conjugates, then if the sign of λ_1 is the opposite of the sign of the real part of λ_2 and λ_3 , the fixed point is called a **Saddle Focus**

- Other stability exists, but are often named after the bifurcations that produce them

Linearization

- Jacobian of system will tell us about behavior of fixed points