

1 First Order Partial Differential Equations

Ex: Solve the following PDE

$$\begin{cases} yu_x + xu_y = 0 \\ u(0, y) = e^{-y^2} \end{cases}$$

Solution: This is a nice PDE; there are two ways to solve it, but we will focus on the easier one. We can see that we have a coefficient on both u_x and u_y . This tells us that when u was differentiated, there was a constant y and x . So now we can say that

$$\frac{dy}{dx} = \frac{x}{y}$$

This is a simple ODE to solve. We can use separation of variables:

$$\begin{aligned} xdx &= ydy \\ \int xdx &= \int ydy \\ \frac{1}{2}x^2 &= \frac{1}{2}y^2 + c \\ x^2 - y^2 &= c_0 \\ y^2 - x^2 &= c_1 \end{aligned}$$

You'll notice that we multiplied both sides by -1 . This is for convenience. Now we can express the answer as

$$u(x, y) = f(y^2 - x^2)$$

And we are done.

Ex: Solve the following PDE

$$u_x + u_y + u = x + y$$

Solution: In this case, we have a non-zero term on the right hand side of the equation. We will use substitution to solve this. For the time being, we will let T be $x + y$. To find our S , we can use a similar method to the one that we used above. We know that the coefficients on the u_x and u_y are 1 in this case. Now we have:

$$\begin{aligned}\frac{dy}{dx} &= 1 \\ dy &= dx \\ \int dy &= \int dx \\ x &= y + c \\ x - y &= c\end{aligned}$$

Now we can say that $S = x - y$. Next we have to actually use the substitution. We need to express u_x and u_y in terms of S and T . The following equations come from the chain rule; when the S and T terms are differentiated, we get u_S and u_T left behind.

$$\begin{aligned}u_x &= u_S \frac{\partial S}{\partial x} + u_T \frac{\partial T}{\partial x} & u_y &= u_S \frac{\partial S}{\partial y} + u_T \frac{\partial T}{\partial y} \\ u_x &= -u_S + u_T & u_y &= u_S + u_T\end{aligned}$$

Now we have our substitution for u_x and u_y . We can now plug in our values into the original equation. Note that we didn't do anything with u ; it gets left alone. In addition, if we needed u_{xx} or u_{yy} , we can just differentiate again. We will see this later when we talk about classifications. Now, back to plugging in:

$$\begin{aligned}(-u_S + u_T) + (u_S + u_T) + u &= T \\ 2u_T + u &= T\end{aligned}$$

This is just a first order ODE. Since it's not homogeneous, we need to solve for the general solution, and then the particular solution. Let's start with the general.

$$\begin{aligned}2y' + y &= 0 \\ 2r + 1 &= 0 \\ r &= -\frac{1}{2} \\ y &= ce^{(-\frac{1}{2})x}\end{aligned}$$

In this case, we know that our x is really T . So we finally have $y = ce^{(-\frac{1}{2})T}$. We will back substitute soon, but let's find the particular solution before we do that. Recall from ODE that we have several cases for the particular solution. In our case, we have a linear equation on the right hand side of our ODE. This lets us say:

$$\begin{aligned}y &= AT + B \\ y' &= A\end{aligned}$$

Now we can substitute again into our non-homogeneous ODE as follows:

$$2y' + y = T \equiv 2A + AT + B = T$$

And we have that $A = 1$, and $B = -2$. Now we have a final solution of:

$$\begin{aligned}y &= ce^{(-\frac{1}{2})T} + T - 2 \\ y &= ce^{(-\frac{1}{2})(x+y)} + (x + y) - 2\end{aligned}$$

We aren't quite done yet, as we have a solution to an ODE, not a PDE. It isn't too difficult to switch back; The c is really the only difference. In our ODE, the c is a constant term, but to our PDE, the c is a function. The c now becomes $f(s)$. We will go ahead and back substitute, which will give us $f(y-x)$. Now we have our final solution:

$$u(x, y) = f(y-x)e^{(-\frac{1}{2})(x+y)} + (x+y) - 2$$

And we are done.

2 Classification of Partial Differential Equations

We mentioned earlier the topic of classification of PDE's. Since we only have this and Fourier Series to talk about, we will go ahead and knock out classification. There's some information that we need to go ahead and put out here. This will be explained in the problem, but it's handy to have it for reference:

$$\begin{aligned}b^2 - ac < 0 &\implies \text{Elliptic with complex roots} \\b^2 - ac = 0 &\implies \text{Parabolic with the same root twice} \\b^2 - ac > 0 &\implies \text{Hyperbolic with two distinct real roots}\end{aligned}$$

Another helpful piece of information is what's known as the *classical trinity* of PDEs.

$$\begin{aligned}u_t - ku_{xx} &= 0 && \text{The heat equation (parabolic)} \\u_{tt} - c^2u_{xx} &= 0 && \text{The wave equation (hyperbolic)} \\u_{xx} + u_{yy} &= 0 && \text{The Laplace equation (elliptic)}\end{aligned}$$

Let's go ahead and look at an example.

Ex. Classify the following PDE:

$$yu_{xx} + u_{yy} = 0$$

Solution: The first thing we need to identify are the coefficients on the u_{xx} , u_{yy} , and u_{xy} or u_{yx} terms. Note that the u_{xy} and u_{yx} are interchangeable. We will assign these coefficients to a , b , and c corresponding to u_{xx} , u_{xy} , and u_{yy} , respectively. Looking at the PDE, we can see that

$$a = y \quad b = 0 \quad c = 1$$

Now we look at $b^2 - ac$. It should be noted that we are really dealing with $b = \frac{b_0}{2}$. In other words, we need to divide our b value by 2. In the case of the first problem, we are dealing with $b = 0$, so it's not evident. So, now we can go ahead and classify our PDE as follows:

$$b^2 - ac = 0 - y < 0 \implies \text{Elliptic}$$

Note that we assume $y > 0$. So our PDE is elliptic. Now we need to make it look like it's corresponding equation from the classical trinity. In this case we have an Elliptic equation, so we want to make it look like Laplace's equation. We need to find values for u_{xx} and u_{yy} , and then substitute those into the equation. This is similar to how we solved the first order PDE's; let's look at our characteristic equation:

$$\begin{aligned}am^2 + bm + c &= 0 \\ym^2 + 1 &= 0 \\m &= \pm \frac{i}{\sqrt{y}}\end{aligned}$$

So, now we have two solutions, and they are both imaginary as expected. Next we need to find values for S and T that we can substitute into the equation. We will do both of them at the same time here. Notice that we will drop the i ; we don't really care about the imaginary part:

$$\begin{aligned}\frac{dy}{dx} &= \frac{1}{\sqrt{y}} & \frac{dy}{dx} &= -\frac{1}{\sqrt{y}} \\ \int \frac{dy}{dx} &= \int \frac{1}{\sqrt{y}} & \int \frac{dy}{dx} &= -\int \frac{1}{\sqrt{y}} \\ S &= \frac{2}{3}y^{(\frac{3}{2})} - x & T &= \frac{2}{3}y^{(\frac{3}{2})} + x\end{aligned}$$

Now we can substitute into our equation. Unfortunately, we will need to take the partial derivative twice in order to get the substitutions we need. Let's start with u_{xx} .

$$\begin{aligned}u_x &= u_S \frac{\partial S}{\partial x} + u_T \frac{\partial T}{\partial x} \\&= -u_S + u_T \\u_{xx} &= \frac{\partial}{\partial x}(-u_S + u_T) \\&= u_{SS} + u_{TT}\end{aligned}$$

That wasn't too bad. We will skip the math for finding u_{yy} , which would give us

$$u_{yy} = y u_{SS} + y u_{TT}$$

Now we plug everything in. Remember that we were under the assumption that $y > 0$:

$$\begin{aligned}y(u_{SS} + u_{TT}) + y u_{SS} + y u_{TT} &= 0 \\2y u_{SS} + 2y u_{TT} &= 0 \\u_{SS} + u_{TT} &= 0\end{aligned}$$

And we are finished.

3 Fourier Series and Partial Differential Equations


It's finally time to talk about Fourier Series. Before we start though, it may be helpful to look at some common patterns amongst the Fourier Series. We have the following information that will help us:

	Cosine	Sine	Full Series
Interval	$[0, L]$	$[0, L]$	$[-L, L]$
Basis	$\{1, \cos(\frac{n\pi x}{L})\}$	$\{\sin(\frac{n\pi x}{L})\}$	$\{1, \sin(\frac{n\pi x}{L}), \cos(\frac{n\pi x}{L})\}$
Series	$\frac{b_0}{2} + \sum_{n=1}^{\infty} [b_n \cos(\frac{n\pi x}{L})]$	$\sum_{n=1}^{\infty} [a_n \sin(\frac{n\pi x}{L})]$	$\frac{b_0}{2} + \sum_{n=1}^{\infty} [a_n \sin(\frac{n\pi x}{L}) + b_n \cos(\frac{n\pi x}{L})]$

To solve for the coefficients, we have:

$$\begin{aligned}
 \text{Sine :} \quad a_n &= \frac{2}{L} \int_0^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx \\
 \text{Cosine :} \quad b_n &= \frac{2}{L} \int_0^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx \\
 &\quad b_0 = \frac{1}{L} \int_0^L f(x) dx \\
 \text{Full :} \quad a_n &= \frac{2}{L} \int_{-L}^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx \\
 &\quad b_n = \frac{2}{L} \int_{-L}^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx \\
 &\quad b_0 = \frac{1}{L} \int_{-L}^L f(x) dx
 \end{aligned}$$

It will also help us to quickly review the concept of odd and even functions, as they will help us to simplify the process of computing Fourier series later. Specifically, let's look at the graphs for sin and cos on $[-\pi, \pi]$, as well as another function that will turn out to be neither even nor odd..



We should recall that a function is *even* when $f(x) = f(-x)$ for all x in the domain of f . Graphically speaking, an *even* function is reflected across only the y -axis. By looking at the above graphs, we can see that $f(x) = \cos x$ is an even function.

A function is *odd* when $-f(x) = f(-x)$ for all x in the domain of f . Graphically speaking, the function is reflected across the x -axis and the y -axis. From the above graphs, we can say that $f(x) = \sin x$ is an odd function. Note that we can have a function that is neither odd nor even, but we won't see much of this in the following problems. The reason that we want to discuss odd and even functions here is because the integration that we will be seeing later will be affected by the symmetry of a function. Let's take a look at how even and odd functions effect integration:

Ex: Compute the following integral:

$$\int_0^{\pi} \cos x dx$$

Solution: Looking at this integral, we can see from the graph of the \cos above that this will be 0. That answer is meaningless to us however. What we are really looking for is

$$2 \int_0^{\frac{\pi}{2}} \cos x dx$$

We are allowed to do this since the \cos function is even. Essentially, we have turned the negative half of the cosine into a positive half. Similarly, we could ask for something like

$$\int_0^{\pi} |\cos x| dx$$

Since the absolute value of a function is not integrable, we drop the absolute value signs. This will leave us with our initial problem, which we will have to change anyways. Our integral becomes:

$$\int_0^{\pi} |\cos x| dx = 2 \int_0^{\frac{\pi}{2}} \cos x dx = 1$$

And we are done.

The first type of series we will look at is the Full Fourier Series. Note that this series is just a combination of a Fourier Sine Series and a Fourier Cosine Series. A typical problem would look something like this:

Ex: Find the corresponding Fourier Series for

$$f(x) = \begin{cases} L & -L \leq x \leq 0 \\ 2x & 0 \leq x \leq L \end{cases}$$

Solution: So we have a piecewise function here, but that shouldn't throw us off too much; all we need to do is combine the integrals for each part of the coefficient solution. Let's start by finding the terms for the *cos* series.

$$\begin{aligned} B_0 &= \frac{1}{2L} \left[\int_{-L}^0 L dx + \int_0^L 2x dx \right] \\ &= \frac{1}{2L} [2L^2] \\ &= L \end{aligned}$$

That wasn't too bad, was it? Now We need to find B_n , and then we will have our cosine part of the series. The integration can get pretty messy, so we will skip some steps.

$$\begin{aligned} B_n &= \frac{1}{L} \left[\int_{-L}^0 L \cos\left(\frac{n\pi x}{L}\right) dx + \int_0^L 2x \cos\left(\frac{n\pi x}{L}\right) dx \right] \\ &= \frac{1}{L} \left[0 + \left(\frac{2L^2}{n^2\pi^2} \right) ((-1)^n - 1) \right] \\ &= \frac{2L}{(n\pi)^2} ((-1)^n - 1) \end{aligned}$$

Note that the sequence will be 0 for all $n = 2k$. This allows us to substitute $2k + 1$ for n , since we don't care about the 0 terms. For our cosine part of the series, we now have:

$$\begin{aligned} B_n &= \frac{2L}{(2k+1)^2\pi^2} ((-1)^{2k+1} - 1) \\ &= \frac{2L}{(2k+1)^2\pi^2} (-2) \\ &= \frac{-4L}{(2k+1)^2\pi^2} \end{aligned}$$

So, now we need to find the sin part of the series. This is easier, since we only need to find A_n .

$$\begin{aligned} A_n &= \frac{1}{L} \left[\int_{-L}^0 L \sin\left(\frac{n\pi x}{L}\right) dx + \int_0^L 2x \sin\left(\frac{n\pi x}{L}\right) dx \right] \\ &= \frac{L}{n\pi} [-1 - (-1)^n] \end{aligned}$$

Note that we can do a similar substitution here as we did with the cosine series. This time, however, we will have 0 for $n = 2k + 1$. We now have:

$$\begin{aligned} A_n &= \frac{L}{2k\pi} [-1 - (-1)^{2k}] \\ &= \frac{L}{2k\pi} [-1 - 1] \\ &= \frac{L}{2k\pi} [-2] \\ &= \frac{-2L}{2k\pi} \end{aligned}$$

Now we have all the parts that we need. Our final answer will be essentially the sum of all the parts:

$$\begin{aligned} f(x) &\sim \frac{L}{2} + \sum_{k=0}^{\infty} \left[\frac{-4L}{(2k+1)^2\pi^2} \cos\left(\frac{(2k+1)\pi x}{L}\right) \right] + \sum_{k=0}^{\infty} \left[\frac{-2L}{2k\pi} \sin\left(\frac{2k\pi x}{L}\right) \right] \\ &\sim \frac{L}{2} + \sum_{k=0}^{\infty} \left[\frac{-4L}{(2k+1)^2\pi^2} \cos\left(\frac{(2k+1)\pi x}{L}\right) + \frac{-2L}{2k\pi} \sin\left(\frac{2k\pi x}{L}\right) \right] \end{aligned}$$

And we are done.

Now let's take a look at a slightly more theoretical Fourier Series.

Ex: Find the corresponding Fourier Series for

$$f(x) = |\cos x| \quad -\pi \leq x \leq \pi$$

Solution: Before we dive into calculations here, let's think about the function at hand. We have the absolute value of $\cos x$. Usually, \cos would be an odd function, which would imply that our \cos series component to the whole Fourier Series would be 0. However, the absolute value makes our function even, and as such the \sin component of the series will be 0. That's good news for us, because now we don't have to find a_n ; we actually only have a Fourier Cosine Series here. Recall that we have a formula for b_n and b_0 , and that our general solution will have the form of

$$\frac{b_0}{2} + \sum_{n=1}^{\infty} [b_n \cos\left(\frac{n\pi x}{L}\right)]$$


Our function is bounded above by π , so we really have $L = \pi$. This gives us:

$$\frac{b_0}{2} + \sum_{n=1}^{\infty} [b_n \cos(nx)]$$


Let's go ahead and find our b_0 :

$$\begin{aligned} b_0 &= \frac{1}{\pi} \int_{-\pi}^{\pi} |\cos x| dx \\ &= \frac{2}{\pi} \int_0^{\pi} |\cos x| dx \\ &= \frac{4}{\pi} \int_0^{\frac{\pi}{2}} |\cos x| dx \\ &= \frac{4}{\pi} \left[\sin \frac{\pi}{2} - \sin 0 \right] \\ &= \frac{4}{\pi} \end{aligned}$$

Let's take a second and discuss some the moves we made here. First of all, we know that since the function is even, we can say that the the integral of the entire range is really twice the integral of half the range. Since we are dealing with the cosine function, we actually have four times a quarter of the range. If we take a look at a graph, this may make a little more sense:



$f(x) = |\cos x|$ on $[0, 2\pi]$



$f(x) = \cos x$ on $[0, 2\pi]$

You can see that the section of the graph from $[\frac{\pi}{2}, \frac{3\pi}{2}]$ is positive for the $|\cos x|$, but negative for $\cos x$. If we simply integrated the $|\cos x|$ over $[0, \pi]$, the result would be 0, but graphically this result would not make any sense. Now let's take a look at the b_n term. Before we jump into the integration, let's think about this function:

$$|\cos x| = \begin{cases} \cos(x) & \text{if } 0 \leq x \leq \frac{\pi}{2} \\ -\cos(x) & \text{if } \frac{\pi}{2} \leq x \leq \pi \end{cases}$$

Now we can set up the formula for the b_n term. We have as follows:

$$\begin{aligned} b_n &= \frac{2}{\pi} \left[\int_0^{\frac{\pi}{2}} \cos x \cos nx dx - \int_{\frac{\pi}{2}}^{\pi} \cos x \cos nx dx \right] \\ &= \frac{2}{\pi} \left[\frac{-2 \cos\left(\frac{n\pi}{2}\right) + \sin(n\pi)}{n^2 - 1} \right] \end{aligned}$$

Note that (as we have seen before) that the sin term disappears; however, the more interesting part is what happens with our cos term. The value of $\cos\left(\frac{n\pi}{2}\right)$ will be 0 at all odd n , or that is to say for all $n = 2k + 1$. So, we will let our $n = 2k$. This gives us:

$$\begin{aligned} b_n &= \frac{2}{\pi} \left[\frac{-2 \cos(k\pi)}{(2k)^2 - 1} \right] \\ &= -\frac{4}{\pi} \left[\frac{(-1)^k}{(2k)^2 - 1} \right] \end{aligned}$$

And that leaves us with the final result

$$f(x) \sim \frac{2}{\pi} - \sum_{k=0}^{\infty} \left[\frac{4}{\pi} \left[\frac{(-1)^k}{(2k)^2 - 1} \right] \cos\left(\frac{2k\pi}{2}\right) \right]$$

Let's recall the 2nd order PDE that we found a series solution for earlier. Now we want to apply the initial condition of $u(x, 0) = x$.

Ex: Find the particular solution for the following PDE:

$$\begin{cases} \frac{\partial u}{\partial t} = k \frac{\partial^2 u}{\partial x^2} \\ u(x, 0) = x \\ u(0, t) = u(L, t) = 0 \end{cases}$$

Solution: Earlier, we found a general solution of the form

$$\begin{aligned} u_n(x, t) &= a_n \sin\left(\frac{n\pi x}{L}\right) e^{-k\left(\frac{n\pi}{L}\right)^2 t} \\ u(x, t) &= \sum_{n=1}^{\infty} \left[a_n \sin\left(\frac{n\pi x}{L}\right) e^{-k\left(\frac{n\pi}{L}\right)^2 t} \right] \end{aligned}$$

Without loss of generality, let's let $L = \pi$. This would change our boundary condition to $u(\pi, t) = 0$, but this wouldn't change the solution that we have here in any way other than replacing the L 's in the equation with π . We will also let $k = 1$. Again this doesn't change anything other than the fact that we will replace k with 1. We now have:

$$u(x, t) = \sum_{n=1}^{\infty} \left[a_n \sin(nx) e^{-(n)^2 t} \right]$$

This is a much nicer sequence for us to deal with. Our goal now is to find a_n , which will give us a particular solution. We know that our answer for the particular solution will be in the form of a series, since our initial condition is not represented in the form of terms of a series (we will see an example of this later). Let's go ahead and apply our initial condition. Doing so will yield:

$$x = \sum_{n=0}^{\infty} [a_n \sin nx]$$

Now, we need to come up with an a_n . We have a Fourier Sine Series here, so we can use the formula for finding a_n that corresponds to our case.

$$\begin{aligned} a_n &= \frac{2}{\pi} \int_0^{\pi} x \sin(nx) dx \\ &= -\frac{2n\pi \cos(n\pi) + 2 \sin(n\pi)}{n^2 \pi} \\ &= -\frac{2(-1)^n}{n} \end{aligned}$$

So our particular solution is:

$$u(x, t) = \sum_{n=1}^{\infty} \left[-\frac{2(-1)^n}{n} \sin(nx) e^{-(n)^2 t} \right]$$

And we are done. We haven't discussed the convergence of series yet, and it turns out that if the series converges, it will converge to the solution. We won't worry about it right now. Instead, let's look at another example

Ex: Solve the following boundary value problem given the general solution:

$$\begin{cases} u_{tt} - u_{xx} = 0 \\ u(0, t) = 0 & u(1, t) = 0 \\ u(x, 0) = f(x) & u_t(x, 0) = g(x) \end{cases}$$

$$f(x) = 2 \sin(\pi x) + 3 \sin(2\pi x) \quad g(x) = 4 \sin(3\pi x) - 7 \sin(5\pi x)$$

$$u(x, t) = \sum_{n=1}^{\infty} [(a_n \cos[n\pi t] + b_n \sin[n\pi t]) \sin[n\pi x]]$$

Solution: There's a lot going on here, so let's break it down. First of all, notice that we have two initial conditions, one of which is a derivative. This should make sense since the equation have two second order partial derivatives included in it. Next notice that we have a Fourier Sine Series, despite both the $\sin[n\pi t]$ and $\cos[n\pi t]$ terms. If we had worked out the entire problem, we would see that our eigenvalue/function for this equation takes the form of

$$\{\lambda_n = n\pi, \sin(n\pi x)\}$$

which tells us that the corresponding series is a Fourier Sine Series. We also see the initial conditions on our equation are already representative of a series of sine functions, which is not just a coincidence. To apply the initial conditions, we will need both the general solution and the derivative with respect to t of the general solution. Let's go ahead and apply the first initial condition, $u(x, 0) = f(x)$. Letting $t = 0$ gives us:

$$u(x, 0) = 2 \sin(\pi x) + 3 \sin(2\pi x) = \sum_{n=1}^{\infty} [a_n \sin(n\pi x)]$$

We see that the term with b_n has dropped out, since $\sin(0) = 0$. Now we just have to match up the coefficients on a_n . We can see that when we have $n = 1$, $a_n = 2$, and when $n = 2$, $a_n = 3$. For all other n , we have $a_n = 0$, so we ignore those terms. Next, we can apply the initial condition for the derivative of the general solution with respect to t . We expect that all the terms related to a_n will disappear, and in fact they will. Let's take a look:

$$u_t(x, t) = \sum_{n=1}^{\infty} [(-a_n n\pi \sin[n\pi t] + b_n n\pi \cos[n\pi t]) \sin[n\pi x]]$$

$$u_t(x, 0) = 4 \sin(3\pi x) - 7 \sin(5\pi x) = \sum_{n=1}^{\infty} [n\pi b_n \sin(n\pi x)]$$

We are going to do the same thing here that we did to find our a_n terms, but notice that we don't just have b_n ; we actually have $n\pi b_n$. So we have the following:

$$\begin{aligned} 3\pi b_3 &= 4 & 5\pi b_5 &= -7 \\ b_3 &= \frac{4}{3\pi} & b_5 &= -\frac{7}{5\pi} \end{aligned}$$

Much like the solutions for a_n , all the other b_n terms are 0. Now we can construct a particular solution. In this case, the answer will not be in the form of a series, since we have definite terms. Our solution is as follows:

$$u(x, t) = 2 \cos(\pi t) \sin(\pi x) + 3 \cos(2\pi t) \sin(2\pi x) + \frac{4}{3\pi} \sin(3\pi t) \sin(3\pi x) - \frac{7}{5\pi} \sin(5\pi t) \sin(5\pi x)$$

And we are done.

Up to this point, we have found a Fourier series and left our solutions at that. We will now look into the convergence of a Fourier series. This topic includes two questions:

1. Does the series converge?
2. If the series converges, to what value does it converge?

Both of these questions are highly theoretical. We will discuss the answers to them, but first we need to go over some terms:

1. ***Jump Discontinuity:*** To say that $f(x)$ has a jump discontinuity at $x = a$ is to say that

$$\lim_{x \rightarrow a^-} f(x) \neq \lim_{x \rightarrow a^+} f(x)$$

Where both the limit from the left and the limit from the right exist.

2. ***Piecewise Smooth:*** A function $f(x)$ is said to be piecewise smooth if the function can be broken into two distinct intervals, and on each interval both $f(x)$ and $f'(x)$ are continuous.
3. ***Periodic Extension:*** The periodic extension of $f(x)$ on $[-L, L]$ is the repetition of the function on intervals on periods to the left and right of $[-L, L]$.

With these definitions, we can lay out a general rule for convergence of a Fourier series. We won't give a proof of these rules for the time being. We just need them to lay out some groundwork for the rest of the information we will present on Fourier series. We have the following:

Suppose $f(x)$ is a function, and that $f(x)$ is piecewise smooth on the interval $[-L, L]$. The corresponding Fourier series for $f(x)$ converges to either

1. the periodic extension of $f(x)$, if $f(x)$ is continuous, or
2. the average of the two one-sided limits, if the periodic extension of $f(x)$ has a jump discontinuity at $x = a$.

We should note that the second condition can be expressed by the formula

$$\frac{1}{2} \left[\lim_{x \rightarrow a^-} f(x) + \lim_{x \rightarrow a^+} f(x) \right]$$

which gives us a nice way to calculate the value to which the series converges if we happen to meet the criteria in (2). Let's take a look at some examples.


Ex: Find the Fourier series for the following equation, and determine if the series converges. If it does, tell where the series converges to.

$$f(x) = \begin{cases} L & -L \leq x \leq 0 \\ 2x & 0 \leq x \leq L \end{cases}$$

Solution: We will go ahead and skip the details of finding the Fourier series for this function. If you do want to find it, you will have to find a full piecewise Fourier series. For this particular $f(x)$, we have:

$$f(x) \sim L + \sum_{n=1}^{\infty} \left[\frac{2L}{n^2\pi^2} ([-1]^n - 1) \cos\left(\frac{n\pi x}{L}\right) \right] - \sum_{n=1}^{\infty} \left[\frac{L}{n\pi} (1 + [-1]^n) \sin\left(\frac{n\pi x}{L}\right) \right]$$

You'll notice that we didn't simplify this solution to a nicer form. If we evaluate this series, we will run across terms where the series is 0, but that is okay for now. You should also notice that this function is defined as a piecewise function, and on both intervals $([-L, 0])$ and $[0, L])$, $f(x)$ is continuous. In other words, $f(x)$ is piecewise smooth, so we know that this series converges. Now let's take a look at the graph of $f(x)$ on an arbitrary interval of $[-1, 1]$.



$$f(x) = \begin{cases} 1 & -1 \leq x \leq 0 \\ 2x & 0 \leq x \leq 1 \end{cases}$$

We can see that there is a jump in the graph at $x = 0$. In other words, there is a jump discontinuity at $x = 0$, and that means that we need to use (2) in order to find the value of convergence. We will let \sum stand in for our entire series. We have:

$$\begin{aligned} \sum &\rightarrow \frac{1}{2} \left[\lim_{x \rightarrow 0^-} L + \lim_{x \rightarrow 0^+} 2x \right] \\ &\rightarrow \frac{1}{2} [L + 0] \\ &\rightarrow \frac{L}{2} \end{aligned}$$

So, at $x = 0$, the series converges to $\frac{L}{2}$. We are not done yet, however. Now, we need to consider the periodic extension of the series. The function will repeat every $[-L, L]$, so we need to look at what the convergence of the series looks like at $-L$ and L . Let's take a look:

$$\begin{aligned} \sum &\rightarrow \frac{1}{2} \left[\lim_{x \rightarrow -L^-} f(x) + \lim_{x \rightarrow -L^+} f(x) \right] & \sum &\rightarrow \frac{1}{2} \left[\lim_{x \rightarrow L^-} f(x) + \lim_{x \rightarrow L^+} f(x) \right] \\ &\rightarrow \frac{1}{2} [2L + L] & &\rightarrow \frac{1}{2} [2L + L] \\ &\rightarrow \frac{3L}{2} & &\rightarrow \frac{3L}{2} \end{aligned}$$

And we are done.

4 Second Order Partial Differential Equations

4.1 The Heat and Wave Equations

Let's take a look at the second order PDE. We will start with a homogeneous equation, which we will tackle with the separation of variables method. The general strategy is that we want to take our PDE and convert it to an ODE, which we know how to solve. The problems presented here also involve boundary conditions, which will require us to solve for eigenvalues/functions. We will also be dealing with series (specifically Fourier Series) to help us solve these problems. Let's start by looking at the homogeneous heat equation.

Ex: Solve the following PDE

$$\begin{cases} \frac{\partial u}{\partial t} = k \frac{\partial^2 u}{\partial x^2} \\ u(x, 0) = x \\ u(0, t) = u(L, t) = 0 \end{cases}$$

Solution: We will use the separation of variable technique to solve this problem. We know to use separation of variable because we are dealing with a homogeneous PDE with homogenous boundary conditions. Separation of variables tells us that our solution will be in the form of $u(x, t) = X(x)T(t)$. If we take this fact at face value, we can use the separation of variables method to solve this. First, we substitute in our solution into the PDE:

$$\begin{aligned} \frac{\partial}{\partial t}(X(x)T(t)) &= k \frac{\partial^2}{\partial x^2}(X(x)T(t)) \\ X(x) \frac{dT}{dt} &= k \frac{d^2 X}{dx^2} T(t) \\ \frac{dT}{dt} \frac{1}{T(t)} &= \frac{d^2 X}{dx^2} \frac{k}{X(x)} \end{aligned}$$

The only way that our function of x and our function of t can be equal is if they are equal to a constant; thus we now have:

$$\frac{dT}{dt} \frac{1}{T(t)} = \frac{d^2 X}{dx^2} \frac{k}{X(x)} = -\lambda$$

And now we have a first order ODE for the time and a second order ODE for the position:

$$\frac{dT}{dt} = -k\lambda T(t) \quad \text{and} \quad \frac{d^2 X}{dx^2} = -\lambda X(x)$$

Our next step is to find the Eigenvalues and Eigenfunctions of the second order ODE; we will come back to the time ODE when we have values for λ .

To solve for these values, we have to find non-trivial eigenvalues that correspond to the boundary values of the ODE. In this case, we know that $X(0) = 0$ and $X(L) = 0$. These boundary conditions come from the BC on the PDE which states $u(0, t) = u(L, t) = 0$. To find the eigenvalues/functions, we need to find the characteristic equation corresponding to the ODE in question. Let's go ahead and find it:

$$\begin{aligned} \frac{d^2 X}{dx^2} &= -\lambda \frac{X(x)}{k} \\ \frac{d^2 X}{dx^2} + \lambda \frac{X(x)}{k} &= 0 \\ r^2 + \lambda &= 0 \\ r^2 &= -\lambda \\ r &= \pm \sqrt{-\lambda} \end{aligned}$$

Now, we can substitute values of λ to find out solutions. Let's start with $\lambda > 0$. Since we have a $-\lambda$ under the radical, we will have $r = \pm \sqrt{\lambda}i$. This will yield the solution of

$$y(x) = c_1 \cos \sqrt{\lambda}x + c_2 \sin \sqrt{\lambda}x$$

Applying our boundary conditions gives us:

$$y(0) = c_1 \cos 0 + c_2 \sin 0 = 0$$

Now we know that $c_1 = 0$. This result came from the fact that $\sin 0 = 0$, so the c_2 term disappears. Now we are left with $\cos 0 = 1$, which tells us that $c_1 = 0$. We can apply this to our second boundary condition like so:

$$y(L) = 0 = c_2 \sin \sqrt{\lambda}L$$

It looks like there will only be trivial solutions, but we know this isn't true due to the Jacobian matrix. Recall that if the determinant of the Jacobian $\neq 0$, then we only have trivial solutions, and if the determinant of the Jacobian $= 0$, then we do not have the trivial solution. Let's take a look at the Jacobian for this case:

$$\begin{aligned} \begin{vmatrix} 1 & 0 \\ \cos \sqrt{\lambda}L & \sin \sqrt{\lambda}L \end{vmatrix} &= 0 \\ \sin \sqrt{\lambda}L &= 0 \\ \sqrt{\lambda}L &= \arcsin 0 \\ \sqrt{\lambda} &= \frac{\arcsin 0}{L} \\ \lambda &= \left(\frac{\arcsin 0}{L} \right)^2 \\ \lambda &= \left(\frac{n\pi}{L} \right)^2 \end{aligned}$$

Notice that we replaced the $\arcsin 0$ term with $n\pi$. Recall from trig that the sin function is 0 at every $n\pi$, starting with $n = 0$. Next, we just need to find our corresponding eigenfunction. Our c_1 was 0, so we can ignore the entire cos term. Thus, we have an eigenfunction from plugging in our λ that looks like:

$$X(x) = \sin \frac{n\pi x}{L}$$

Unfortunately, we aren't done yet. Now we have to find the corresponding eigenvalues/functions for $\lambda = 0$. This isn't too hard, and can be done in a few lines if we recall the characteristic equation. Plugging in 0 for λ gives us that $r = 0$. Using this piece of information, we now know that $y(x) = c_1 + xc_2$. Applying our boundary conditions gives us $c_1 = 0, c_2 = 0$. I'll leave you to do the math (try plugging into the Jacobian; you'll see that there is no way to force the determinant to be 0), but we can see that we have only trivial solutions for $\lambda = 0$.

Our final case is for $\lambda < 0$. I'll go ahead and tell you that this will yield another trivial solution, but we can take a look at the work behind it. Since $\lambda > 0$, we know that our r from the characteristic equation will be $r = \pm\sqrt{\lambda}$. From here we are going to do something a little unconventional that will help us. Usually, we would have a solution of the form $y(x) = c_1 e^{\sqrt{\lambda}x} + c_2 x e^{-\sqrt{\lambda}x}$. In our case, it will be more convenient to use hyperbolic trig functions. It's not too far off from our solution from $\lambda > 0$:

$$\begin{aligned} y(x) &= c_1 e^{\sqrt{\lambda}x} + c_2 e^{-\sqrt{\lambda}x} \\ &= c_1 \cosh \sqrt{\lambda}x + c_2 \sinh \sqrt{\lambda}x \end{aligned}$$

I'll leave you to apply the boundary conditions (remember that $\sinh 0 = 0$ and $\cosh 0 = 1$), but you will find that you have only trivial solutions.

So, we are done with the second order ODE, and now we turn our eye to the first order time equation. We know our λ , so it isn't too hard to solve:

$$\begin{aligned} \frac{dT}{dt} &= -\lambda T(t) \\ T(t) &= ce^{-k\left(\frac{n\pi}{L}\right)^2 t} \end{aligned}$$

Finally we can put it all together. Our penultimate solution will be

$$u_n(x, t) = B_n \sin\left(\frac{n\pi x}{L}\right) e^{-k\left(\frac{n\pi}{L}\right)^2 t}$$

We can analyze this solution and tell where each piece comes from. The B_n is a coefficient that results from our c' 's in the general solutions of the differential equations. The *sin* part results from our eigen vector. Remember, we get a different value for each n , so there may or may not be 0 terms in the series. Finally, we have the exponential term. This term comes from the solution to our time equation. When we put it all together, we can see that this is, of course a series; for every n , we get a different solution. From here we can find the Fourier Series (which will be a Fourier Sine Series). We won't worry about that yet, but we will come back to it.

It will benefit us to take a moment and discuss how boundary conditions effect our Fourier series. We have three different combinations of boundary conditions, where each one will produce a different result in our series solution.

1. *Dirichlet Boundary Conditions*

Dirichlet boundary conditions involve the function for which we are looking. In other words, the boundary conditions are presented in terms of the functions we are trying to find, rather than the derivative of that function. Typically, this type of boundary condition will take the form of (in the homogeneous case)

$$u(0, t) = u(L, t) = 0.$$

In regards to our Fourier series, we expect to see $\sin\left(\frac{n\pi x}{L}\right)$. Our basis will look like

$$\left\{ \lambda_n = \left(\frac{n\pi}{2}\right)^2, \phi_n(x) = \sin\left(\frac{n\pi x}{L}\right) \right\}$$

2. *Neumann Boundary Conditions*

Neumann boundary conditions involve the derivatives of the function for which we are solving. Neumann boundary conditions will look like

$$u_x(0, t) = u_x(L, t) = 0.$$

This will cause our Fourier series to contain $\cos\left(\frac{n\pi x}{L}\right)$. Our basis will look like

$$\left\{ \lambda_n = \left(\frac{n\pi}{2}\right)^2, \phi_n(x) = \cos\left(\frac{n\pi x}{L}\right) \right\}$$

3. *Robin Boundary Conditions*

Robin boundary conditions will contain a mixture of the function and it's derivative. Typically, we will see

$$u_x(0, t) = u(L, t) = 0 \quad \text{or} \quad u(0, t) = u_x(L, 0) = 0.$$

These boundary conditions will give us two different results, depending on whether we have the derivative first or second. We will have

$$\begin{aligned} u_x(0, t) = u(L, t) = 0 &\implies \left\{ \lambda_n = \left(\left[n + \frac{1}{2}\right] \frac{\pi}{L}\right)^2, \phi_n(x) = \cos\left(\left[n + \frac{1}{2}\right] \frac{\pi}{L} x\right) \right\} \\ u(0, t) = u_x(L, t) = 0 &\implies \left\{ \lambda_n = \left(\left[n + \frac{1}{2}\right] \frac{\pi}{L}\right)^2, \phi_n(x) = \sin\left(\left[n + \frac{1}{2}\right] \frac{\pi}{L} x\right) \right\} \end{aligned}$$

Now we can take a look at a slightly harder PDE. We still have the heat equation, but we will not have homogeneous boundary conditions. Our goal here is to find a way to transform our boundary conditions into a homogenous form and take it from there.

Ex: Solve the following PDE

$$\begin{cases} u_t - u_{xx} = 0 \\ u(0, t) = 0, \quad u(1, t) = t \\ u(x, 0) = x^2 \end{cases}$$

Solution: We would like to just jump into this PDE and solve it, but it's not quite that simple. We have to find a way to express our boundary conditions in a homogeneous manner. We can do this through a clever substitution; we will create a new PDE, denoted Υ . To find Υ , we use the following formula:

$$\Upsilon = B_1 + \frac{x}{l} \Delta B,$$

where B is a boundary condition. In our case, we have as follows:

$$\begin{aligned} \Upsilon &= B_1 + \frac{x}{l} \Delta B \\ &= 0 + \frac{x}{1}(t - 0) \\ &= xt \end{aligned}$$

Now, we find a new PDE, denoted by v . We have another formula that tells us

$$v(x, t) = u(x, t) - \Upsilon(x, t)$$

Applying this formula gives us

$$\begin{aligned} v(x, t) &= u(x, t) - \Upsilon \\ &= u(x, t) - xt \end{aligned}$$

Let's take a moment to analyze the result. We wanted to find a new PDE in terms of our old one that would allow us to have homogeneous boundary conditions. We found a solution, called v , in terms of our original PDE, called u . Our original PDE was $u_t - u_{xx} = 0$, so to properly perform our substitution, we need to find v_t and v_{xx} . Through some careful differentiation, we have

$$v_t = u_t - x \quad \text{and} \quad v_{xx} = u_{xx}$$

To find our new PDE, we simply evaluate.

$$v_t - v_{xx} = u_t - x - u_{xx} = u_t - u_{xx} - x = 0 - x = -x$$

So we finally have a new PDE that looks like

$$\begin{cases} v_t - v_{xx} = -x \\ v(0, t) = v(1, t) = 0 \\ v(x, 0) = x^2 \end{cases}$$

You may wonder how this has made anything simpler, since the PDE itself is no longer homogeneous. By forcing the boundary conditions to be homogeneous, we can now use a Fourier series to find our solution. Recall that earlier we discussed the boundary condition and their resulting eigenvalue/eigenfunction. Here, we have Dirichlet boundary conditions, which tells us that we have a Fourier sine series. We have that

$$\begin{aligned} v(x, t) &= \sum_{n=1}^{\infty} [a_n(t) \sin(n\pi x)] \\ v_t(x, t) &= \sum_{n=1}^{\infty} [a'_n(t) \sin(n\pi x)] \\ v_{xx}(x, t) &= - \sum_{n=1}^{\infty} [a_n(t)(n\pi)^2 \sin(n\pi x)] \end{aligned}$$

Which tells us that

$$v_t - v_{xx} = \sum_{n=1}^{\infty} [(a'_n(t) + (n\pi)^2 a_n(t)) \sin(n\pi x)] = -x$$

This may look rather complicated, but all we have is a function's Fourier sine series. Essentially, we have

$$-x = \sum_{n=1}^{\infty} [b_n \sin(n\pi x)]$$

To find b_n , we use the formula that was discussed in the earlier section about Fourier series. We have that

$$\begin{aligned} a'_n(t) + (n\pi)^2 a_n(t) &= \int_0^1 -x \sin(n\pi x) dx \\ &= \left. \frac{2x}{n\pi} \cos(n\pi x) \right|_0^1 \\ &= \frac{2}{n\pi} \end{aligned}$$

We now have an ODE which we can solve for a_n . Note that from our PDE, we do have an initial condition for the ODE, but it is not as simple as just saying $a_n(0) = x^2$. At $t = 0$, we have the following series:

$$\sum_{n=1}^{\infty} a_n(0) \sin(n\pi x)$$

We got this by taking our original $v(x, t)$ and plugging in 0 for t . We know how to get $a_n(0)$, since this is just a simple Fourier sine series. It is given by

$$\begin{aligned} a_n(0) &= 2 \int_0^1 x^2 \sin(n\pi x) dx \\ &= \frac{(-1)^{(n+1)}}{n\pi} + \frac{2}{(n\pi)^3} [(-1)^n - 1] \end{aligned}$$

We now have all we need to solve the ODE. We will omit the work for the solution, but the final answer for a_n is

$$a_n(t) = \frac{4[(-1)^n - 1]}{(n\pi)^3} e^{-(n\pi)^2 t} + \frac{2(-1)^n}{(n\pi)^3}$$

Now all that's left to do is go back to u from v . This may seem hard, but we saw earlier that $v = u - \Upsilon$. Solving for u gives us $u = v + \Upsilon$. Recall that Υ was xt , and v was the series for which we just found a_n . Putting it all together gives us

$$u(x, t) = xt + \sum_{n=1}^{\infty} \left[\frac{4[(-1)^n - 1]}{(n\pi)^3} e^{-(n\pi)^2 t} + \frac{2(-1)^n}{(n\pi)^3} \right] \sin(n\pi x)$$

And we are done.

Ex: Solve the following PDE

$$\begin{cases} u_{tt} - u_{xx} = 1 + t \cos(2\pi x) \\ u_x(0, t) = u_x(1, t) = 0 \\ u(x, 0) = 2 \cos(2\pi x), \quad u_t(x, 0) = 1 + \cos(\pi x) - \cos(2\pi x) \end{cases}$$

Solution: This problem is an example of the wave equation with a non-homogeneous solution. We are lucky this time; the boundary conditions are already homogeneous, so there is no need to change our variable. We know that since both of the boundary conditions are derivatives, we are dealing with a Fourier cosine series. This yields the eigenvalue/function of

$$\{\lambda_n = (n\pi)^2, \cos(n\pi x)\}$$

Since we are dealing with a cosine series, we also have a leading term, a_0 . Our full series looks like

$$u(x, t) = a_0(t) + \sum_{n=1}^{\infty} a_n(t) \cos(n\pi x)$$

We need u_{tt} and u_{xx} , so we have:

$$u_{tt}(x, t) = a_0''(t) + \sum_{n=1}^{\infty} a_n''(t) \cos(n\pi x) \quad \text{and} \quad u_{xx}(x, t) = - \sum_{n=1}^{\infty} (n\pi)^2 a_n(t) \cos(n\pi x)$$

All we have to do now is simply plug into our original equation. This gives us

$$\begin{aligned} a_0''(t) + \sum_{n=1}^{\infty} a_n''(t) \cos(n\pi x) + \sum_{n=1}^{\infty} (n\pi)^2 a_n(t) \cos(n\pi x) &= 1 + t \cos(2\pi x) \\ a_0''(t) + \sum_{n=1}^{\infty} (a_n''(t) + (n\pi)^2 a_n(t)) \cos(n\pi x) &= 1 + t \cos(2\pi x) \end{aligned}$$

Now we can play the matching game. We have a *constant + series term* on the right side, and *constant + series* on the left, so we can match up the constants. This gives us

$$a_0''(t) = 1$$

Now we match up the series term. On the right hand side, we have $\cos(2\pi x)$, so we know that for $n = 2$, we have

$$a_2''(t) + 4\pi^2 a_2(t) = t$$

a_0 and a_2 are the only terms that appear, so we can assume for now that $\forall n \neq 2$ and $3, a_n = 0$. We are left with two ODE's, so we now need to find the corresponding initial conditions. Let's apply the first initial condition to our PDE. This gives us

$$a_0(0) + \sum_{n=1}^{\infty} a_n(0) \cos(n\pi x) = 2 \cos(2\pi x)$$

Which tells us

$$a_0 = 0 \qquad a_2 = 2$$

Now we can apply the second initial condition. After taking the partial derivative of u with respect to t , we have

$$a_0'(0) + \sum_{n=1}^{\infty} a_n'(0) \cos(n\pi x) = 1 + \cos(\pi x) - \cos(2\pi x)$$

Some more clever pattern matching give us

$$a_0'(0) = 1 \qquad a_1'(0) = 1 \qquad a_2'(0) = -1$$

Notice that we have a term a_1' . We cannot simply ignore this value; rather we have to take it into account when we find our solution. We know that we have an ODE for $a_1(t)$, but it may not be evident how to express it. Recall that earlier we said $\forall n \neq 2 \text{ and } 3, a_n = 0$. This means that $a_1 = 0$, and since a_1 is a coefficient in the Fourier series, we know that $a_1''(t) + \pi^2 a_1(t) = 0$. We now have three different ODE's that we must solve:

$$\begin{cases} a_0''(t) = 1 \\ a_0'(0) = 1, \quad a_0(0) = 1 \end{cases} \quad \begin{cases} a_1''(t) + \pi^2 a_1(t) = 0 \\ a_1'(0) = 1, \quad a_1(0) = 0 \end{cases} \quad \begin{cases} a_2''(t) + 4\pi^2 a_2(t) = t \\ a_2'(0) = 1, \quad a_2(0) = 2 \end{cases}$$


Luckily, Mathematica is here to save the day. We will skip the work and let the computer do the thinking for us here. We get that

$$\begin{aligned} a_0(t) &= \frac{1}{2}t^2 + t \\ a_1(t) &= \frac{\pi \cos(t\pi) + \sin(t\pi)}{\pi} \\ a_2(t) &= \frac{2\pi t + 16\pi^3 \cos(2\pi t) - \sin(2\pi t) + 4\pi^2 \sin(2\pi t)}{8\pi^3} \end{aligned}$$

Now all we need to do is plug back in to our u . Our final answer becomes

$$u(x, t) = \frac{1}{2}t^2 + t + \frac{\pi \cos(t\pi) + \sin(t\pi)}{\pi} \cos(\pi x) + \frac{2\pi t + 16\pi^3 \cos(2\pi t) - \sin(2\pi t) + 4\pi^2 \sin(2\pi t)}{8\pi^3} \cos(2\pi x)$$

Just for interest, the plot of this solution looks like



On $-\pi \leq \theta \leq \pi$ and $-\pi \leq x \leq \pi$; And we are done.

4.2 The Laplace Equation

Up to this point, we have only looked at the wave equation and the heat equation. There is, however, one other equation in the classical trinity: the Laplace Equation. We haven't dealt with this equation yet because we have to be very careful in our approach to solving it. When we looked at the homogeneous wave and heat equations, we used the general approach called *Separation of Variables*. In the case of the homogeneous Laplace equation, we may not be able to tackle the problem using separation of variables. The problem arises from the various types of boundaries that effect the domain. We can either have a 'rectangular' bound, which will be separable, or a 'circular' bound which is **not** separable. Imagine that we have the following two problems:

$$A = \begin{cases} u_{xx} + u_{yy} = 0 & \text{on } [0, 1] \times [0, 2] \\ u(x, 0) = u(x, 2) = 0 \\ u(0, y) = y, u(1, y) = 0 \end{cases} \quad B = \begin{cases} u_{xx} + u_{yy} = 0 & \text{on } x^2 + y^2 < a^2 \\ u(x, y) \big|_{x^2 + y^2 = a^2} = f(x, y) \end{cases}$$

A is an example of a separable Laplace equation. We can see graphically that the domain $[0, 1] \times [0, 2]$ is rectangular. B 's domain is not separable, since it is circular. Let's take a look A .

Ex: Solve the following PDE

$$\begin{cases} u_{xx} + u_{yy} = 0 & \text{on } [0, 1] \times [0, 2] \\ u(x, 0) = u(x, 2) = 0 \\ u(0, y) = y, u(1, y) = 0 \end{cases}$$

Solution: Earlier, we discussed the different boundary conditions we can encounter and their corresponding eigenvalues/functions. In this problem, we have Dirichlet boundary conditions. This tells us that

$$\left\{ \lambda_n = \left(\frac{n\pi}{2} \right)^2, \phi_n(y) = \sin \left(\frac{n\pi y}{2} \right) \right\}$$

Notice that we used y here instead of x . We did this because the boundary conditions are defined in terms of y , not x . Now we can continue to solve our problem. Recall that our separation of variables gives us two second order ODE's,

$$X'' - \lambda X = 0 \quad \text{and} \quad \lambda Y - Y'' = 0$$

We already solved for $Y(y)$ by finding the corresponding eigenvalues/vectors. Now we plug in our λ_n into our X ODE. Solving the resulting ODE will give us a basis of

$$\left\{ e^{\frac{n\pi}{2}x}, e^{-\frac{n\pi}{2}x} \right\}$$

Now we could continue at this point, but if we think ahead, we can see that when we eventually apply our initial conditions, nothing will cancel out. At this point, we want to transform our basis. We have as follows

$$\left\{ e^{\frac{n\pi}{2}x}, e^{-\frac{n\pi}{2}x} \right\} \implies \left\{ \cosh \left(\frac{n\pi x}{2} \right), \sinh \left(\frac{n\pi x}{2} \right) \right\}$$

All we did here is apply an identity. Now we have everything we need to set up a solution. Our $u(x, y)$ will be

$$u(x, y) = \sum_{n=1}^{\infty} \left[a_n \cosh \left(\frac{n\pi x}{2} \right) + b_n \sinh \left(\frac{n\pi x}{2} \right) \right] \sin \left(\frac{n\pi y}{2} \right)$$

We can apply our initial conditions, which gives us b_n and a_n :

$$a_n = \frac{4(-1)^{n+1}}{n\pi}$$

$$b_n = -\coth \left(\frac{n\pi}{2} \right) \frac{4(-1)^{n+1}}{n\pi}$$

This yields a final solution of

$$u(x, y) = \sum_{n=1}^{\infty} \left[\frac{4(-1)^{n+1}}{n\pi} \cosh \left(\frac{n\pi x}{2} \right) - \coth \left(\frac{n\pi}{2} \right) \frac{4(-1)^{n+1}}{n\pi} \sinh \left(\frac{n\pi x}{2} \right) \right] \sin \left(\frac{n\pi y}{2} \right)$$

And we are done

Now we turn our attention to the Laplace Equation with a circular domain. As we discussed earlier, we can't simply say that our solution will be in the form of $X(x)Y(y)$. However, we can force the domain to be rectangular if we use polar coordinates. Let's consider a variation of B from above.

Ex: Solve the following PDE

$$\begin{cases} u_{xx} + u_{yy} = 0 & \text{on } x^2 + y^2 < a^2 \\ u(a, \theta) = U_1 & \text{on } 0 \leq \theta \leq \pi \\ u(a, \theta) = U_2 & \text{on } \pi \leq \theta \leq 2\pi \end{cases}$$

Solution: We want to express the partial derivatives here in terms of r and θ . Recall that we have

$$x = r \cos \theta \quad \text{and} \quad y = r \sin \theta$$

First, we will find the partial derivatives of x and y with respect to r and θ . Since $x = r \cos \theta$,

$$\frac{\partial x}{\partial r} = \cos \theta \quad \frac{\partial x}{\partial \theta} = -r \sin \theta$$

We can compute the derivatives for y in a similar fashion:

$$\frac{\partial y}{\partial r} = \sin \theta \quad \frac{\partial y}{\partial \theta} = r \cos \theta$$

Now we need to find u_{rr} and $u_{\theta\theta}$. Let's focus on u_{rr} first. We will need to start by finding u_r . We have as follows

$$\begin{aligned} \frac{\partial u}{\partial r} &= \frac{\partial u}{\partial x} \frac{\partial x}{\partial r} + \frac{\partial u}{\partial y} \frac{\partial y}{\partial r} \\ &= \cos \theta \frac{\partial u}{\partial x} + \sin \theta \frac{\partial u}{\partial y} \end{aligned}$$

We have u_r , so now we need to compute u_{rr} . We have as follows:

$$\begin{aligned} \frac{\partial^2 u}{\partial r^2} &= \cos \theta \frac{\partial}{\partial r} \frac{\partial u}{\partial x} + \sin \theta \frac{\partial}{\partial r} \frac{\partial u}{\partial y} \\ &= \cos^2 \theta \frac{\partial^2 u}{\partial x^2} + 2 \cos \theta \sin \theta \frac{\partial^2 u}{\partial x \partial y} + \sin^2 \theta \frac{\partial^2 u}{\partial y^2} \end{aligned}$$

We won't delve into finding $u_{\theta\theta}$, as the partials get a little out of hand. However, we end up with

$$\begin{aligned} \frac{\partial^2 u}{\partial r^2} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} &= -\frac{1}{r} \frac{\partial u}{\partial r} + \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \\ u_{xx} + u_{yy} &= u_{rr} + \frac{1}{r} u_r + \frac{1}{r^2} u_{\theta\theta} = 0 \end{aligned}$$

Lets go ahead and formalize it:

$$\nabla^2 u = u_{rr} + \frac{1}{r} u_r + \frac{1}{r^2} u_{\theta\theta}$$

Now we can treat this as a 2D problem; in other words, we can now use separation of variables to solve for u . As before, we know our solution will look like

$$u(r, \theta) = R(r)T(\theta)$$

Plugging everything in and realizing by the usual trick that everything's equal to a constant gives us

$$-\frac{T''}{T} = \frac{R'' + \frac{R'}{r}}{\frac{R}{r^2}} = \lambda$$

Since we are dealing in polar coordinates, we will go ahead and walk through the eigenvalues/functions. We will be solving for $T(\theta)$, since our boundary conditions are in terms of θ . We also should note that the solutions will repeat themselves every 2π , since we are in the polar system. In addition, our u should be finite at $r = 0$. Let's take a look at the eigenvalues for the Laplace equation.

$$\underline{\lambda = 0}$$

When $\lambda = 0$, we have

$$T(\theta) = a\theta + b.$$

This also tells us that

$$R(r) = c \ln(r) + d.$$

Putting everything together gives us

$$u(r, \theta) = (a\theta + b)(c \ln r + d).$$

Since the solution is periodic on 2π , we know that $a = 0$. Since the solution has to remain finite at $r = 0$, we know that $c = 0$. This gives us that $u = bd$, or that u is equal to a constant.

So we know that the only λ that yields eigenvalues is $\lambda = 0$, and we know that this particular eigenvalue yields a constant solution. If we solve the differential equation for R , we will find that the only solution we care about is r^n . We can finally say that our solution will look like

$$u(r, \theta) = \frac{a_0}{2} + \sum_{n=1}^{\infty} r^n [a_n \cos(n\theta) + b_n \sin(n\theta)]$$

This solution is worth memorizing, as it will generally be the same for all variations of the Laplace equation similar to this one. Now we can apply the conditions for θ . We have two conditions, but it is really just a piecewise function which we will call $f(\theta)$. We also know that our bound on the domain is a , or the length of the radius. This is enough information to find all the various variables in our $u(r, \theta)$. We have as follows:

$$u(a, \theta) = f(\theta) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a^n [a_n \cos(n\theta) + b_n \sin(n\theta)]$$

Note that the a which is the length of the radius is *not* the same a as a_n . Now we can solve for a_0 and a_n .

$$\begin{aligned} a_0 &= \frac{1}{\pi} \int_0^{2\pi} f(\theta) d\theta & a_n &= \frac{1}{\pi} \left[\int_0^{\pi} U_1 \cos(n\theta) d\theta + \int_{\pi}^{2\pi} U_2 \cos(n\theta) d\theta \right] \\ &= U_1 + U_2 & &= 0 \end{aligned}$$

For b_n , we have

$$\begin{aligned} a^n b_n &= \frac{1}{\pi} \left[\int_0^{\pi} U_1 \sin(n\theta) d\theta + \int_{\pi}^{2\pi} U_2 \sin(n\theta) d\theta \right] \\ &= \frac{1 - (-1)^n}{n\pi} (U_1 - U_2) \\ b_n &= \frac{[1 - (-1)^n] (U_1 - U_2)}{n\pi a^n} \\ &= \frac{2(U_1 - U_2)}{(2k + 1)\pi a^{2k+1}} \end{aligned}$$

Putting it all together gives us a final solution of

$$u(r, \theta) = \frac{U_1 + U_2}{2} + \sum_{k=0}^{\infty} r^k \frac{2(U_1 - U_2)}{(2k + 1)\pi a^{2k+1}}$$

And we are done.

Now let's look at another Laplace equation in polar coordinates. Again, we will be dealing with Dirichlet boundary conditions, so we will be able to jump straight into the solution. This one won't take us as long to solve, as we have already done a lot of the work. We are asked the following:

Ex: Solve the following PDE

$$\begin{cases} \nabla^2 u = 0 \\ u(3, \theta) = \begin{cases} 1 & 0 \leq \theta \leq \pi \\ \sin^2 \theta & \pi \leq \theta \leq 2\pi \end{cases} \end{cases}$$

Solution: We already know what our separable equation will look like:

$$\frac{r^2 R'' + r R'}{R} = -\frac{T''}{T} = -\lambda$$

This yields an eigenvalue of $\lambda = n^2$. We know this because our function T must be periodic on 2π , and as such we get the following piecewise function:

$$T(\theta) = \begin{cases} a_0 & \text{for } n = 0 \\ a_n \cos(n\theta) + b_n \sin(n\theta) & \text{for } n > 0 \end{cases}$$

The solution to our equation for R yields

$$R(r) = \begin{cases} c_1 + c_2 \ln r & \text{for } n = 0 \\ c_1 r^n + c_2 r^{-n} & \text{for } n > 0 \end{cases}$$

We can disregard c_2 for this equation, since it would cause our solution to go to ∞ as $r \rightarrow 0$, at the center of the disk. Putting everything together gives us

$$\begin{aligned} u(r, \theta) &= R(r)T(\theta) \\ &= \begin{cases} a_0 & \text{for } n = 0 \\ a_n r^n \cos(n\theta) + b_n r^n \sin(n\theta) & \text{for } n > 0 \end{cases} \end{aligned}$$

We should probably construct a series out of this, since that is what we usually do. The corresponding series is

$$u(r, \theta) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n r^n \cos(n\theta) + b_n r^n \sin(n\theta)$$

And applying the boundary conditions gives us

$$u(3, \theta) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n 3^n \cos(n\theta) + b_n 3^n \sin(n\theta)$$

For a_0 , we have

$$\begin{aligned} a_0 &= \frac{1}{\pi} \left[\int_0^{\pi} 1 d\theta + \int_{\pi}^{2\pi} \sin^2 \theta d\theta \right] \\ &= \frac{3}{2} \end{aligned}$$

Solving for a_n gives us

$$\begin{aligned} a_n &= \frac{1}{3^n \pi} \left[\int_0^{\pi} \cos(n\theta) d\theta + \int_{\pi}^{2\pi} \cos(n\theta) \sin^2 \theta d\theta \right] \\ &= \frac{[n^2 - 2 - 4 \cos(n\pi)] \sin(n\pi)}{3^n n(n^2 - 4) \pi} \\ &= 0 \end{aligned}$$


Our b_n term is

$$\begin{aligned}
b_n &= \frac{1}{3^n \pi} \left[\int_0^\pi \sin(n\theta) d\theta + \int_\pi^{2\pi} \sin(n\theta) \sin^2 \theta d\theta \right] \\
&= \frac{2 \left[n^2 - 6 - 4 \cos(n\pi) \right] \sin^2 \left(\frac{n\pi}{2} \right)}{3^n n \pi (n^2 - 4)} \\
&= \frac{2 \left[n^2 - 6 - 4(-1)^n \right] \sin^2 \left(\frac{n\pi}{2} \right)}{3^n n \pi (n^2 - 4)} \\
&= \begin{cases} 0 & n \in \{2k \mid k \in \mathbb{Z}\} \\ \frac{1}{3^n \pi} \left[\frac{2}{n} + \frac{4}{n(n^2-4)} \right] & n \in \{2k+1 \mid k \in \mathbb{Z}\} \end{cases}
\end{aligned}$$


We essentially have our solution. All we have to do is combine everything:

$$u(r, \theta) = \frac{3}{4} + \sum_{k=0}^{\infty} \frac{r^{2k+1}}{3^{2k+1} \pi} \frac{2(4k^2 + 4k - 1)}{(2k+3)(4k^2 - 1)} \sin[(2k+1)\theta]$$


Let's take a look at some of the plots for this equation.



The first 3 terms



The first 30 terms



The first 50 terms

Plugging this infinite sum into mathematica tells us that it does not converge; and we are done.