

Steven Rosendahl
Homework 8

1. Determine whether each of the following polynomials has a zero in the given \mathbb{Z}_p . Either find a zero or prove that no zero exists.

(a) $f(x) = x^3 + x^2 + x + 1$ in \mathbb{Z}_2 .

We can factor $x^3 + x^2 + x + 1$ into $(x^2 + 1)(x + 1)$. The only two options for a root in \mathbb{Z}_2 are 1 and 0. If we plug in zero we get $(0 + 1)(0 + 1) \equiv 1 \pmod{2}$, so 1 is not a zero of the polynomial. If we try 1, we get $(1 + 1)(1 + 1) \equiv (2)(2) \equiv (0)(0) \equiv 0 \pmod{2}$, so 1 is a zero of the polynomial.

(b) $f(x) = x^3 + x^2 + x + 1$ in \mathbb{Z}_3 .

$f(x)$ factors into $(x^2 + 1)(x + 1) \equiv (x^2 - 2)(x - 2) \pmod{3}$. We have one zero at $x = 2$. If we test 0, we get $(0 + 1)(0 + 1) \equiv 1 \pmod{3}$, so 0 is not a zero of the polynomial, and if we test 1 we get $(-1)(-1) \equiv 1 \pmod{3}$, so 1 is not a zero either.

(c) $f(x) = x^2 + 2x + 3$ in \mathbb{Z}_5 .

The polynomial $f(x) = x^2 + 2x + 3$ cannot be factored in \mathbb{Z}_5 . Therefore, it has no zeros since we cannot express it in the form $g(x)(x - \alpha)$.

2. Let $f(x) = x^2 + 1$. Prove that f has a zero in \mathbb{Z}_5 , but not in \mathbb{Z}_7 .

If we consider the polynomial in \mathbb{Z}_5 , we have that

$$x^2 + 1 \equiv x^2 - 4 \equiv (x + 2)(x - 2) \equiv (x - 3)(x - 2) \pmod{5}.$$

Since f was reducible in \mathbb{Z}_5 , we know it has at least one zero, and in this case it has two zeros, $x = 3$ and $x = 5$. In \mathbb{Z}_7 , we have

$$x^2 + 1 \equiv x^2 - 6 \pmod{7}.$$

The polynomial $x^2 - 6$ is irreducible in \mathbb{Z}_7 , so there are no roots.

3. Suppose that p is prime, k is a non-zero element of \mathbb{Z}_p , and $f_k(x) = x^2 - k$.

- (a) Prove that s is a zero of f_k if and only if $-s$ is also a zero of f_k .

\Rightarrow) Suppose s is a zero of f_k . Then $s^2 - k \equiv 0 \pmod{p}$. If $-s$ is not a zero, then we have $(-s)^2 - k \not\equiv 0 \pmod{p}$, so $s^2 - k \not\equiv 0 \pmod{p}$, which is a contradiction.

\Leftarrow) Suppose $-s$ is a zero of f_k . Then $(-s)^2 - k \equiv 0 \pmod{p}$. Then $s^2 - k \equiv 0 \pmod{p}$, so s is a zero of f_k .

- (b) Further assuming that $p > 2$ prove that f_k either has no zeros in \mathbb{Z}_p or exactly two zeros in \mathbb{Z}_p .

Suppose there was only one zero in \mathbb{Z}_p . We know this cannot be the case, since by (a) we showed that if s is a zero, then $-s$ is also a zero. If we suppose there are more than 2 zeros, then we can consider another zero, σ . By (a), we know that $-\sigma$ is also a zero. Since $\deg(f) = 2$, we know that there are at most 2 zeros in \mathbb{Z}_p . There cannot be just one zero, so there must be only two zeros, or no zeros.

4. Let p be a prime with $p > 2$ and assume that $a, b, c, r \in \mathbb{Z}_p$ with $a \not\equiv 0 \pmod{p}$. Further, we define the polynomial $f(x) = ax^2 + bx + c \in \mathbb{Z}_p[x]$.

- (a) Prove that r is a zero of f in \mathbb{Z}_p if and only if $(2ar + b)^2 \equiv b^2 - 4ac \pmod{p}$.

\Rightarrow) Since r is a zero of $f(x)$ and f is a quadratic polynomial, we can create the expression

$$\begin{aligned} ar^2 + br + c &\equiv 0 \pmod{p} \\ ar^2 + br &\equiv -c \pmod{p}. \end{aligned}$$

Consider the term $(2ar + b)^2$. Then

$$\begin{aligned}
 (2ar + b)^2 &= 4a^2r^2 + 4bar + b^2 \\
 &= 4a(ar^2 + br) + b^2 \\
 &= 4a(-c) + b^2 \\
 &= b^2 - 4ac \\
 &\equiv b^2 - 4ac \pmod{p}
 \end{aligned}$$

\Leftarrow) Suppose $(2ar + b)^2 \equiv b^2 - 4ac \pmod{p}$. Then

$$\begin{aligned}
 4a^2r^2 + 4arb + b^2 &\equiv b^2 - 4ac \pmod{p} \\
 4a^2r^2 + 4arb &\equiv -4ac \pmod{p} \\
 4a^2r^2 + 4arb + 4ac &\equiv 0 \pmod{p} \\
 4a(ar^2 + br + c) &\equiv 0 \pmod{p}.
 \end{aligned}$$

Since we are in \mathbb{Z}_p , we know all elements have an inverse. We can multiply both sides of the congruence by $a^{-1}4^{-1}$, which gives us $ar^2 + br + c \equiv 0 \pmod{p}$. Therefore, r is a root of f .

(b) A point $y \in \mathbb{Z}_p$ is called a *perfect square* if there exists $z \in \mathbb{Z}_p$ such that $z^2 = y$.

i. If f has at least one zero in \mathbb{Z}_p , prove that $b^2 - 4ac$ is a perfect square.

Suppose r is a zero of f . Then by (a) we have

$$(2ar + b)^2 \equiv b^2 - 4ac \pmod{p},$$

which implies $b^2 - 4ac$ is a perfect square.

ii. If $b^2 - 4ac \equiv 0 \pmod{p}$, prove that f has a unique zero in f . Find a formula for that zero in terms of a and b .

Suppose $b^2 - 4ac \equiv 0 \pmod{p}$. We know by (a) that r is root, and therefore $(2ar + b)^2 \equiv b^2 - 4ac \pmod{p}$. Suppose ρ is a root of f . then $(2a\rho + b)^2 \equiv b^2 - 4ac \pmod{p}$, so $(2ar + b)^2 \equiv (2a\rho + b)^2$. Since $b^2 - 4ac$ is a perfect square, then $2ar + b \equiv 2a\rho + b$. Then $2ar \equiv 2a\rho$, and multiplying by $2^{-1}a^{-1}$ yields $r \equiv \rho$. To find an expression for r , we know that $(2ar + b)^2 = 0$, so if we multiply by $(2ar + b)^{-1}$, we have $2ar + b = 0$. Then $r = -b2^{-1}a^{-1}$.

iii. If $b^2 - 4ac \not\equiv 0 \pmod{p}$ and $b^2 - 4ac$ is a perfect square, prove that f has exactly two distinct zeros in \mathbb{Z}_p .

We know that we have the equivalence $(2ar + b)^2 \equiv b^2 - 4ac$. Let $x = (2ar + b)^2$ and $k = b^2 - 4ac$. Then we can rearrange the congruence to say $x^2 - k \equiv 0 \pmod{p}$. We know $x^2 - k$ has either 0 or 2 zeros. Since k is a perfect square, we can let $k = j^2$ for some $j \in \mathbb{Z}_p$. Then $x^2 - k = x^2 - j^2 = (x + j)(x - j)$, which has two distinct roots in \mathbb{Z}_p .

5. Suppose that $g(x) = x^2 + 1 \in \mathbb{Z}_3[x]$ and let ϕ be a zero of g .

(a) Prove that $\phi \notin \mathbb{Z}_3$.

(b) Define the set $\mathbb{F}_9 = \{0, 1, 2, \phi, \phi + 1, \phi + 2, 2\phi, 2\phi + 1, 2\phi + 2\}$. Assuming that the multiplication and addition in \mathbb{F}_9 obeys the distributive law, prove that every element of \mathbb{F}_9 has a multiplicative inverse.