

Principles of Counting and Introduction to Probability Theory

Sourav Karmakar

- 1 Chapter 1: Counting Principles (Combinatorics)
- 2 Chapter 2: Introduction to Probability Theory

Counting Principles

The Mathematics of "How Many?"

1. The Multiplication Rule of Counting

Fundamental Counting Principle

If a task can be broken down into a sequence of k steps, where:

- Step 1 can be done in n_1 ways,
- Step 2 can be done in n_2 ways, ...,
- Step k can be done in n_k ways,


Then the total number of ways to perform the task is:

$$N = n_1 \times n_2 \times \cdots \times n_k$$

Multiplication Rule: Visual Example

Scenario: Ordering a "Combo Meal".

- **Main Dish:** Burger or Sandwich (2 options).
- **Drink:** Coke, Water, or Juice (3 options).



Total Combinations = $2 \times 3 = 6$ distinct meals.

Multiplication Rule: Numerical Examples

Example 1: The ATM Pin

A bank card requires a 4-digit PIN (digits 0-9). How many unique PINs are possible?

Multiplication Rule: Numerical Examples

Example 1: The ATM Pin

A bank card requires a 4-digit PIN (digits 0-9). How many unique PINs are possible?

- Pos 1: 10 options (0-9)
- Pos 2: 10 options
- Pos 3: 10 options
- Pos 4: 10 options
- **Total:** $10 \times 10 \times 10 \times 10 = 10,000$ PINs.

Multiplication Rule: Numerical Examples

Example 1: The ATM Pin

A bank card requires a 4-digit PIN (digits 0-9). How many unique PINs are possible?

- Pos 1: 10 options (0-9)
- Pos 2: 10 options
- Pos 3: 10 options
- Pos 4: 10 options
- **Total:** $10 \times 10 \times 10 \times 10 = 10,000$ PINs.

Example 2: Buying a Car

You are configuring a new car online. The options are: 1. Model: (Sedan, SUV, Hatchback) \rightarrow 3 options. 2. Color: (Red, Black, White, Silver) \rightarrow 4 options. 3. Transmission: (Manual, Automatic) \rightarrow 2 options.

Total Configurations: $3 \times 4 \times 2 = 24$ unique cars.

2. The Addition Rule

Rule of Sum

If a task can be done in n_A ways OR in n_B ways, and these sets of ways are mutually exclusive (you cannot do both at once), then:

$$\text{Total ways} = n_A + n_B$$

2. The Addition Rule

Rule of Sum

If a task can be done in n_A ways OR in n_B ways, and these sets of ways are mutually exclusive (you cannot do both at once), then:

$$\text{Total ways} = n_A + n_B$$

Example: Commute Options

You want to go to the office.

- You can take a **Bus** (3 different routes).
- OR you can take a **Train** (2 different lines).


You cannot take both simultaneously.

Total Choices: $3 + 2 = 5$ ways to get to work.

Combining Addition and Multiplication

Problem: Travel from City A to City C via City B.


- **A to B:** 2 Flights OR 3 Trains.
- **B to C:** 2 Buses.



Combining Addition and Multiplication

Problem: Travel from City A to City C via City B.

- **A to B:** 2 Flights OR 3 Trains.
- **B to C:** 2 Buses.




Solution:

- ① **A to B (Addition):**
 $2 \text{ (F)} + 3 \text{ (T)} = 5 \text{ ways.}$

Combining Addition and Multiplication

Problem: Travel from City A to City C via City B.

- **A to B:** 2 Flights OR 3 Trains.
- **B to C:** 2 Buses.




Solution:

- 1 **A to B (Addition):**
 $2 \text{ (F)} + 3 \text{ (T)} = 5 \text{ ways.}$
- 2 **B to C:**
2 ways.

Combining Addition and Multiplication

Problem: Travel from City A to City C via City B.

- **A to B:** 2 Flights OR 3 Trains.
- **B to C:** 2 Buses.



Solution:

- 1 **A to B (Addition):**
 $2 \text{ (F)} + 3 \text{ (T)} = 5 \text{ ways.}$
- 2 **B to C:**
 2 ways.
- 3 **Total (Multiplication):**
 $5 \times 2 = 10 \text{ routes.}$

Factorials

Before we discuss Permutations, we must define the **Factorial**.

Definition

The factorial of a non-negative integer n , denoted by $n!$, is the product of all positive integers less than or equal to n .

$$n! = n \times (n - 1) \times (n - 2) \times \cdots \times 2 \times 1$$

Factorials

Before we discuss Permutations, we must define the **Factorial**.

Definition

The factorial of a non-negative integer n , denoted by $n!$, is the product of all positive integers less than or equal to n .

$$n! = n \times (n - 1) \times (n - 2) \times \cdots \times 2 \times 1$$

Recursion Rule: $n! = n \times (n - 1)!$

Factorials

Before we discuss Permutations, we must define the **Factorial**.

Definition

The factorial of a non-negative integer n , denoted by $n!$, is the product of all positive integers less than or equal to n .

$$n! = n \times (n - 1) \times (n - 2) \times \cdots \times 2 \times 1$$

Recursion Rule: $n! = n \times (n - 1)!$

The Special Case of Zero: Why is $0! = 1$?

$$\text{Since } n! = n \times (n - 1)! \implies 1! = 1 \times 0! \implies 1 = 1 \times 0! \implies 0! = 1$$

Examples:

- $3! = 3 \times 2 \times 1 = 6$
- $5! = 5 \times 4 \times 3 \times 2 \times 1 = 120$


3. Permutations (Order Matters)

Key Intuition

Permutation is about **arrangement**. The order in which you place objects changes the outcome (e.g., Password "123" \neq "321").

Formula: The number of ways to arrange r objects chosen from n distinct objects:

$$P(n, r) = \frac{n!}{(n - r)!}$$



Permutations: Numerical Examples

Example 1: [Ranking] 10 participants in a race. How many ways can we award Gold, Silver, and Bronze?

Permutations: Numerical Examples

Example 1: [Ranking] 10 participants in a race. How many ways can we award Gold, Silver, and Bronze?

$$P(10, 3) = \frac{10!}{(7)!} = 10 \times 9 \times 8 = 720 \text{ ways.}$$

Permutations: Numerical Examples

Example 1: [Ranking] 10 participants in a race. How many ways can we award Gold, Silver, and Bronze?

$$P(10, 3) = \frac{10!}{(7)!} = 10 \times 9 \times 8 = 720 \text{ ways.}$$

Example 2: [Bookshelf] You have 5 different books. In how many ways can you arrange all of them on a shelf?

Permutations: Numerical Examples

Example 1: [Ranking] 10 participants in a race. How many ways can we award Gold, Silver, and Bronze?

$$P(10, 3) = \frac{10!}{(7)!} = 10 \times 9 \times 8 = 720 \text{ ways.}$$

Example 2: [Bookshelf] You have 5 different books. In how many ways can you arrange all of them on a shelf?

$$P(5, 5) = \frac{5!}{0!} = 5! = 120 \text{ ways.}$$

Permutations: Numerical Examples

Example 1: [Ranking] 10 participants in a race. How many ways can we award Gold, Silver, and Bronze?

$$P(10, 3) = \frac{10!}{(7)!} = 10 \times 9 \times 8 = 720 \text{ ways.}$$

Example 2: [Bookshelf] You have 5 different books. In how many ways can you arrange all of them on a shelf?

$$P(5, 5) = \frac{5!}{0!} = 5! = 120 \text{ ways.}$$

Example 3: [Corporate Board] A board has 12 members. They need to elect a Chairman, a Secretary, and a Treasurer. No person can hold two posts.

Permutations: Numerical Examples

Example 1: [Ranking] 10 participants in a race. How many ways can we award Gold, Silver, and Bronze?

$$P(10, 3) = \frac{10!}{(7)!} = 10 \times 9 \times 8 = 720 \text{ ways.}$$

Example 2: [Bookshelf] You have 5 different books. In how many ways can you arrange all of them on a shelf?

$$P(5, 5) = \frac{5!}{0!} = 5! = 120 \text{ ways.}$$

Example 3: [Corporate Board] A board has 12 members. They need to elect a Chairman, a Secretary, and a Treasurer. No person can hold two posts.

$$P(12, 3) = 12 \times 11 \times 10 = 1,320 \text{ ways.}$$

4. Combinations (Order Doesn't Matter)

Key Intuition

Combination is about **selection**. "Fruit Salad" logic: Apple and Banana is the same as Banana and Apple.

Formula: The number of ways to choose r objects from n distinct objects:

$$C(n, r) = \binom{n}{r} = \frac{n!}{r!(n-r)!}$$

Note: Combinations result in fewer counts than permutations because duplicates (e.g., AB vs BA) are removed/divided out.

Combinations: Numerical Examples

Example 1: [Team Selection] A manager needs to pick 3 engineers from a pool of 10 to go on a trip.

Combinations: Numerical Examples

Example 1: [Team Selection] A manager needs to pick 3 engineers from a pool of 10 to go on a trip.

$$C(10, 3) = \frac{10 \times 9 \times 8}{3 \times 2 \times 1} = 120 \text{ teams.}$$

Combinations: Numerical Examples

Example 1: [Team Selection] A manager needs to pick 3 engineers from a pool of 10 to go on a trip.

$$C(10, 3) = \frac{10 \times 9 \times 8}{3 \times 2 \times 1} = 120 \text{ teams.}$$

Example 2: [Handshakes] In a meeting of 6 people, if everyone shakes hands once with everyone else, how many handshakes occurred?

Combinations: Numerical Examples

Example 1: [Team Selection] A manager needs to pick 3 engineers from a pool of 10 to go on a trip.

$$C(10, 3) = \frac{10 \times 9 \times 8}{3 \times 2 \times 1} = 120 \text{ teams.}$$

Example 2: [Handshakes] In a meeting of 6 people, if everyone shakes hands once with everyone else, how many handshakes occurred?

$$C(6, 2) = \frac{6 \times 5}{2 \times 1} = 15 \text{ handshakes.}$$

Combinations: Numerical Examples

Example 1: [Team Selection] A manager needs to pick 3 engineers from a pool of 10 to go on a trip.

$$C(10, 3) = \frac{10 \times 9 \times 8}{3 \times 2 \times 1} = 120 \text{ teams.}$$

Example 2: [Handshakes] In a meeting of 6 people, if everyone shakes hands once with everyone else, how many handshakes occurred?

$$C(6, 2) = \frac{6 \times 5}{2 \times 1} = 15 \text{ handshakes.}$$

Example 3: [Lottery] A lottery ticket requires you to choose 6 numbers from 1 to 49 (order of drawing doesn't matter). How many possible tickets exist?

$$C(49, 6) = \frac{49!}{6!43!} = 13,983,816 \text{ tickets.}$$

Quiz: Counting Principles

Q1: [License Plates]. A plate has 3 letters followed by 3 digits. Letters cannot repeat, but digits can. How many plates?

Quiz: Counting Principles

Q1: [License Plates]. A plate has 3 letters followed by 3 digits. Letters cannot repeat, but digits can. How many plates?

- Letters (Permutation): $26 \times 25 \times 24$
- Digits (Multiplication): $10 \times 10 \times 10$
- **Ans:** $15,600 \times 1,000 = 15,600,000$

Quiz: Counting Principles

Q1: [License Plates]. A plate has 3 letters followed by 3 digits. Letters cannot repeat, but digits can. How many plates?

- Letters (Permutation): $26 \times 25 \times 24$
- Digits (Multiplication): $10 \times 10 \times 10$
- **Ans:** $15,600 \times 1,000 = 15,600,000$

Q2: [Committee Restrictions]. Form a committee of 4 from 5 Men and 6 Women. The committee **must** contain exactly 2 Men and 2 Women.

Quiz: Counting Principles

Q1: [License Plates]. A plate has 3 letters followed by 3 digits. Letters cannot repeat, but digits can. How many plates?

- Letters (Permutation): $26 \times 25 \times 24$
- Digits (Multiplication): $10 \times 10 \times 10$
- **Ans:** $15,600 \times 1,000 = 15,600,000$

Q2: [Committee Restrictions]. Form a committee of 4 from 5 Men and 6 Women. The committee **must** contain exactly 2 Men and 2 Women.

- Choose Men: $C(5, 2) = 10$. Choose Women: $C(6, 2) = 15$.
- **Ans:** $10 \times 15 = 150$ ways.

Quiz: Counting Principles

Q1: [License Plates]. A plate has 3 letters followed by 3 digits. Letters cannot repeat, but digits can. How many plates?

- Letters (Permutation): $26 \times 25 \times 24$
- Digits (Multiplication): $10 \times 10 \times 10$
- **Ans:** $15,600 \times 1,000 = 15,600,000$

Q2: [Committee Restrictions]. Form a committee of 4 from 5 Men and 6 Women. The committee **must** contain exactly 2 Men and 2 Women.

- Choose Men: $C(5, 2) = 10$. Choose Women: $C(6, 2) = 15$.
- **Ans:** $10 \times 15 = 150$ ways.

Q3: [Seating]. In how many ways can 5 people sit in a row if Person A and Person B **must** sit next to each other?

Quiz: Counting Principles

Q1: [License Plates]. A plate has 3 letters followed by 3 digits. Letters cannot repeat, but digits can. How many plates?

- Letters (Permutation): $26 \times 25 \times 24$
- Digits (Multiplication): $10 \times 10 \times 10$
- **Ans:** $15,600 \times 1,000 = 15,600,000$

Q2: [Committee Restrictions]. Form a committee of 4 from 5 Men and 6 Women. The committee **must** contain exactly 2 Men and 2 Women.

- Choose Men: $C(5, 2) = 10$. Choose Women: $C(6, 2) = 15$.
- **Ans:** $10 \times 15 = 150$ ways.

Q3: [Seating]. In how many ways can 5 people sit in a row if Person A and Person B **must** sit next to each other?

- Treat (AB) as 1 unit. We arrange 4 units: $4! = 24$.
- A and B can swap places inside the unit: $2! = 2$.
- **Ans:** $24 \times 2 = 48$ ways.

Introduction to Probability Theory

Quantifying Uncertainty

1. Random Experiment & Sample Space

Random Experiment: A process where the outcome cannot be predicted with certainty.

Sample Space (S): The set of **all** possible outcomes.

Single Coin Toss:

$$S = \{H, T\}$$

$$|S| = 2$$

Rolling a Die:

$$S = \{1, 2, 3, 4, 5, 6\}$$

$$|S| = 6$$

The elements of the Sample Space follows the *MECE Principle*: Mutually Exclusive (outcomes don't overlap) and Collectively Exhaustive (cover all possibilities).

2. Defining Probability

Probability of an Event A: For a finite sample space where all outcomes are equally likely (Count-based):

$$P(A) = \frac{\text{Number of favorable outcomes}}{\text{Total number of possible outcomes}} = \frac{|A|}{|S|}$$

$$0 \leq P(A) \leq 1$$

Probability Examples (Count Based)

Example 1: [Tossing Two Coins] What is the probability of getting **at least one Head**?

Probability Examples (Count Based)

Example 1: [Tossing Two Coins] What is the probability of getting **at least one Head**?

- $S = \{HH, HT, TH, TT\} \implies |S| = 4$.
- Favorable outcomes $A = \{HH, HT, TH\}$.
- $P(A) = 3/4 = 0.75$.

Probability Examples (Count Based)

Example 1: [Tossing Two Coins] What is the probability of getting **at least one Head**?

- $S = \{HH, HT, TH, TT\} \implies |S| = 4$.
- Favorable outcomes $A = \{HH, HT, TH\}$.
- $P(A) = 3/4 = 0.75$.

Example 2: [Rolling Two Dice] What is the probability that the **sum of the dice is 7**?

Probability Examples (Count Based)

Example 1: [Tossing Two Coins] What is the probability of getting **at least one Head**?

- $S = \{HH, HT, TH, TT\} \implies |S| = 4$.
- Favorable outcomes $A = \{HH, HT, TH\}$.
- $P(A) = 3/4 = 0.75$.

Example 2: [Rolling Two Dice] What is the probability that the **sum of the dice is 7**?

- Total outcomes: $6 \times 6 = 36$.
- Favorable pairs (x, y) : $\{(1, 6), (2, 5), (3, 4), (4, 3), (5, 2), (6, 1)\}$.
- Count = 6.
- $P(\text{Sum} = 7) = 6/36 = 1/6 \approx 0.167$.

3. Kolmogorov's Axioms of Probability

The foundation of modern probability.

- ① **Non-negativity:** For any event E , $P(E) \geq 0$.
- ② **Normalization:** The probability of the sample space is 1. $P(S) = 1$.
- ③ **Additivity:** If A and B are mutually exclusive (cannot happen together), then:

$$P(A \cup B) = P(A) + P(B)$$

3. Kolmogorov's Axioms of Probability

The foundation of modern probability.

- ① **Non-negativity:** For any event E , $P(E) \geq 0$.
- ② **Normalization:** The probability of the sample space is 1. $P(S) = 1$.
- ③ **Additivity:** If A and B are mutually exclusive (cannot happen together), then:


$$P(A \cup B) = P(A) + P(B)$$

Useful Corollaries:

- **Complement Rule:** $P(A^c) = 1 - P(A)$.
- **Subset Rule:** If $A \subset B$, then $P(A) \leq P(B)$.
- (If A is part of B , A cannot be more probable than B).


4. Events and Venn Diagrams

Union ($A \cup B$): OR



Any shaded area

Intersection ($A \cap B$): AND




Only the overlap

Complementary Events

Complement (A' or A^c): NOT A

$$P(A') = 1 - P(A)$$




Physically: If probability of system failure is 0.01, reliability is 0.99.

5. Real World Example: Parallel Redundancy

Problem: Three servers (A, B, C) are connected in **Parallel**. System fails only if **ALL** servers fail.

- $P(\text{Fail}_A) = 0.1$
- $P(\text{Fail}_B) = 0.1$
- $P(\text{Fail}_C) = 0.1$

Find the probability of System Success.



Step 1: Find Probability of Total Failure Assuming failures are independent:

$$P(\text{System Fail}) = P(F_A) \times P(F_B) \times P(F_C)$$

Step 1: Find Probability of Total Failure Assuming failures are independent:

$$P(\text{System Fail}) = P(F_A) \times P(F_B) \times P(F_C)$$

$$P(\text{System Fail}) = 0.1 \times 0.1 \times 0.1 = 0.001$$

Step 1: Find Probability of Total Failure Assuming failures are independent:

$$P(\text{System Fail}) = P(F_A) \times P(F_B) \times P(F_C)$$

$$P(\text{System Fail}) = 0.1 \times 0.1 \times 0.1 = 0.001$$

Step 2: Find Probability of Success (Complement Rule)

$$P(\text{Success}) = 1 - P(\text{System Fail})$$

$$P(\text{Success}) = 1 - 0.001 = 0.999$$

Conclusion: Redundancy improves reliability from 90% (single server) to 99.9%.

Quiz: Probability

Q1: [Defective Items]. A box has 5 Good and 3 Bad parts. You pick 2 parts at random (without replacement). What is the probability **both are Bad**?

Quiz: Probability

Q1: [Defective Items]. A box has 5 Good and 3 Bad parts. You pick 2 parts at random (without replacement). What is the probability **both are Bad**?

- Total ways to pick 2: $C(8, 2) = 28$.
- Ways to pick 2 Bad: $C(3, 2) = 3$.
- **Ans:** $3/28 \approx 0.107$.

Quiz: Probability

Q1: [Defective Items]. A box has 5 Good and 3 Bad parts. You pick 2 parts at random (without replacement). What is the probability **both are Bad**?

- Total ways to pick 2: $C(8, 2) = 28$.
- Ways to pick 2 Bad: $C(3, 2) = 3$.
- **Ans:** $3/28 \approx 0.107$.

Q2: [At Least One]. A shooter hits the target with $P = 0.4$. If he shoots 3 times, what is the probability he hits the target **at least once**?

Quiz: Probability

Q1: [Defective Items]. A box has 5 Good and 3 Bad parts. You pick 2 parts at random (without replacement). What is the probability **both are Bad**?

- Total ways to pick 2: $C(8, 2) = 28$.
- Ways to pick 2 Bad: $C(3, 2) = 3$.
- **Ans:** $3/28 \approx 0.107$.

Q2: [At Least One]. A shooter hits the target with $P = 0.4$. If he shoots 3 times, what is the probability he hits the target **at least once**?

- Use Complement. $P(\text{Miss}) = 1 - 0.4 = 0.6$.
- $P(\text{Miss All}) = 0.6 \times 0.6 \times 0.6 = 0.216$.
- **Ans:** $1 - 0.216 = 0.784$.

Quiz: Probability

Q1: [Defective Items]. A box has 5 Good and 3 Bad parts. You pick 2 parts at random (without replacement). What is the probability **both are Bad**?

- Total ways to pick 2: $C(8, 2) = 28$.
- Ways to pick 2 Bad: $C(3, 2) = 3$.
- **Ans:** $3/28 \approx 0.107$.

Q2: [At Least One]. A shooter hits the target with $P = 0.4$. If he shoots 3 times, what is the probability he hits the target **at least once**?

- Use Complement. $P(\text{Miss}) = 1 - 0.4 = 0.6$.
- $P(\text{Miss All}) = 0.6 \times 0.6 \times 0.6 = 0.216$.
- **Ans:** $1 - 0.216 = 0.784$.

Q3: [Difference]. Two dice are rolled. What is the probability the **difference between the numbers is exactly 3**?

Quiz: Probability

Q1: [Defective Items]. A box has 5 Good and 3 Bad parts. You pick 2 parts at random (without replacement). What is the probability **both are Bad**?

- Total ways to pick 2: $C(8, 2) = 28$.
- Ways to pick 2 Bad: $C(3, 2) = 3$.
- **Ans:** $3/28 \approx 0.107$.

Q2: [At Least One]. A shooter hits the target with $P = 0.4$. If he shoots 3 times, what is the probability he hits the target **at least once**?

- Use Complement. $P(\text{Miss}) = 1 - 0.4 = 0.6$.
- $P(\text{Miss All}) = 0.6 \times 0.6 \times 0.6 = 0.216$.
- **Ans:** $1 - 0.216 = 0.784$.

Q3: [Difference]. Two dice are rolled. What is the probability the **difference between the numbers is exactly 3**?

- Pairs: (1,4), (2,5), (3,6), (4,1), (5,2), (6,3). Total 6 pairs.
- Total Outcomes: 36.
- **Ans:** $6/36 = 1/6$.

Thank You!

Next Steps: Conditional Probability and Bayes' Theorem.