



Comprehensive study of tRNA-derived fragments in plants for biotic stress responses

Supriya P. Swain¹ · Niyati Bisht¹ · Shailesh Kumar¹

Received: 12 January 2025 / Revised: 28 February 2025 / Accepted: 11 March 2025
© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2025

Abstract

Plant growth and development are often disrupted by biological stressors as they interfere with the regulatory pathways. Among the key regulators, transfer-RNA-derived fragments (tRFs) have emerged as key players in plant defense mechanisms. While tRF-mediated responses to abiotic stress have been well studied, their role in biotic stress remains less understood, as various stressors may elicit different regulatory systems. In this study, tRF-mediated biotic responses in three species, viz. *Arabidopsis thaliana*, *Oryza sativa*, and *Solanum lycopersicum* are investigated using *in-silico* approaches. Analysis of predicted tRFs across various biotic stress conditions reveals specific interactions with mRNA targets, microRNAs (miRNAs), and transposable elements (TEs), highlighting their regulatory significance in plant adaptation mechanisms. These findings provide new insights into tRF-mediated stress responses and establish a computational framework for further functional studies. The study's database is publicly available at <http://www.nipgr.ac.in/PbtRFdb>.

Keywords tRFs · tRNA fragments · TncRNAs · Biotic stress · Stress response · Genome regulation · Plant database

Introduction

The majority of the human genome encodes for non-coding RNAs (ncRNAs) which are now extensively studied for their crucial roles in regulating gene expression and genome organization (Nemeth et al. 2023). The endonucleolytic cleavage of precursor tRNA (pre-tRNA) or mature tRNAs, which constitute 4–10% of total cellular RNA, results in the formation of a pool of tRNA-derived fragments (Lee et al. 2009; Zhao et al. 2023). It constitutes the short fragments of length 14 to 30 nucleotides (nt) called tRNA derived fragments (tRFs or tDRs), longer tRNA halves (tRHs or tiRNAs) of approximately 30 to 50 nt (Sun et al. 2020; Zhu et al. 2019a, b) and other tRFs from the internal region of tRNAs. All the classes termed as tRNA-derived non-coding RNAs (tncRNAs) are conserved across most organisms.

These genome regulatory molecules are further classified based on the tRNA from which they are derived (S. Li et al.

2018). The tRFs derived from the 5' end are tRF-5, and the 3' end generates tRF-3, resulting from cleavage in the D and T regions, respectively (Ma et al. 2021). Additionally, pre-tRNA generates 5' U-tRFs (leader tRF) and tRF-1/tsRNA (30 U-tRFs) from the 5' leader and 3' trailer regions, respectively (Fig. 1) (Zahra et al. 2021). In the case of tRNA halves, 5' and 3' tRHs are produced from cleavage in the anticodon region, containing the 5' and 3' portions of the mature tRNA (Ma et al. 2021). Therefore, the identified tRFs are classified into tRF-5, tRF-3, tRF-1, leader-tRF, 5' tRH, 3' tRH, and a miscellaneous group labeled as other tRFs (Zahra et al. 2021).

The originating tRNA type, cell type, developmental stage, and stress conditions influence the specific cleavage of tRNA that generates these fragments. (Zahra et al. 2021). These tRFs can be produced in both DICER-dependent and -independent manner. RNase P eliminates the leader sequence and RNase Z cleaves the trailer sequence precisely at the discriminator base of the precursor-tRNA leading to the production of leader-tRF and tRF-1/tsRNA, respectively (Kim et al. 2020). Processing of certain types of tRNA halves requires both Dicer and angiogenin (Liu et al. 2018). The regulatory role of tRFs includes 1) regulation of mRNA stability as well as their potential to induce mRNA cleavage (Yu et al. 2021), 2) translation inhibition (Kumar et al.

Supriya P. Swain and Niyati Bisht contributed equally to this work.

✉ Shailesh Kumar
shailesh@nipgr.ac.in

¹ Bioinformatics Lab, National Institute of Plant Genome Research, Aruna Asaf Ali Marg, New Delhi 110067, India




Fig. 1 Classification and biogenesis of tRNA-derived fragments (tRFs). These fragments are generated through specific cleavage mechanisms involving enzymes such as DICER and angiogenin (ANG). tRFs are produced under various biotic conditions, includ-

ing those caused by nematodes, fungi, viruses, bacteria, and insects, which influence tRNA cleavage and tRF production. The resulting tRFs play diverse biological roles, such as association with RISC complexes, regulation of translation, and apoptosis

2016), 3) regulation of ribosome production (H. K. Kim et al. 2017), 4) control on apoptosis of cells depending on the diseases and target mRNA (Cui et al. 2022; Gong et al. 2023), and 5) facilitates amplification of viral propagation (Fu et al. 2015).

Exploration of tRFs in plants has been growing rapidly due to the exposure of plants to various biotic and abiotic stresses, which significantly impact their development (C. Wang et al. 2023). To manage these challenging conditions, plants have developed complex response mechanisms that involve multiple gene regulatory processes, including transcriptional (Chang et al. 2020) and post-transcriptional regulation (Jones-Rhoades et al. 2006). It is shown that tRFs in plants operate similarly to those in animals which could be due to their similarity in biogenesis, suggesting an evolutionary connection (Cao et al. 2022; Lalande et al. 2020). The studies on tRFs related to biotic stresses are relatively less developed compared to research on abiotic stresses (C. Wang et al. 2023). Nonetheless, the biological functionality of tRFs produced by biotic stress is better understood than that of abiotically induced tRFs (C. Wang et al. 2023). Their functional roles were thoroughly investigated in both the leaves and roots of black pepper affected by *Phytophthora*

capsici (Asha & Soniya 2016). In *A. thaliana*, *Botrytis cinerea* infection was seen to be significantly downregulating the 5' tRNA-derived small RNAs (Gu et al. 2022) whereas in *S. lycopersicum*, differentially expressed tRFs were characterized in plants subjected to *Tomato Mosaic Virus* (ToMV) (Zahra et al. 2021). Identification and characterization of tRFs and their specific possible targets in plants, particularly under biotic stress conditions, alongside an in-depth understanding of tRF-guided stress regulatory networks, can greatly contribute to enduring agricultural production. In this study, we identified a diverse range of tRFs across three angiosperms viz. *A. thaliana*, *O. sativa*, and *S. lycopersicum* using publicly available small RNA-seq (sRNA-seq) datasets of plants under various biotic stresses. Differential expression (DE) analysis revealed stress-specific tRFs, and common tRNA sources, followed by their mRNA targets, complemented by functional enrichment and pathway analysis. Analysis of RNA-Seq datasets further highlighted the DE patterns of mRNA targets, and interactions between tRFs and TEs, offering insights into their roles in stress adaptation. These findings were integrated into PbtRFdb, a comprehensive database with advanced search features, visualization tools, and dynamic graphs for

exploring tRFs under biotic stress conditions. An overview of each step followed during the study is shown in Fig. 2.

Methodology

Data retrieval and processing

A total of 112, 188, and 44 sRNA-seq samples of *A. thaliana*, *O. sativa*, and *S. lycopersicum*, sequenced under various biotic stress conditions, were downloaded from the NCBI SRA Toolkit (v3.0.2) (S. Sherry et al., 2023). Genome files for the respective plants viz. *A. thaliana* (TAIR10.1), *O. sativa japonica* (Build 4.0; organellar: IRGSP-1.0), and *S. lycopersicum* (nuclear: SL4.0; organellar: SL3.0), were obtained from EnsemblPlants (<https://plants.ensembl.org/index.html>) (Yates et al. 2022). To facilitate efficient secondary analysis, the headers for the corresponding plant genome files were substituted with “chr” followed by the chromosome numbers (Zahra et al. 2021). The raw FASTQ reads were processed by Trim Galore v0.6.6 (<https://github.com/FelixKrueger/TrimGalore>) to eliminate the low-quality bases and adapter contaminations (J. Wang et al. 2022). The FASTQ data quality

comparison was then performed with FastQC v.0.11.5 (<https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>).

tRF identification and filtration

For each sRNA-Seq sample, the tncRNA toolkit (<https://github.com/skbinfo/tncRNA-Toolkit.git>) was employed to identify the tRFs and their associated tRNAs (Zahra et al. 2023). The genomes were indexed using the integrated Bowtie v.1.3.0 tool in this toolkit, and this indexed genome was utilized to further locate the tRFs in the sequencing samples. The tRFs identified from each sample of each plant were concatenated, resulting in three merged files, one for each species. The tRFs with individual expression values of ≥ 10 per million reads were shortlisted (Alves et al. 2017). Unique sequences were subsequently identified, with the criterion that tRFs with identical sequences but differing modification sites were treated as distinct tRFs. The tRFs were named according to their respective species i.e. *A. thaliana* tRFs were designated as tRFs; *O. sativa* tRFs were labeled as OStRFs; and *S. lycopersicum* tRFs were referred to as SLtRFs (Table S1).




Fig. 2 Workflow for the Identification and Analysis of tRNA-derived Fragments (tRFs). The workflow outlines a systematic pipeline for identifying and analyzing tRFs under biotic conditions. Initial data-

sets consist of 112 samples across 13 biotic conditions for *Arabidopsis thaliana*, 188 samples across 8 biotic conditions for *Oryza sativa*, and 44 samples across 5 biotic conditions for *Solanum lycopersicum*

Differential expression of tRFs

A matrix was constructed in R, comprising tRFs and their corresponding read counts across the various samples (Table S1). This matrix was utilized as “countData” for DE analysis, with sample names and their descriptions designated as “colData”, as detailed in Table S2. For the identification of differentially expressed tRFs under various stress conditions, the DESeq2 R package was used (Love et al. 2014) with the R script mentioned in the Supplementary file 2. Only samples with corresponding treated and control pairs for each condition were considered. The threshold to identify the DE genes was an absolute fold change (\log_2FC) ≥ 1 or ≤ -1 , along with an adjusted P -value < 0.05 (Kim et al. 2024a, b) by using the Benjamini–Hochberg correction. The bar plots and heatmaps were created using the “ggplot2” package in R.

Target prediction and functional enrichment analysis

The highly, differentially expressed tRFs that were present in at least two infections were considered as queries for target prediction against the cDNA sequence of the considered plants. The tRFtarget-pipeline v0.3.0 was used to detect the respective transcripts for the tRFs at default parameters (<https://github.com/ZWang-Lab/tRFtarget-pipeline.git>) (N. Li et al. 2021). Moreover, it also facilitates the computational calculation of the binding energy between the tRFs and their target transcript (Tan et al. 2024). Based on minimum free energy (mfe) for each tRF, the top 10 targets were subjected to GO and pathway enrichment followed by further characterization. Functional enrichment of the target genes was done on g:profiler (<https://biit.cs.ut.ee/gprofiler/gost>) (Raudvere et al. 2019); Database for Annotation, Visualization, and Integrated Discovery (DAVID) (Dennis et al. 2003) (Saif et al. 2022); UniProt (<https://www.uniprot.org/uniprotkb/>) (Bateman et al. 2021), and KEGG (<https://www.genome.jp/kegg/>) (Kanehisa 2000). The significance threshold while using g: profiler was set to Benjamini–Hochberg FDR with a user threshold of 0.05 (Garay-Baquero et al. 2020). The DAVID analysis parameters were configured with an EASE score threshold of 0.1, the inclusion of FDR for result display, and a requirement for at least two hub genes in the analysis (Mukherjee & Sudandiradoss 2021). We employed a combination of databases for functional analysis since many tomato genes have not yet been annotated in g: profiler and DAVID. To address this, individual genes were further analyzed using UniProt which also helped in minimizing false positives. Only the most highly enriched terms across the databases were selected for downstream analysis (Saif et al. 2022).

RNA-Seq analysis and co-expression profiling of differentially expressed genes

The reference genome was indexed using HISAT2 (D. Kim et al. 2019), which was also utilized to align the filtered reads to the indexed genome. The aligned reads were assembled into potential transcripts using StringTie (Pertea et al. 2015), and the transcript abundance was quantified as FPKM values [total exon fragments/mapped reads (millions) \times exon length (kB)]. The number of mapped reads and transcript length in the samples were normalized to accurately reflect each transcript’s expression level and a gene count matrix consisting of raw read counts was constructed (Kim et al. 2024a, b). To evaluate whether the target genes were upregulated or downregulated during the infection, differential expression analysis was performed using DESeq2, employing the same parameters previously used for the DE tRF study. The volcano plot was generated using the SRplot web server (<https://www.bioinformatics.com.cn/srplot>) (Tang et al. 2023). The target genes that were downregulated in the samples were shortlisted and further underwent co-expression analysis. The Venn diagrams were generated using the jvenn web server (<https://jvenn.toulouse.inrae.fr/app/index.html>) (Bardou et al. 2014). The Pearson pairwise correlation analysis was conducted on the normalized data obtained from the differential expression study using the “corrplot” and “hclust” R packages. The threshold for the Pearson correlation coefficient was set at ≥ 0.9 between genes with a significance P -value of 0.05 to ensure that only highly co-expressed genes are considered (Pirone et al. 2023). The functional correlation of the co-expressed genes in *A. thaliana* and *O. sativa* was investigated using ATTED-II (<https://atted.jp/>) (Obayashi et al. 2022) and RiceFREND (<https://ricefrend.dna.affrc.go.jp/>) (Sato et al. 2013), respectively. Genes that were positively co-expressed with the target gene and also showed downregulation under the same infection condition in *S. lycopersicum* were selected for GO enrichment using UniProt. Cytoscape software version 3.10.2 was used to visualize and construct the various biological networks required for the study (Shannon et al. 2003).

Identification of potential transposable element targets

Details of Transposable Elements for the respective organism were obtained from the APTedB website (<http://apte.cp.utfrpr.edu.br/>) (Pedro et al. 2021). Only known and characterized TEs were retained for further analysis. To determine the DE status of the genes, their coordinates were cross-referenced with the DE data. The interaction of the tRFs with the TE was visualized using tRFtarget-pipeline v0.3.0. To categorize the classes of TEs involved in interactions

with tRFs, a chord diagram was created using the "circlize" package in R. (H. Wang et al. 2019).

Detection and verification of miRNAs for the target mRNA

The top 10 targets were shortlisted based on 'mfe' of each tRF and were also examined for potential miRNA interactions. The miRNA information for *A. thaliana* and *O. sativa* was sourced from the PMRD database (<https://bioinformatics.cau.edu.cn/PMRD/>) (Z. Zhang et al. 2010) while data for *S. lycopersicum* was obtained from the TarDB database (<http://www.biosequencing.cn/TarDB/>) (Liu et al. 2021a, b) respectively. After conducting the mRNA-miRNA interaction study using the tRFtarget pipeline v0.3.0, the interactions between tRFs and miRNAs with their respective target genes were compared based on binding energy and the location of binding on the mRNA. Subsequently, the presence of the miRNA in the sample was detected using the BrumiR toolkit (<https://github.com/camoragaq/BrumiR>) (Moraga et al. 2022).

Database development

The data analysis was compiled into a database called "PbtRFdb". Web development for this database was carried out using the XAMPP package (v.8.0.30) (<https://www.apachefriends.org/>), which facilitates the integration of Apache, MySQL, and PHP in a local server environment. The front-end web interface was designed using HTML, CSS, and JavaScript. MySQL handled data management and processing, providing a robust platform for storing and efficiently retrieving data. The PHP scripts interacted with MySQL to deliver query results to the user.

Results

Comprehensive profiling and differential expression of tRFs under biotic stress conditions across three plants

Using the tncRNA-toolkit, a total of 277,183 tRFs were identified in *A. thaliana*, 268,032 in *O. sativa*, and 92,263 in *S. lycopersicum*. After an initial filtration based on individual expression level (Fig. S1), and retaining the top 50% of tRFs, a final unique set of 2,505 tRFs for *A. thaliana*, 3,078 for *O. sativa*, and 843 for *S. lycopersicum* was obtained. Based on individual expression levels, other-tRFs categories were more abundant (Table S1). The number of tRFs remaining after each filtration step is shown in Fig. 2. We analyzed the expression of these tRFs under various biotic stress conditions in plants, revealing distinct variations in

both expression patterns and the types of tRFs across different groups of biotic stress infections (3), and in response to each specific infection (Fig. S2). A significant number of tRFs, across various types, were predominantly upregulated in all the plants (Fig. S3). The classes tRF-5, other-tRFs and 5' tRH showed significant differential expression. Alongside the stress response, we also observed tissue-specific variations in tRFs expression. Notably, tRF-5 was highly upregulated and downregulated in leaf tissue at different developmental stages under various infections (Fig. S4). The majority of the tRF-5 fragments were found in the developmental stages ranging from mature leaf (21–30 days old) to young leaf (15 days). Additionally, other tRFs were found to be DE across major tissue types under prominent biotic stress conditions, including various developmental stages of leaves and in 15-day-old roots of *O. sativa*.

GluTTC exhibits maximum tRNA cleavage in response to biotic stress

All the DE tRFs were analyzed for the tRNA origin information. It has been found that the GluTTC generated the highest number of tRFs across various biotic stress conditions in all studied plants (Fig. S5). Different isoacceptors produced various types of tRFs and showed differences between plants (Zahra et al. 2021). Specifically, the isoacceptors of tRNA-Glu i.e., TTC and CTC, contributed to the majority of DE tRF production across all plants, with GluTTC producing the highest amount of tRF-5 in *A. thaliana*, *O. sativa*, and *S. lycopersicum*, respectively (Fig. S6). Other tRFs were also observed to be produced from Glu-TTC in both *A. thaliana* and *S. lycopersicum*. Various anticodons contributed to the production of tRF-5 across all plants. For example, in *A. thaliana*, tRF-5 was also generated by GlyTCC, AspGTC, GlyGCC, and MetCAT, among others. To assess the impact of tRFs generated from GluTTC, their mRNA targets were identified and shortlisted based on binding energy. The identified targets were then subjected to GO enrichment analysis, revealing a strong enrichment for the terms "response to heat" and "cellular component organization" in *A. thaliana* (Fig. S7). These terms fall under the broader categories of "response to stimulus" and "cellular process". The targets of tRFs generated by GluTTC in *O. sativa* were found to play a role in "galactose metabolism", suggesting that these tRFs may regulate genes involved in this metabolic pathway. None of the genes showed enrichment in *S. lycopersicum*.

Enrichment of mRNA targets in distinct stress response mechanisms

tRFs may function as post-transcriptional regulators by binding complementarily to their mRNA targets, leading to the degradation of these mRNAs (Alves et al. 2017; Kusc

et al. 2018). To explore the functional roles of tRFs, we conducted target predictions (Fig. 3, Table S3) and pathway enrichment analyses, for tRFs expressed in at least two infectious conditions in plants. Our analysis revealed that the majority of target genes were associated with responses to various stress types, including cold stress (abiotic) and bacterial stress (biotic) (Fig. S8). The cellular mechanisms activated to cope with stressors, such as "signal transduction", "transport", and various "metabolic processes", were found to be enriched across all plant species analyzed. Within the transport processes, "microtubule-based movement" was particularly prominent. Among biological processes, the term "biosynthetic process" was observed to be the most highly enriched across the plants. Under molecular functions, "catalytic activity" was extensively abundant. Consistent with the findings of Zahra et al. 2021, our tRF targets were also prominently associated with "transferase" and "kinase" activities. Additionally, "binding" was highly enriched in *A. thaliana* and *O. sativa*. "ATP binding" was significantly enriched and the most prevalent term, while "microtubule binding" exhibited the highest level of significance. In *A. thaliana* and *O. sativa*, the term "membrane" was highly and significantly enriched, whereas in *S. lycopersicum*, the term "nucleus" was extensively enriched. These

findings suggest distinct functional adaptations among plant species, reflecting their specific stress response mechanisms and highlighting the role of tRFs in regulating various biological processes (Fig. S9).

Identification of differentially expressed mRNA targets

RNA-seq analysis followed by DE gene identification was conducted to examine the expression of target genes in the samples under biotic stress conditions. This approach also aimed to assess the potential influence of tRFs on regulating these genes within the samples. In *O. sativa* and *S. lycopersicum*, most of the genes were observed to be downregulated, whereas, in *A. thaliana*, the majority of the genes were predominantly upregulated (Fig. 4). No common genes were identified across all conditions in all plant species; however, overlapping genes in at least two conditions are evident in Fig. S10. Most of the 76 genes in *A. thaliana* were found to overlap between those upregulated during *Tobacco Mosaic Virus* (TMV) infection and those downregulated in response to *Turnip Crinkle Virus* (TCV). A similar trend was observed in *O. sativa* and *S. lycopersicum*, where gene overlaps were detected in at least two infection conditions. In *O. sativa*,




Fig. 3 mRNA targets of the most recurrent tRFs identified through tRFtarget-pipeline v0.3.0 (<https://github.com/ZWang-Lab/tRFtarget-pipeline.git>). The figure highlights the top three mRNA targets for

each species with their minimum free energy (mfe): **a** *Arabidopsis thaliana*, **b** *Oryza sativa*, and **c** *Solanum lycopersicum*




Fig. 4 Volcano plots showing the gene expression in response to biotic stress conditions. The plots illustrate the upregulation and downregulation of genes for each species: **a** *Arabidopsis thaliana*, **b** *Oryza sativa*, and **c** *Solanum lycopersicum*

11 genes predominantly overlapped between those downregulated by *Brown Planthopper* (BP) infection and *Xanthomonas oryzae* (XO) infection. Meanwhile, in *S. lycopersicum*, a substantial overlap of 150 genes was noted between those downregulated during *Meloidogyne incognita* (MI) infection and those upregulated in response to *ToMV*. In *A. thaliana*, the overlapping genes were significantly enriched in pathways related to "intracellular protein transport," "ARF protein signal transduction," and "post-transcriptional gene silencing" (Fig. S11). These terms generally fall under the category of stress response. In *O. sativa*, "monoterpenoid biosynthesis" was prominently featured, while in *S. lycopersicum*, the "ubiquitin conjugation pathway" was notably highlighted which also indicated the stress-related process.

Upregulated tRFs may contribute to the downregulation of target mRNAs and co-expressed genes, thereby influencing shared biological pathways

We observed that the majority of shortlisted mRNA targets were downregulated during DE analysis. With the hypothesis that the upregulated tRF could be a contributing factor to the downregulation of its target mRNA (Zong et al. 2021), we selected the downregulated target mRNAs from the list of DE genes in each plant species. The co-expression analysis revealed that genes co-expressed with the target gene were also downregulated in response to the infection. AT2G39950, a flocculation protein and a potential target mRNA for tRF1492 in *A. thaliana*, was significantly downregulated during infection with *Alternaria brassicicola* (AB), showing a log2FC value of -1.66 . Additionally, its co-expressed gene AT5G60170 was also found to be downregulated during the same infection, showing a log2FC value of -1.05 (Table S4). The functional analysis revealed

that genes AT5G60170 and AT2G28540 both belong to the RNA-binding protein family. At the same time, AT3G22270 is associated with topoisomerase II which may facilitate mRNA decay and enhance post-transcriptional reprogramming (<https://www.arabidopsis.org/locus?name=AT3G22270>) (Fig. 5a). In *O. sativa*, the gene OsCesA5 (Os03g0837100), which encodes a cellulose synthase A5 protein, was significantly downregulated during the BP infection with a notable negative log2FC value of -3.075 . This gene is targeted by OstRF1920 and OstRF1928, which were significantly upregulated during the same infection. All the co-expressed genes, including CESA6 (Os07g0252400), OsGT47A (Os01g0926600), and OsFLA3 (Os08g0321000), were significantly downregulated during BP infection; however, the Os07g0633600 gene did not show any DE in the samples. The CESA6 and OsGT47A are involved in "plant-type cell wall biogenesis" and "polysaccharide biosynthetic process" (Fig. 5b). These genes along with the target mRNA, OsCesA5, play a role in the biosynthesis and modification of the plant cell wall. The other co-expressed gene sets for the mRNA targets OsWD40-125 (Os06g0143900) and Os09g0283600 did not show DE under any of the infectious conditions. In *S. lycopersicum*, the majority of significant co-expression was observed for the target gene *Solyc02g080340.3* with SLtRF725 and SLtRF26 under *ToMV* infection, specifically in processes related to "fatty acid metabolism and lipid degradation" and "defense response to other organisms". The target gene was downregulated with a log2FC value of -0.96 , while the tRFs were upregulated with log2FC values of 28.054 and 36.855, respectively, during the infection. All the co-expressed genes were downregulated under *ToMV* stress. Among them, the *Solyc01g104210.2* gene is also involved in "fatty acid metabolism (elongation)" and the "sphingolipid biosynthetic process" (Fig. 5c). Two of the co-expressed




Fig. 5 Flowchart illustrating the co-expression of mRNA targets with other genes and their functional annotation for each species: **a** *Arabidopsis thaliana*, **b** *Oryza sativa*, and **c** *Solanum lycopersicum*

genes, *Solyc06g082340.2* and *Solyc12g056130.2*, are involved in ubiquitin-dependent pathways, specifically the ubiquitin-dependent protein catabolic process and the ubiquitin ligase conjugation pathway, respectively. In contrast, most of the remaining genes were not found to be significantly enriched. Moreover, the gene *Solyc06g048950.3*, which encodes a non-specific serine/threonine protein kinase (STPK), was significantly downregulated during infection by the *Alternaria* fungus (AF). This gene showed considerable co-expression with a retrotransposon-derived protein that was also downregulated under the same infection. Additionally, the target gene *Solyc01g106770.3*, another STPK-TOR, which is targeted by the upregulated SLTrF723, was downregulated under ToMV infection, along with its co-expressed gene *Solyc01g008740.2*, a calcium-dependent kinase involved in intracellular signal transduction.

miRNAs did not influence target mRNA expression

Fig. S12a illustrates that several mRNA transcripts in *A. thaliana* have the potential to be regulated by specific miRNAs capable of binding to them. In contrast, in *O. sativa*, two genes were identified to bind to miRNAs, as depicted in Fig. S12b potentially. No miRNAs were found to potentially bind to any predicted target genes in *S. lycopersicum*. A comparison of the binding energies and binding coordinates between miRNA and tRF reveals that, although both exhibited highly negative binding energies, the binding coordinates on the target gene differed (Table 1). Additionally, these mRNA targets did not show DE under any infection conditions in either *A. thaliana* or *O. sativa*. Furthermore, none of these miRNAs were detected in the samples considered in this study. Moreover, this supports the previous

hypothesis that the downregulated potential target mRNA and its corresponding co-expressed downregulated mRNA may be DE primarily due to their interaction with the tRFs, which are upregulated during various biotic stresses.

tRFs regulate the activity of the transposable elements

The chord diagram in Fig. 6 collectively demonstrates that most tRFs across all plant species exhibited a strong affinity for binding to LTR/gypsy TE, supporting the findings of (Zahra et al. 2021). In *A. thaliana*, tRF-5 was predominantly associated with LTR/gypsy elements, while tRF-3-CCA primarily interacted with SINE/Unknown elements (Table S5). It was observed that TE transcripts were predominantly downregulated during TCV infection, with tRF-5 and 5'tRH showing strong interactions with TE transcripts in *O. sativa*. Notably, tRF-5 mainly interacted with Helitron/Helitron elements, while 5'tRH was primarily associated with TIR/Dada elements. Significant downregulation of TE transcripts was observed during BP infection, with some downregulation also occurring in XO and *Meloidogyne graminicola* (MG) infections. In *S. lycopersicum*, most tRF-5 s and other-tRFs showed a preference for interacting with LTR/gypsy elements, with some interaction observed with LTR/Copia and LTR/Unknown elements. The strongest downregulation of TE transcripts was observed during AF infection, with notable downregulation also seen in MI and ToMV infections. This supports the conclusion highlighted in the review article of (Y. Zhang et al. 2024), that tsRNA can regulate cellular responses by affecting the activity of retrotransposons. It was observed that tRNAMetCAT generates

Table 1 Comparative analysis of miRNA and tRF targets in *Arabidopsis thaliana* and *Oryza sativa*

Species	Target	miRNA	SubseqDP	BE miRNA	Start target	End target	tRF ID	SubseqDP	BE tRF	Start target	End target
<i>Arabidopsis thaliana</i>	AT3G10290.1	ath-miRf10578-akr	AUAUAAGCG AAGAGUGCA GU&ACUGCA ACUGCUUGUAU	-23.2	1413	1432	tRF1319	UCCUUUAUCAA GCCGUUGGU GCUACGACG C&GGCUCUGUA GUCCAACGG UUAGGA	-36.8	451	481
	AT5G58190.2	ath-miRf10373-akr	AGCUCUUCU UUGGGAGGU UA&UGACCU GCCAAAGGA GAGUU	-34.3	1474	1493	tRF365	UGGCGUCCAACA GCAGG&CCU GCUGUUGAC GCCA	-33.3	81	97
	AT1G05580.1	ath-miR414	GCGGUGAUGGUC UGAA&UUCAUC AUAUCGU	-20.6	1154	1169	tRF1319	GUCCUAUGCGGC CUGAUUACG GACGC&GCG UCUGUAGUC CAACGGUUA GGAU	-38.5	874	899
	AT1G05580.2	ath-miR414	GCGGUGAUGGUC UGAA&UUCAUC AUAUCGU	-20.6	1077	1092	tRF1319	GUCCUAUGCGGC CUGAUUACG GACGC&GCG UCUGUAGUC CAACGGUUA GGAU	-38.5	874	899
<i>Arabidopsis thaliana</i>	AT1G72120.1	ath-miRf11033-akr	GAUGUUGAA GAUGCCACA GC&GUGUG GCAUCAUCA AGAUU	-32.8	1212	1231	tRF1492	UUGAGCUAGAGG CAAAUACAU CCCC&GGGGAU GUAGCUCAA	-28.5	821	845
	AT3G04200.1	ath-miR163	AUCGAAGGUUC GAUCCAAA ACCGCGUCU GCAA&UUGAAG AGGACUUGG AACUUCGAU	-21.5	217	250	tRF1319	AUCCUGGUACCA UUAGGAUUG CAGACGC&GCG UCUGUAGUC CAACGGUUA GGAU	-38.5	545	572
<i>Arabidopsis thaliana</i>	AT1G56080.1	ath-miRf10720-akr	UCCGCGUAUAC CGCGG&CCC GCGGUUAC CACGGG	-37.7	111	128	tRF1492	GAGCUGCAUCUC C&GGGGAUGUA GCUC	-29.4	614	626

Table 1 (continued)

Species	Target	miRNA	SubseqDP	BE miRNA	Start target	End target	tRF ID	SubseqDP	BE tRF	Start target	End target
<i>Oryza sativa</i>	Os05g0590000	osa-miR2103	UGGGUGGCGCAGG GCCUUGGAG	-36.8	1526	1547	OstRF1274	GCGACGUUCUAC UACGUGCAC	-45.4	961	1002
			AAGGCGGCG					CUCCUGGGC			
			GGGAGAG&UUU CCCUCCCG UGC GGCUC					UUCAGCGUG GGC&GCCAC UUAGCUCAG AGGUUAGAG CAUCGC			
	Os05g0590000	osa-miR2103	UGGGUGGCGCAGG GCCUUGGAG	-36.8	1526	1547	OstRF1530	CCUACGCCGCGG UGGACUUGCG	-33.2	530	556
			AAGGCGGCG					UCGCGC&GCG			
			GGGAGAG&UUU CCCUCCCG UGC GGCUC					UUUGUAGUC CAACGGUUAGG			
	Os06g0611000	osa-miRf11622-akr	GGCUGCUCGCGU GAA&UGCCAU	-30.2	4090	4112	OstRF2652	GACCAGCUA GCUGCAGAG	-34.1	1060	1082
			CACCGGUAG					CAGGA&UCC			
			CAGCACUU					GUUGUAGUC UAGCUGGUU			

The table displays target gene identifiers, miRNA and tRF ID, and the subsequence of duplex pairing (SubseqDP), their binding energy (BE), and start and end target positions within the gene




Fig. 6 Chord diagram shows that the majority of tRFs across all plant species display a strong binding affinity to transposable elements, LTR/Gypsy TEs

tRF-s which specifically targets the LTR retrotransposon, Athila6A (Martinez et al. 2017). Since both tRFs and TEs are known to be expressed and accumulate under stress, it would seem that the regulation of tRF production is connected to TE activity.

PbtRF database

A total of 637,476 entries are stored in PbtRFdb, comprising 277,181 entries for *A. thaliana*, 268,032 for *O. sativa*, and 92,263 for *S. lycopersicum*. For differentially expressed

tRFs, *A. thaliana* has 2,260 entries, *O. sativa* has 6,663, and *S. lycopersicum* has 901 entries. Detailed information on tRFs can be retrieved through two main modules: the “Search” and “Browse” modules. The Search module allows users to extract detailed information based on sequences, tRF types, amino acids, anticodons, modifications, and differentially expressed tRFs. The “DE tRFs” section enables users to query biotic stress-specific studies across the three species, with filters for various stress conditions and expression levels. The Browse module allows users to extract information across the three species via four categories: tissue type, amino acid, anticodon, and stress. Each query provides users with specific details such as SRA accession, organism, tissue, developmental stage, tRF type, locus, position, length, stress, sequence, and modification. The query also provides log2fold change data and expression information for differential expression, including upregulation and down-regulation. Each SRA accession is hyperlinked to the NCBI SRA page, offering additional information. Users can visualize tRFs using the “JBrowse” module for the species of interest. The module allows users to select tracks, including the reference sequence, general feature file track (GFF), and tRF track. Users can zoom in on specific locations to view tRFs and gene features. The tRF track includes sequence information, tRF type, tRNA derivatives, and the length of the tRF. PbtRFdb also includes a “Statistics” module, which allows users to visualize tRFs across various stress conditions. The most abundant classes of tRFs and parental tRNAs under different conditions are displayed using graphs. DE tRFs are presented with user-friendly visualizations such as line graphs, bar plots, and pie charts. Additional features include a “Download” module, allowing users to download results for each plant species separately. A “Help” module is also provided to guide users through navigating the database.

Discussion

tRNA-derived fragments are emerging as essential regulators of plant defense responses and stress-related adaptations. However, their functional roles in biotic stress responses remain underexplored. In this study, we address this gap by using computational methods to comprehensively profile tRFs across the plant species—*A. thaliana*, *O. sativa*, and *S. lycopersicum* under biotic stress conditions. Furthermore, we studied the regulatory networks of the majority of recurring tRFs, by examining how they interact with mRNA targets, miRNA, and TEs to gain a better understanding of their functional importance in plant stress response.

A significant number of other-tRFs amongst the tRF subclasses may indicate that less-defined tRFs are important mediators of plant responses to different biotic stresses. The findings of Thompson et al. 2008, showcased that oxidative

stress causes the endonucleolytic cleavage of tRNAs in plants, was consistent with the significant differential expressions shown in the tRF-5, other-tRFs, and 5'tRH classes. Likewise, Alves et al. 2017 emphasized how tRF-5 is linked to certain plant AGOs, promoting its synthesis under abiotic stress. Thus, it is reasonable to assume that during biotic stress conditions, tRF-5 actively participates in post-transcriptional gene regulation. Plant immunity is based on the different pathways that it may activate during infection.

In our study, for differential expressions we considered biotrophic pathogens (*Xanthomonas oryzae*, *Heterodera schachtii*, *African Xanthomonas*, *Meloidogyne graminicola*, *Meloidogyne incognita*) (Galeng-Lawilao et al. 2018; Haidar et al. 2016; Panthapulakkal Narayanan et al. 2020), necrotrophic pathogens (*Botrytis cinerea*, *Sclerotinia sclerotiorum*, *Alternaria brassicicola*, *Fusarium oxysporum*) (Cho 2015; Pandey et al. 2016; Panthapulakkal Narayanan et al. 2020; Petrasch et al. 2019) and hemibiotrophs (*Phytophthora capsici*, *Magnaporthe oryzae*) (Barraza et al. 2022; Panthapulakkal Narayanan et al. 2020). Biotrophic pathogens activate salicylic acid (SA)-dependent immunity (Yang et al. 2015) and necrotrophic pathogens activate jasmonic acid (JA)- and ethylene (ET)-dependent immunity (Macioszek et al. 2023; Rahman et al. 2012; Sanchez et al. 2012). NRP1 is a gene that regulates salicylic acid-mediated immunity in plants (Mishra et al. 2024). In one of the studies, cleavage mediated by the 5'Ala^{CGC} tRF was experimentally mapped at the tRF binding sites on the mRNA targets of non-expresser of pathogenesis-related protein (NPR1), which showed a down-regulation during *Phytophthora capsici* infection (Asha & Soniya 2016). Another study demonstrates that 5' tRNA-Ala-derived tRFs regulate the defense against fungal responses, possibly due to the upregulation of jasmonic acid-mediated immunity (Gu et al. 2022). These pathways can function both synergistically and antagonistically, adding complexity to plant immune responses (Ghazlan et al. 2020; N. Li et al. 2019; Mur et al. 2006). Our results highlighted that tRF-5, tRF-3-CCA, and other-tRFs were major subclasses that were differentially expressed in all the infections suggesting that they are involved in the specific adjustments of plant immunity (Figure S1). As viruses are obligate biotrophs, their relationship with SA-dependent defense may influence tRF biogenesis. The increased levels of specific tRFs in *A. thaliana* as a response to bacterial, fungal, and viral infections could indicate a potential activation of SA-mediated immunity. In *O. sativa*, the decrease of specific tRFs in fungi and virus infections might suggest a suppression or diminished activation of JA/ET-mediated immunity. In *S. lycopersicum*, the increase of specific tRFs during fungal and viral infections could indicate a response similar to that seen in *A. thaliana*. Gaining insight into the responses of tRF subclasses to biotrophic and necrotrophic pathogens within these different immune networks may help

us understand their possible regulatory functions in plant defense.

The regulatory role of tRF-5 is evident, as a large amount of tRF-5 is cleaved from tRNA-GluTTC across all species (Figure S6). tRNA-GluTTC translates to the glutamate (Glu) amino acid, which is a key metabolite in stress-related pathways such as signaling and metabolic regulation. Therefore, tRNA-GluTTC is highly abundant during abiotic stress across species (Qiu et al. 2020; Zahra et al. 2021). tRNA-GluTTC serves as a stress-responsive biomarker in human gastric and thyroid cancers (Mao et al. 2024; Shan et al. 2021; Zhu et al. 2019a, b). Using GO enrichment analysis, we derived some functional relevance of tRFs cleaved from tRNA-GluTTC. In *A. thaliana*, the tRFs were linked with UDP glucose 4-epimerase activity (essential for cell wall biosynthesis), ATP-dependent chromatin remodeling activity (related to plant development and stress response), and cytoplasmic stress granule (early stress response), all of which contribute to stress adaptation (Guo & He 2024; Maruri-López et al. 2021; Reuper et al. 2021; Rösti et al. 2007) (Figure S7a). In *O. sativa*, tRFs cleaved from tRNA-GluTTC were associated with galactose metabolism (Figure S7b), a process that plays a role in cellular homeostasis during stress conditions (Wang et al. 2024; Höftberger et al. 2022; Liu et al. 2021a, b). However, no significant enrichment patterns were observed for *S. lycopersicum*, possibly due to the limited functional annotation data. This indicates that tRFs derived from tRNA-GluTTC may play an active role in responses to biotic stress by regulating metabolic pathways through gene expression or interactions with proteins that respond to stress, as part of a larger plant defense system (Yagyū & Yoshimoto 2024).

Apart from their function in the metabolic process, tRFs also affect gene expression via interacting with splicing factors and mRNA regulation (George et al. 2022; Y. Li et al. 2023). Differential gene expression patterns were observed across species in response to biotic stress. During *Alternaria brassicicola* infection, the target mRNA and co-expressed genes were downregulated in *A. thaliana*, whereas during *Brown planthopper* infestation, the target mRNA co-expressed genes showed a comparable downregulation in *O. sativa*. The target mRNA and co-expressed genes in *S. lycopersicum* were upregulated after *ToMV* infection and downregulated in *Meloidogyne incognita*. The upregulation of the target genes under specific conditions may enhance stress responses, by increasing the metabolism activity and increasing the production of stress-related proteins (Hillwig et al. 2008). Conversely, the downregulation of target genes may suppress certain stress responses therefore increasing the susceptibility to pathogens (Krishna et al. 2021; Park & Kim 2018). The regulatory patterns observed indicate that tRFs might play a role in stress adaptation by modulating mRNA

expression and the co-expressed gene network, possibly via RNA interference or splicing modulation. Further experimental validation is, however, required to understand gene regulation mediated by tRFs and its conservation among species.

Considering the established function of small RNAs in gene regulation, it is crucial to observe how tRFs compare or interact with other small RNA pathways, like miRNA-mediated regulation. Mechanisms like translation repression of mRNAs and their degradation have defined the role of miRNAs in regulating mRNA stability (Dalmay 2013). In *A. thaliana*, studies reveal miRNAs play a role in post-transcriptional regulation by targeting the mRNAs of transcription factors (Bartel 2004). The co-expression of the target genes and miRNA has been linked to various metabolic processes and signaling systems. For instance, ath-miR414 participates in the phosphatidylinositol signaling system (Xie et al. 2015), ath-miR163 is potentially linked to adaptations in nutrient-deficient environments (Fasani et al. 2021), and osa-miR2103 was found to play a critical role in rice immunity against *Magnaporthe oryzae* (Fan et al. 2020). The co-expression of tRFs and miRNAs hypothesizes that tRFs can function in miRNA canonical pathways by competing with the AGO association acting as global regulators of RNA-mediated silencing (Alves et al. 2017; Alves & Nogueira 2021). However, despite the well-documented role of miRNAs in gene regulation, our study did not detect miRNAs in the analyzed samples. The absence of detectable miRNAs in our datasets may be due to a reduced count of highly abundant miRNAs. We can hypothesize the possibility that tRFs might serve as alternative regulators of mRNA stability and gene expression. Since tRFs are known to interact with AGO proteins and affect translation in a manner comparable to miRNAs, they could play a role in mRNA silencing via different or independent mechanisms. It is necessary to conduct further research to establish whether tRFs and miRNAs work together in post-transcriptional regulation or if tRFs function as independent regulators under certain stress conditions.

Given the ability of tRFs to regulate gene expression through the interactions with mRNAs, and miRNAs, it is important to extend its regulatory to non-coding elements such as TEs. The interaction between tRFs and TEs has been explored in only a few studies across both plants and mammals, yet their functional relationship remains unclear (Martinez 2018; Slotkin et al. 2009; Zahra et al. 2021). TEs engage with transcripts, modifying expression levels through their role as new promoters or by affecting regulatory pathways (Hirsch & Springer 2017). Our results showed various TE families among species, with some directly associated with stress responses and adaptations to challenging environments (Bolger et al. 2014; Pietzenuk et al. 2016). The findings suggest that tRFs could play a role in the regulation

of TEs, potentially by influencing their expression or affecting their function as regulatory elements.

We also developed a database to catalog and analyze tRF profiles across *A. thaliana*, *O. sativa*, and *S. lycopersicum*, which allows for detailed investigations of tRFs under different biotic stresses. This resource offers a foundation for future investigations into tRF-mediated regulatory mechanisms during biotic stress conditions.

The insights gained from our *in-silico* analysis provided an emphasis on the regulatory roles of tRNA-derived fragments in plants under biotic stress. The identification of distinct tRF profiles and their potential interactions with mRNA targets, miRNAs, and transposable elements highlights the complexity and specificity of stress response pathways across various plant species. The results emphasize the possible function of tRFs in cellular adaption under biotic stress. To verify these interactions and clarify the molecular pathways by which tRFs aid in cellular adaptation to biotic stress conditions, additional experimental validation and thorough mechanistic investigations are necessary.

Conclusion

tRFs are small non-coding RNAs that can act as novel regulators, impacting transcriptional and post-transcriptional processes (Martinez et al. 2017). Several studies have characterized tRFs in viruses, fungi, and plants (Asha & Soniya 2016; Gu et al. 2022; Gupta et al. 2018; Zahra et al. 2021, 2022), making it a critical regulators of stress mechanisms in plants. Our study emphasizes the function of tRFs in biotic stress across three species- *A. thaliana*, *O. sativa*, and *S. lycopersicum* as well as their regulatory network with mRNA, microRNAs, and transposable elements.

Notably, our findings reveal that tRNA-GluTTC is extensively cleaved in response to biotic stress and plays a unique role in stress-response pathways. Most recurrent tRFs show interaction with mRNA targets essential for the plant adaptation to stress, indicating that differential tRF expression may influence mRNA regulation. Additionally, post-transcriptional gene silencing may be facilitated by tRF-mediated stress regulation mechanisms that function independently of miRNA pathways. A complex regulatory network affecting stress adaption may result from the interaction of tRFs and TEs. This study's database is accessible at <http://www.nipgr.ac.in/PbtRFdb> and serves as a valuable resource for future studies on tRFs in plant biotic stress response and provides a background for further experimental studies for their regulatory role.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s10142-025-01576-3>.

Acknowledgements The authors are thankful to the Department of Biotechnology (DBT)-eLibrary Consortium, India, for providing access to e-resources. SPS and NB are thankful to the University Grants Commission (UGC), India, and the Department of Biotechnology (DBT), India, respectively, for research fellowships. The authors acknowledge the Computational Biology & Bioinformatics Facility (CBBF) of the National Institute of Plant Genome Research (NIPGR).

Author contributions Data curation: SPS and NB.

Formal analysis: SPS and NB.

Web-interface development: NB.

Funding acquisition, Project administration, Resources, and Supervision: SK.

Roles/Writing—Original draft: SPS and NB; and Writing—review & editing: SK.

Funding This research is supported by the BT/PR40146/BTIS/137/4/2020, BT/PR40169/BTIS/137/71/2023, BT/PR40160/BTIS/137/64/2023, BT/PR40261/BTIS/137/55/2023 project grants by Department of Biotechnology (DBT), Government of India and by the core grant of the National Institute of Plant Genome Research (NIPGR) in the laboratory of SK.

Data availability No datasets were generated or analysed during the current study.

Declarations

Competing interests The authors declare no competing interests.

References

- Alves CS, Nogueira FTS (2021) Plant small RNA world growing bigger: tRNA-derived fragments, longstanding players in regulatory processes. *Front Mol Biosci* 8:638911. <https://doi.org/10.3389/fmolb.2021.638911>
- Alves CS, Vicentini R, Duarte GT, Pinoti VF, Vincentz M, Nogueira FTS (2017) Genome-wide identification and characterization of tRNA-derived RNA fragments in land plants. *Plant Mol Biol* 93(1–2):35–48. <https://doi.org/10.1007/s11103-016-0545-9>
- Asha S, Soniya EV (2016) Transfer RNA derived small RNAs targeting defense responsive genes are induced during phytophthora capsici infection in black pepper (*Piper nigrum* L.). *Front Plant Sci* 7. <https://doi.org/10.3389/fpls.2016.00767>
- Bardou P, Mariette J, Escudé F, Djemiel C, Klopp C (2014) jvenn: an interactive Venn diagram viewer. *BMC Bioinforma* 15(1):293. <https://doi.org/10.1186/1471-2105-15-293>
- Barraza A, Núñez-Pastrana R, Loera-Muro A, Castellanos T, Aguilar-Martínez CJ, Sánchez-Sotelo IS, Caamal-Chan MG (2022) Elicitor Induced JA-Signaling Genes Are Associated with Partial Tolerance to Hemibiotrophic Pathogen *Phytophthora capsici* in *Capsicum chinense*. *Agronomy* 12(7):1637. <https://doi.org/10.3390/agronomy12071637>
- Bartel DP (2004) MicroRNAs. *Cell* 116(2):281–297. [https://doi.org/10.1016/S0092-8674\(04\)00045-5](https://doi.org/10.1016/S0092-8674(04)00045-5)
- Bateman A, Martin M-J, Orchard S, Magrane M, Agivetova R, Ahmad S, Alpi E, Bowler-Barnett EH, Britto R, Bursteinas B, Bye-A-Jee H, Coetzee R, Cukura A, Da Silva A, Denny P, Dogan T, Ebenezer T, Fan J, Castro LG, ..., Teodoro D (2021) UniProt: the universal protein knowledgebase in 2021. *Nucl Acids Res* 49(D1):D480–D489. <https://doi.org/10.1093/nar/gkaa1100>
- Bolger A, Scossa F, Bolger ME, Lanz C, Maumus F, Tohge T, Quesneville H, Alseekh S, Sørensen I, Lichtenstein G, Fich EA, Conte

- M, Keller H, Schneeberger K, Schwacke R, Ofner I, Vrebalov J, Xu Y, Osorio S, ... Fernie AR (2014) The genome of the stress-tolerant wild tomato species *Solanum pennellii*. *Nat Genet* 46(9):1034–1038. <https://doi.org/10.1038/ng.3046>
- Cao K-Y, Yan T-M, Zhang J-Z, Chan T-F, Li J, Li C, Lai-Han Leung E, Gao J, Zhang B-X, Jiang Z-H (2022) A tRNA-derived fragment from Chinese yew suppresses ovarian cancer growth via targeting TRPA1. *Mol Ther Nucleic Acids* 27:718–732. <https://doi.org/10.1016/j.omtn.2021.12.037>
- Chang Y, Zhu C, Jiang J, Zhang H, Zhu J, Duan C (2020) Epigenetic regulation in plant abiotic stress responses. *J Integr Plant Biol* 62(5):563–580. <https://doi.org/10.1111/jipb.12901>
- Cho Y (2015) How the Necrotrophic Fungus *Alternaria brassicicola* Kills Plant Cells Remains an Enigma. *Eukaryot Cell* 14(4):335–344. <https://doi.org/10.1128/EC.00226-14>
- Cui H, Li H, Wu H, Du F, Xie X, Zeng S, Zhang Z, Dong K, Shang L, Jing C, Li L (2022) A novel 3'tRNA-derived fragment tRF-Val promotes proliferation and inhibits apoptosis by targeting EEF1A1 in gastric cancer. *Cell Death Dis* 13(5):471. <https://doi.org/10.1038/s41419-022-04930-6>
- Dalmay T (2013) Mechanism of miRNA-mediated repression of mRNA translation. *Essays Biochem* 54:29–38. <https://doi.org/10.1042/bse0540029>
- Dennis G, Sherman BT, Hosack DA, Yang J, Gao W, Lane HC, Lempicki RA (2003) DAVID: Database for Annotation, Visualization, and Integrated Discovery. *Genome Biol* 4(9):R60. <https://doi.org/10.1186/gb-2003-4-9-r60>
- El Rahman TA, Oirdi M, El, Gonzalez-Lamothe R, Bouarab K (2012) Necrotrophic Pathogens Use the Salicylic Acid Signaling Pathway to Promote Disease Development in Tomato. *Mol Plant-Microbe Interact* 25(12):1584–1593. <https://doi.org/10.1094/MPMI-07-12-0187-R>
- Fan J, Quan W, Li G-B, Hu X-H, Wang Q, Wang H, Li X-P, Luo X, Feng Q, Hu Z-J, Feng H, Pu M, Zhao J-Q, Huang Y-Y, Li Y, Zhang Y, Wang W-M (2020) circRNAs are involved in the rice-*Magnaporthe oryzae* Interaction. *Plant Physiol* 182(1):272–286. <https://doi.org/10.1104/pp.19.00716>
- Fasani E, DalCorso G, Zorzi G, Vitulo N, Furini A (2021) Comparative analysis identifies micro-RNA associated with nutrient homeostasis, development and stress response in *Arabidopsis thaliana* upon high Zn and metal hyperaccumulator *Arabidopsis halleri*. *Physiol Plant* 173(3):920–934. <https://doi.org/10.1111/ppl.13488>
- Fu Y, Lee I, Lee YS, Bao X (2015) Small Non-coding Transfer RNA-Derived RNA Fragments (tRFs): Their Biogenesis, Function and Implication in Human Diseases. *Genomics & Informatics* 13(4):94. <https://doi.org/10.5808/GI.2015.13.4.94>
- Galeng-Lawilao J, Kumar A, De Waele D (2018) QTL mapping for resistance to and tolerance for the rice root-knot nematode. *Meloidogyne Graminicola BMC Genetics* 19(1):53. <https://doi.org/10.1186/s12863-018-0656-1>
- Garay-Baquero DJ, White CH, Walker NF, Tebruegge M, Schiff HF, Ugarte-Gil C, Morris-Jones S, Marshall BG, Manousopoulou A, Adamson J, Vallejo AF, Bielecka MK, Wilkinson RJ, Tezera LB, Woelk CH, Garbis SD, Elkington P (2020) Comprehensive plasma proteomic profiling reveals biomarkers for active tuberculosis. *JCI Insight* 5(18). <https://doi.org/10.1172/jci.insight.137427>
- George S, Rafi M, Aldarmaki M, ElSiddig M, Al Nuaimi M, Amiri KMA (2022) tRNA derived small RNAs—Small players with big roles. *Front Genet* 13:997780. <https://doi.org/10.3389/fgene.2022.997780>
- Ghoshan MH, EL-Argawy E, Tokgöz S, Lakshman DK, Mitra A (2020) Plant Defense against Necrotrophic Pathogens. *Am J Plant Sci* 11(12):2122–2138. <https://doi.org/10.4236/ajps.2020.1112149>
- Gong M, Deng Y, Xiang Y, Ye D (2023) The role and mechanism of action of tRNA-derived fragments in the diagnosis and treatment of malignant tumors. *Cell Commun Signal* 21(1):62. <https://doi.org/10.1186/s12964-023-01079-3>
- Gu H, Lian B, Yuan Y, Kong C, Li Y, Liu C, Qi Y (2022) A 5' tRNA-Ala-derived small RNA regulates anti-fungal defense in plants. *Sci China Life Sci* 65(1):1–15. <https://doi.org/10.1007/s11427-021-2017-1>
- Guo J, He X-J (2024) Composition and function of plant chromatin remodeling complexes. *Curr Opin Plant Biol* 81:102613. <https://doi.org/10.1016/j.pbi.2024.102613>
- Gupta N, Singh A, Zahra S, Kumar S (2018) tRFdb: a database for plant transfer RNA-derived fragments. *Database* 2018. <https://doi.org/10.1093/database/bay063>
- Haidar AM, Al-Assas KM, Dawabah AAM (2016) Prevalence, distribution and intraspecific variation of *Heterodera schachtii* populations from semiarid environment. *Saudi Journal of Biological Sciences* 23(2):293–299. <https://doi.org/10.1016/j.sjbs.2015.06.017>
- Hillwig MS, LeBrasseur ND, Green PJ, MacIntosh GC (2008) Impact of transcriptional, ABA-dependent, and ABA-independent pathways on wounding regulation of RNS1 expression. *Mol Genet Genomics* 280(3):249. <https://doi.org/10.1007/s00438-008-0360-3>
- Hirsch CD, Springer NM (2017) Transposable element influences on gene expression in plants. *Biochimica et Biophysica Acta (BBA) - Gene Regulatory Mechanisms* 1860(1):157–165. <https://doi.org/10.1016/j.bbagr.2016.05.010>
- Höftberger M, Althammer M, Foissner I, Tenhaken R (2022) Galactose induces formation of cell wall stubs and cell death in *Arabidopsis* roots. *Planta* 256(2):26. <https://doi.org/10.1007/s00425-022-03919-x>
- Jones-Rhoades MW, Bartel DP, Bartel B (2006) MicroRNAs and their regulatory roles in plants. *Annu Rev Plant Biol* 57(1):19–53. <https://doi.org/10.1146/annurev.arplant.57.032905.105218>
- Kanehisa M (2000) KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Res* 28(1):27–30. <https://doi.org/10.1093/nar/28.1.27>
- Kim D, Paggi JM, Park C, Bennett C, Salzberg SL (2019) Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* 37(8):907–915. <https://doi.org/10.1038/s41587-019-0201-4>
- Kim HK, Fuchs G, Wang S, Wei W, Zhang Y, Park H, Roy-Chaudhuri B, Li P, Xu J, Chu K, Zhang F, Chua M-S, So S, Zhang QC, Sarnow P, Kay MA (2017) A transfer-RNA-derived small RNA regulates ribosome biogenesis. *Nature* 552(7683):57–62. <https://doi.org/10.1038/nature25005>
- Kim HK, J.-H. Y. M. A. K. (2020) Transfer RNA-derived small RNAs: another layer of gene regulation and novel targets for disease therapeutics. *Mol Ther* 28(11):2340–2357. <https://doi.org/10.1016/j.ymthe.2020.09.013>
- Kim K, Kim Y, Kim SH, Moon JH, Kim EH, Lee EJ, Oh C-M, Ku CR (2024a) Differential gene expression and pathway analysis in growth hormone-secreting pituitary tumors according to granulation pattern. *Frontiers in Oncology* 14:1423606. <https://doi.org/10.3389/fonc.2024.1423606>
- Kim N, Lee J, Yeom S-I, Kang N-J, Kang W-H (2024b) The landscape of abiotic and biotic stress-responsive splice variants with deep RNA-seq datasets in hot pepper. *Scientific Data* 11(1):381. <https://doi.org/10.1038/s41597-024-03239-7>
- Krishna S, Raghavan S, DasGupta R, Palakodeti D (2021) tRNA-derived fragments (tRFs): establishing their turf in post-transcriptional gene regulation. *Cell Mol Life Sci* 78(6):2607–2619. <https://doi.org/10.1007/s00018-020-03720-7>
- Kumar P, Kusc C, Dutta A (2016) Biogenesis and Function of Transfer RNA-Related Fragments (tRFs). *Trends Biochem Sci* 41(8):679–689. <https://doi.org/10.1016/j.tibs.2016.05.004>

- Kuscu C, Kumar P, Kiran M, Su Z, Malik A, Dutta A (2018) tRNA fragments (tRFs) guide Ago to regulate gene expression post-transcriptionally in a Dicer-independent manner. *RNA* 24(8):1093–1105. <https://doi.org/10.1261/rna.066126.118>
- Lalande S, Merret R, Salinas-Giegé T, Drouard L (2020) Arabidopsis tRNA-derived fragments as potential modulators of translation. *RNA Biol* 17(8):1137–1148. <https://doi.org/10.1080/15476286.2020.1722514>
- Lee YS, Shibata Y, Malhotra A, Dutta A (2009) A novel class of small RNAs: tRNA-derived RNA fragments (tRFs). *Genes Dev* 23(22):2639–2649. <https://doi.org/10.1101/gad.1837609>
- Li N, Han X, Feng D, Yuan D, Huang L-J (2019) Signaling Crosstalk between Salicylic Acid and Ethylene/Jasmonate in Plant Defense: Do We Understand What They Are Whispering? *International Journal of Molecular Sciences* 20(3):671. <https://doi.org/10.3390/ijms20030671>
- Li N, Shan N, Lu L, Wang Z (2021) tRFtarget: a database for transfer RNA-derived fragment targets. *Nucleic Acids Res* 49(D1):D254–D260. <https://doi.org/10.1093/nar/gkaa831>
- Li S, Xu Z, Sheng J (2018) tRNA-Derived Small RNA: A Novel Regulatory Small Non-Coding RNA. *Genes* 9(5):246. <https://doi.org/10.3390/genes9050246>
- Li Y, Gao J, Wang Y, Cai J, Wu D, Wang L, Pu W, Yu F, Zhu S (2023) The functions of a 5' tRNA-Ala-derived fragment in gene expression. *Plant Physiol* 193(2):1126–1141. <https://doi.org/10.1093/plphys/kiad361>
- Liu J, Liu X, Zhang S, Liang S, Luan W, Ma X (2021b) TarDB: an online database for plant miRNA targets and miRNA-triggered phased siRNAs. *BMC Genomics* 22(1):348. <https://doi.org/10.1186/s12864-021-07680-5>
- Liu L, Wu X, Sun W, Yu X, Demura T, Li D, Zhuge Q (2021a) Galactinol synthase confers salt-stress tolerance by regulating the synthesis of galactinol and raffinose family oligosaccharides in poplar. *Ind Crops Prod* 165:113432. <https://doi.org/10.1016/j.indcrop.2021.113432>
- Liu S, Chen Y, Ren Y et al (2018) A tRNA-derived RNA fragment plays an important role in the mechanism of arsenite -induced cellular responses. *Sci Rep* 8:16838
- Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* 15(12):550. <https://doi.org/10.1186/s13059-014-0550-8>
- Ma X, Liu C, Cao X (2021) Plant transfer RNA-derived fragments: Biogenesis and functions. *J Integr Plant Biol* 63(8):1399–1409. <https://doi.org/10.1111/jipb.13143>
- Macioszek VK, Jęcz T, Ciereszko I, Kononowicz AK (2023) Jasmonic Acid as a Mediator in Plant Response to Necrotrophic Fungi. *Cells* 12(7):1027. <https://doi.org/10.3390/cells12071027>
- Mao C, Yuan W, Fang R, Wu Y, Zhang Z, Cong H (2024) Transfer RNA-derived small RNAs: A class of potential biomarkers in multiple cancers (Review). *Oncol Lett* 28(1):293. <https://doi.org/10.3892/ol.2024.14427>
- Martinez G (2018) tRNA-derived small RNAs: New players in genome protection against retrotransposons. *RNA Biol* 15(2):170–175. <https://doi.org/10.1080/15476286.2017.1403000>
- Martinez G, Choudury SG, Slotkin RK (2017) tRNA-derived small RNAs target transposable element transcripts. *Nucleic Acids Res* 45(9):5142–5152. <https://doi.org/10.1093/nar/gkx103>
- Maruri-López I, Figueroa NE, Hernández-Sánchez IE, Chodasiewicz M (2021) Plant Stress Granules: Trends and Beyond. *Frontiers in Plant Science* 12:722643. <https://doi.org/10.3389/fpls.2021.722643>
- Mishra S, Roychowdhury R, Ray S, Hada A, Kumar A, Sarker U, Aftab T, Das R (2024) Salicylic acid (SA)-mediated plant immunity against biotic stresses: An insight on molecular components and signaling mechanism. *Plant Stress* 11:100427. <https://doi.org/10.1016/j.stress.2024.100427>
- Moraga C, Sanchez E, Ferrarini MG, Gutierrez RA, Vidal EA, Sagot M-F (2022) BrumiR: A toolkit for de novo discovery of micro-RNAs from sRNA-seq data. *GigaScience* 11:giac093. <https://doi.org/10.1093/gigascience/giac093>
- Mukherjee S, Sudandiradoss C (2021) Transcriptomic analysis of castration, chemo-resistant and metastatic prostate cancer elucidates complex genetic crosstalk leading to disease progression. *Funct Integr Genomics* 21(3–4):451–472. <https://doi.org/10.1007/s10142-021-00789-6>
- Mur LAJ, Kenton P, Atzorn R, Miersch O, Wasternack C (2006) The Outcomes of Concentration-Specific Interactions between Salicylate and Jasmonate Signaling Include Synergy, Antagonism, and Oxidative Stress Leading to Cell Death. *Plant Physiol* 140(1):249–262. <https://doi.org/10.1104/pp.105.072348>
- Nemeth K, Bayraktar R, Ferracin M, Calin GA (2023) Non-coding RNAs in disease: from mechanisms to therapeutics. *Nat Rev Genet*. <https://doi.org/10.1038/s41576-023-00662-1>
- Obayashi T, Hibara H, Kagaya Y, Aoki Y, Kinoshita K (2022) ATTED-II v11: A Plant Gene Coexpression Database Using a Sample Balancing Technique by Subagging of Principal Components. *Plant Cell Physiol* 63(6):869–881. <https://doi.org/10.1093/pcp/pcac041>
- Pandey D, Rajendran SRCK, Gaur M, Sajeesh PK, Kumar A (2016) Plant Defense Signaling and Responses Against Necrotrophic Fungal Pathogens. *J Plant Growth Regul* 35(4):1159–1174. <https://doi.org/10.1007/s00344-016-9600-7>
- Panthapulakkal Narayanan S, Lung S-C, Liao P, Lo C, Chye M-L (2020) The overexpression of OsACBP5 protects transgenic rice against necrotrophic, hemibiotrophic and biotrophic pathogens. *Sci Rep* 10(1):14918. <https://doi.org/10.1038/s41598-020-71851-9>
- Park E, Kim T-H (2018) Fine-Tuning of Gene Expression by tRNA-Derived Fragments during Abiotic Stress Signal Transduction. *Int J Mol Sci* 19(2):518. <https://doi.org/10.3390/ijms19020518>
- Pedro DLF, Amorim TS, Varani A, Guyot R, Domingues DS, Paschoal AR (2021) An Atlas of Plant Transposable Elements. *F1000Research* 10:1194. <https://doi.org/10.12688/f1000research.74524.1>
- Pertea R, Pertea GM, Antonescu CM, Chang T-C, Mendell JT, Salzberg SL (2015) StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. *Nat Biotechnol* 33(3):290–295. <https://doi.org/10.1038/nbt.3122>
- Petrash S, Knapp SJ, van Kan JAL, Blanco-Ulate B (2019) Grey mould of strawberry, a devastating disease caused by the ubiquitous necrotrophic fungal pathogen *Botrytis cinerea*. *Mol Plant Pathol* 20(6):877–892. <https://doi.org/10.1111/mpp.12794>
- Pietzenk B, Markus C, Gaubert H, Bagwan N, Merotto A, Bucher E, Pecinka A (2016) Recurrent evolution of heat-responsiveness in Brassicaceae COPIA elements. *Genome Biol* 17(1):209. <https://doi.org/10.1186/s13059-016-1072-3>
- Pirone R, Frugis G, Locatelli F, Mattana M, Genga A, Baldoni E (2023) Transcriptomic analysis reveals the gene regulatory networks involved in leaf and root response to osmotic stress in tomato. *Frontiers in Plant Science* 14:1155797. <https://doi.org/10.3389/fpls.2023.1155797>
- Qiu X-M, Sun Y-Y, Ye X-Y, Li Z-G (2020) Signaling Role of Glutamate in Plants. *Frontiers in Plant Science* 10:1743. <https://doi.org/10.3389/fpls.2019.01743>
- Raudvere U, Kolberg L, Kuzmin I, Arak T, Adler P, Peterson H, Vilo J (2019) g:Profiler: a web server for functional enrichment analysis and conversions of gene lists (2019 update). *Nucleic Acids Res* 47(W1):W191–W198. <https://doi.org/10.1093/nar/gkz369>
- Reuper H, Amari K, Krenz B (2021) Analyzing the G3BP-like gene family of Arabidopsis thaliana in early turnip mosaic virus infection. *Sci Rep* 11(1):2187. <https://doi.org/10.1038/s41598-021-81276-7>

- Rösti J, Barton CJ, Albrecht S, Dupree P, Pauly M, Findlay K, Roberts K, Seifert GJ (2007) UDP-Glucose 4-Epimerase Isoforms UGE2 and UGE4 Cooperate in Providing UDP-Galactose for Cell Wall Biosynthesis and Growth of *Arabidopsis thaliana*. *Plant Cell* 19(5):1565–1579. <https://doi.org/10.1105/tpc.106.049619>
- Saif R, Mahmood T, Ejaz A, Zia S (2022) Pathway enrichment and network analysis of differentially expressed genes in pashmina goat. *Gene Reports* 27:101606. <https://doi.org/10.1016/j.genrep.2022.101606>
- Sanchez L, Courteaux B, Hubert J, Kauffmann S, Renault J-H, Clément C, Baillieul F, Dorey S (2012) Rhamnolipids elicit defense responses and induce disease resistance against biotrophic, hemibiotrophic, and necrotrophic pathogens that require different signaling pathways in *Arabidopsis* and highlight a central role for salicylic acid. *Plant Physiol* 160(3):1630–1641. <https://doi.org/10.1104/pp.112.201913>
- Sato Y, Namiki N, Takehisa H, Kamatsuki K, Minami H, Ikawa H, Ohyanagi H, Sugimoto K, Itoh J-I, Antonio BA, Nagamura Y (2013) RiceFRIEND: a platform for retrieving coexpressed gene networks in rice. *Nucleic Acids Res* 41(D1):D1214–D1221. <https://doi.org/10.1093/nar/gks1122>
- Shan S, Wang Y, Zhu C (2021) A comprehensive expression profile of tRNA-derived fragments in papillary thyroid cancer. *J Clin Lab Anal* 35(3). <https://doi.org/10.1002/jcla.23664>
- Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B, Ideker T (2003) Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res* 13(11):2498–2504. <https://doi.org/10.1101/gr.1239303>
- Sherry S et al (2023) NCBI SRA toolkit technology for next generation sequence data. <http://www.ncbi.nlm.nih.gov/Traces/sra/sra.cgi?view=software>. Accessed 11 Aug 2023
- Slotkin RK, Vaughn M, Borges F, Tanurdzić M, Becker JD, Feijó JA, Martienssen RA (2009) Epigenetic reprogramming and small RNA silencing of transposable elements in pollen. *Cell* 136(3):461–472. <https://doi.org/10.1016/j.cell.2008.12.038>
- Sun X, Yang J, Yu M, Yao D, Zhou L, Li X, Qiu Q, Lin W, Lu B, Chen E, Wang P, Chen W, Tao S, Xu H, Williams A, Liu Y, Pan X, Cowley AW, Lu W, ... Lu Y (2020) Global identification and characterization of tRNA-derived RNA fragment landscapes across human cancers. *NAR Cancer* 2(4). <https://doi.org/10.1093/narcan/zcaa031>
- Tan L, Wu X, Tang Z, Chen H, Cao W, Wen C, Zou G, Zou H (2024) The tsRNAs (tRFdb-3013a/b) serve as novel biomarkers for colon adenocarcinomas. *Aging*. <https://doi.org/10.18632/aging.205590>
- Tang D, Chen M, Huang X, Zhang G, Zeng L, Zhang G, Wu S, Wang Y (2023) SRplot: A free online platform for data visualization and graphing. *PLoS ONE* 18(11):e0294236. <https://doi.org/10.1371/journal.pone.0294236>
- Thompson DM, Lu C, Green PJ, Parker R (2008) tRNA cleavage is a conserved response to oxidative stress in eukaryotes. *RNA* 14(10):2095–2103. <https://doi.org/10.1261/rna.1232808>
- Wang C, Chen W, Aili M, Zhu L, Chen Y (2023) tRNA-derived small RNAs in plant response to biotic and abiotic stresses. *Frontiers in Plant Science* 14:1131977. <https://doi.org/10.3389/fpls.2023.1131977>
- Wang H, Huang H, Ni X, Zeng W (2019) Revealing spatial-temporal characteristics and patterns of urban travel: a large-scale analysis and visualization study with taxi GPS data. *ISPRS Int J Geo Inf* 8(6):257. <https://doi.org/10.3390/ijgi8060257>
- Wang H, Lu T, Yan W, Yu P, Fu W, Li J, Su X, Chen T, Fu G, Wu Z, Feng B (2024) transcriptome and metabolome analyses reveal ascorbic acid ameliorates cold tolerance in rice seedling plants. *Agronomy* 14(4):659. <https://doi.org/10.3390/agronomy14040659>
- Wang J, Liu X, Cui W, Xie Q, Peng W, Zhang H, Gao Y, Zhang C, Duan C (2022) Plasma tRNA-derived small RNAs signature as a predictive and prognostic biomarker in lung adenocarcinoma. *Cancer Cell Int* 22(1):59. <https://doi.org/10.1186/s12935-022-02481-6>
- Xie K, Zhang T, Zhou X, Chen G (2015) Genome-wide analyses of abiotic stress-related microRNAs and their targets in *Arabidopsis thaliana*. Southern Cross Publishing Group. *Plant Omics J* 8(3):238–243. https://www.pomics.com/zhou_8_3_2015_238_243.pdf
- Yagyu M, Yoshimoto K (2024) New insights into plant autophagy: molecular mechanisms and roles in development and stress responses. *J Exp Bot* 75(5):1234–1251. <https://doi.org/10.1093/jxb/erad459>
- Yang L, Li B, Zheng X, Li J, Yang M, Dong X, He G, An C, Deng XW (2015) Salicylic acid biosynthesis is enhanced and contributes to increased biotrophic pathogen resistance in *Arabidopsis* hybrids. *Nat Commun* 6(1):7309. <https://doi.org/10.1038/ncomms8309>
- Yates AD, Allen J, Amode RM, Azov AG, Barba M, Becerra A, Bhai J, Campbell LI, Carbajo Martinez M, Chakiachvili M, Chougule K, Christensen M, Contreras-Moreira B, Cuzick A, Da Rin Fioretto L, Davis P, De Silva NH, Diamantakis S, Dyer S, ... Flicek P (2022) Ensembl Genomes 2022: an expanding genome resource for non-vertebrates. *Nucleic Acids Res* 50(D1):D996–D1003. <https://doi.org/10.1093/nar/gkab1007>
- Yu X, Xie Y, Zhang S, Song X, Xiao B, Yan Z (2021) tRNA-derived fragments: Mechanisms underlying their regulation of gene expression and potential applications as therapeutic targets in cancers and virus infections. *Theranostics* 11(1):461–469. <https://doi.org/10.7150/thno.51963>
- Zahra S, Bhardwaj R, Sharma S, Singh A, Kumar S (2022) PtnCRNAdb: plant transfer RNA-derived non-coding RNAs (tncRNAs) database. 3. *Biotech* 12(5):105. <https://doi.org/10.1007/s13205-022-03174-7>
- Zahra S, Singh A, Kumar S (2023) tncRNA Toolkit: A pipeline for convenient identification of RNA (tRNA)-derived non-coding RNAs. *MethodsX* 10:101991. <https://doi.org/10.1016/j.mex.2022.101991>
- Zahra S, Singh A, Poddar N, Kumar S (2021) Transfer RNA-derived non-coding RNAs (tncRNAs): Hidden regulation of plants' transcriptional regulatory circuits. *Comput Struct Biotechnol J* 19:5278–5291. <https://doi.org/10.1016/j.csbj.2021.09.021>
- Zhang Y, Gu X, Li Y, Huang Y, Ju S (2024) Multiple regulatory roles of the transfer RNA-derived small RNAs in cancers. *Genes & Diseases* 11(2):597–613. <https://doi.org/10.1016/j.gendis.2023.02.053>
- Zhang Z, Yu J, Li D, Zhang Z, Liu F, Zhou X, Wang T, Ling Y, Su Z (2010) PMRD: plant microRNA database. *Nucleic Acids Research* 38(suppl_1):D806–D813. <https://doi.org/10.1093/nar/gkp818>
- Zhao R, Yang Z, Zhao B, Li W, Liu Y, Chen X, Cao J, Zhang J, Guo Y, Xu L, Wang J, Sun Y, Liu M, Tian L (2023) A novel tyrosine tRNA-derived fragment, tRFTyr, induces oncogenesis and lactate accumulation in LSCC by interacting with LDHA. *Cell Mol Biol Lett* 28(1):49. <https://doi.org/10.1186/s11658-023-00463-8>
- Zhu L, Ge J, Li T, Shen Y, Guo J (2019a) tRNA-derived fragments and tRNA halves: The new players in cancers. *Cancer Lett* 452:31–37. <https://doi.org/10.1016/j.canlet.2019.03.012>
- Zhu L, Li T, Shen Y, Yu X, Xiao B, Guo J (2019b) Using tRNA halves as novel biomarkers for the diagnosis of gastric cancer. *Cancer Biomark* 25(2):169–176. <https://doi.org/10.3233/CBM-182184>

Zong T, Yang Y, Zhao H, Li L, Liu M, Fu X, Tang G, Zhou H, Aung LHH, Li P, Wang J, Wang Z, Yu T (2021) tsRNAs: Novel small molecules from cell function and regulatory mechanism to therapeutic targets. *Cell Proliferation* 54(3):e12977. <https://doi.org/10.1111/cpr.12977>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.