

Discharge Instructions

CHOC Children's Hospital

1201 West La Veta Avenue, Orange, CA 92868

Patient: Tina Johnson

MRN#: 664372

Preferred Language: English

Weight(kg): 9.4 kg

Allergies: NKA

Date of Admission: 04/01/2026

Date of Discharge: 04/03/2026

Admitting Diagnosis: Seizure Disorder

Discharge Diagnosis: Seizure Disorder

Condition on Discharge: Good

Primary Care Physician: Dr. Burch

Notable or Outside Test/Results:

Diet: Regular, Age Appropriate

Seek Medical Attention for: Persistent Temp > 100.4 F, New or Worsened Seizures, Persistent Vomiting, Breathing Difficulties

Special Instructions:

Please follow up with pediatrician as needed.

Follow up with subspecialists as scheduled. Neuro will reschedule the next appointment.

Additional RN Instructions:

Instruction Forms Given:

MEDICATION INSTRUCTIONS (Follow pharmacy instructions):

Continue the Following Home Medications:

Diastat AcuDial 10 mg rectal kit

7.5 mg Per rectum Once (when indicated) PRN for seizure greater than 5 minutes or more than 3 seizures in 1 hour, 2 refills, CVS Pharmacy #8821

Polyethylene glycol 3350 powder

As directed Oral Once daily PRN for constipation, CVS Pharmacy #8821

New Prescriptions:

No New Prescription

Pre-Existing Appointments: To contact scheduling call 1-888-770-CHOC (2462)

No scheduled appointments at this time.

Immunizations:

Procedures:

Electroencephalogram (EEG), continuous recording, physician or other qualified health care professional review of recorded events, analysis of spike and seizure detection, interpretation and report, 2-12 hours of EEG recording; with video (VEEG)

Consulting Physicians: Dr. Campos, MD

Discharge Authenticated by: Dr. Tran, MD

AUTHENTICATED BY ATTENDING

How to Give Diazepam Rectal Gel to Your Child

At today's visit, the health care provider talked with you about how and when to give your child diazepam rectal gel. This medicine helps to stop some seizures. It is a liquid medicine given in the rectum (where poop comes out). Because you may need to use the gel in an emergency, it's important to know what to do so you're ready if your child needs it.

When you pick up diazepam rectal gel at the pharmacy:

- Tell the pharmacist about all other medicines, herbs, or supplements your child takes.

Open the box and:

- Check the expiration date on both syringes.
- Check that the right dose appears in the window on the side of each syringe and you can see the green "READY" band.
- Make sure the instructions with pictures on how to use the medicine are included.

At home:

- Read the instructions on how to use the medicine so you're ready if you need to give it to your child. Keep the instructions with the syringes so you can look at them when it's time to use the medicine.

Keep the medicine where you can get it quickly if you need it:

- Choose a dry place that doesn't get too hot or too cold (never keep diazepam rectal gel in a bathroom, a car, etc.).
- Put the medicine where young children can't reach it, like in a childproof or high cabinet.
- Keep a notebook or paper and pen with the syringes so you can write down seizure timing and when the medicine was given.
- Follow the health care provider's instructions for when and how often to give your child diazepam rectal gel.
- Follow the instructions included in the box for how to give diazepam rectal gel.
- Keep giving your child their normal seizure medicines. Diazepam rectal gel is not a replacement for these medicines.
- If your child is in school, talk to the school nurse about keeping a dose there with instructions on how and when to use it.

After giving diazepam rectal gel:

- Keep your child on their side facing you.
- Follow the instructions for throwing away the syringe safely.
- Wash your hands with soap and water.
- Stay with your child for 4 hours to make sure that the seizures stop and to watch for any changes in breathing or skin color or side effects.
- Let your child rest. Diazepam can make kids sleepy.

Call your health care provider if:

- You have any questions about when or how to give diazepam rectal gel.
- Your child needs diazepam rectal gel more often than directed.

Call 911 if:

- Your child is still having seizures 15 minutes after you gave the diazepam rectal gel, unless your health care provider gave you other directions.
- Your child has trouble breathing or turns blue.
- Your child's seizures seem different or more serious than usual.
- Your child was injured during the seizure.

More to Know

Should I keep giving my child their daily seizure medicines? Yes. Diazepam rectal gel does not replace your child's other medicines.

Why does diazepam gel need to go in the rectum? It is not safe to give medicine in the mouth during a seizure because your child could choke.

Does the diazepam rectal gel syringe have a needle? No. The tip that goes into the rectum is plastic and not sharp.

Does diazepam rectal gel have side effects? Kids usually feel sleepy after getting diazepam. Sometimes kids get a rash, dizziness, headache, belly pain, nervousness, or wheezing. Let your health care provider know if you notice any of these side effects.

When should I give my child diazepam rectal gel? Diazepam rectal gel is used only for kids who have long seizures or lots of seizures. Give the medicine to your child exactly as your health care provider told you.

How To Monitor, Secure, And Destroy Your Medications

Preventing prescription drug abuse starts in your home. According to the U.S. Drug Enforcement Administration, more people die from over-the-counter and prescription drug abuse than from cocaine, heroin, and ecstasy combined. Safeguard your family by following these steps.

MONITOR

Take inventory. Count your medications from the pharmacist. Count them as you use them. Take note of missing and quickly used medications.

SECURE

Lock Medications and reduce access. Don't leave them in a bathroom. Keep them away from children and youth.

DESTROY

Never dispose of medications in a sink or toilet. Dissolve unused or expired medications in a baggie with kitty litter.

How To Properly Destroy Your Expired And Unwanted Medications At Home

What You Will Need:

- Expired and unwanted medications
- Zip baggie
- Hot water (over 110°F)
- Kitty litter

Steps To Follow:

- 1) Pour unwanted/expired medications out of their original containers into a zip baggie.
- 2) Pour hot water (over 110°F - about as hot as a cup of coffee) into the baggie.
- 3) Insert kitty litter into the baggie. Seal baggie. Place in trash bin.

Shred prescription labels or use a black marker to cross out label information.

For more information, visit <http://www.choc.org/programs-services/pharmacy/>

For those living outside of Orange County, please call your county public health agency or search on the web for "[Name of County] medication or drug disposal". i.e. "Riverside county medication or drug disposal"

Adapted from Orange County RX Abuse Prevention Coalition, County of Orange Health Care Agency

How to Address Your Child's Discomfort and/or Pain While at Home

Your healthcare provider will tell you if your child will need any medication for discomfort and/or pain while at home.

There are many other ways you can help keep your child comfortable without using medications or with using less medications.

If okay with your healthcare provider, this can include:

- Use of cool and/or warm packs
- Help your child change position in bed or chair
- Encourage your child to be up in a chair or walking
- Massage your child's feet, hands, and back with lotion
- Distraction; such as art, crafts, music, TV/videos

CHOC Link and COVID-19 Resources

CHOC Link is an online tool to manage your child's medical records and communicate with your providers and clinics. If interested in getting access, please provide a valid email address to your CHOC provider office or to the Admissions Department, and we'll send you an invitation to create your account. To learn more, go to www.choc.org/choclink.

Please visit 'Coronavirus (COVID-19): A Resource for Parents' at choc.org/coronavirus for frequently asked questions and up to date information from Centers for Disease Control and Prevention, Orange County Health Care Agency, California Department of Health, and CHOC Children's including visitor information.

If you have questions about your child's health and COVID-19, call our 24/7 Nurse line at **1-844-GET-CHOC**.