

Discharge Instructions

CHOC Children's Hospital

1201 West La Veta Avenue, Orange, CA 92868

Patient: Crystal King

MRN#: 533414

Preferred Language: English

Weight(kg): 19.0 kg

Allergies: Penicillin

Date of Admission: 03/20/2026

Date of Discharge: 03/27/2026

Admitting Diagnosis: Allergic Reaction

Discharge Diagnosis: Allergic Reaction

Condition on Discharge: Good

Primary Care Physician: Dr. Garza

Notable or Outside Test/Results:

Diet: Regular, Age Appropriate

Seek Medical Attention for: Difficulty breathing, Swelling of face/lips/tongue, Hives spreading, Dizziness, Repeated vomiting

Special Instructions:

Please follow up with pediatrician as needed.

Avoid known allergens. Carry EpiPen at all times. Follow up with allergy specialist as scheduled.

Additional RN Instructions:

Instruction Forms Given:

MEDICATION INSTRUCTIONS (Follow pharmacy instructions):

Continue the Following Home Medications:

EpiPen Jr 0.15 mg auto-injector

0.15 mg Intramuscular PRN for severe allergic reaction, 1 refill, Walgreens #5821

Diphenhydramine 12.5 mg/5 mL elixir

As directed by weight Oral Every 6 hours PRN for itching/hives, 0 refills, Walgreens #5821

New Prescriptions:

Azithromycin 200 mg/5 mL suspension

As directed Oral Once daily for 5 days, 0 refills, Walgreens #5821

Pre-Existing Appointments: To contact scheduling call 1-888-770-CHOC (2462)

No scheduled appointments at this time.

Immunizations:

Procedures:

Allergy testing panel

Complete blood count

Consulting Physicians: Dr. Pierce, MD

Discharge Authenticated by: Dr. Bennett, MD

AUTHENTICATED BY ATTENDING

Managing Your Child's Severe Allergies

At today's visit, the health care provider talked with you about managing your child's severe allergies. Here's what you need to know to keep your child safe.

Avoiding Allergens:

- Read all food labels carefully.
- Ask about ingredients when eating out.
- Inform school, caregivers, and family members about your child's allergies.
- Your child should wear a medical alert bracelet.

Recognizing Anaphylaxis (Severe Allergic Reaction):

- Hives, itching, flushed skin
- Swelling of lips, tongue, or throat
- Difficulty breathing, wheezing
- Dizziness, fainting
- Stomach pain, vomiting, diarrhea
- Feeling of doom

Using the EpiPen:

1. Remove the blue safety cap.
2. Hold the orange tip against the outer thigh (through clothing is OK).
3. Press firmly until you hear a click.
4. Hold in place for 10 seconds.
5. Remove and massage the area.
6. Call 911 immediately - your child needs to go to the hospital even if symptoms improve.

Important Tips:

- Always carry two EpiPens with your child.
- Check expiration dates regularly.
- Keep EpiPens at room temperature (not in hot cars or refrigerators).
- Make sure school has an EpiPen and action plan on file.

Call 911 immediately if:

- Your child shows any signs of anaphylaxis.
- You have given the EpiPen (always go to the ER after using an EpiPen).

How To Monitor, Secure, And Destroy Your Medications

Preventing prescription drug abuse starts in your home. According to the U.S. Drug Enforcement Administration, more people die from over-the-counter and prescription drug abuse than from cocaine, heroin, and ecstasy combined. Safeguard your family by following these steps.

MONITOR

Take inventory. Count your medications from the pharmacist. Count them as you use them. Take note of missing and quickly used medications.

SECURE

Lock Medications and reduce access. Don't leave them in a bathroom. Keep them away from children and youth.

DESTROY

Never dispose of medications in a sink or toilet. Dissolve unused or expired medications in a baggie with kitty litter.

How To Properly Destroy Your Expired And Unwanted Medications At Home

What You Will Need:

- Expired and unwanted medications
- Zip baggie
- Hot water (over 110°F)
- Kitty litter

Steps To Follow:

- 1) Pour unwanted/expired medications out of their original containers into a zip baggie.
- 2) Pour hot water (over 110°F - about as hot as a cup of coffee) into the baggie.
- 3) Insert kitty litter into the baggie. Seal baggie. Place in trash bin.

Shred prescription labels or use a black marker to cross out label information.

For more information, visit <http://www.choc.org/programs-services/pharmacy/>

For those living outside of Orange County, please call your county public health agency or search on the web for "[Name of County] medication or drug disposal". i.e. "Riverside county medication or drug disposal"

Adapted from Orange County RX Abuse Prevention Coalition, County of Orange Health Care Agency

How to Address Your Child's Discomfort and/or Pain While at Home

Your healthcare provider will tell you if your child will need any medication for discomfort and/or pain while at home.

There are many other ways you can help keep your child comfortable without using medications or with using less medications.

If okay with your healthcare provider, this can include:

- Use of cool and/or warm packs
- Help your child change position in bed or chair
- Encourage your child to be up in a chair or walking
- Massage your child's feet, hands, and back with lotion
- Distraction; such as art, crafts, music, TV/videos

CHOC Link and COVID-19 Resources

CHOC Link is an online tool to manage your child's medical records and communicate with your providers and clinics. If interested in getting access, please provide a valid email address to your CHOC provider office or to the Admissions Department, and we'll send you an invitation to create your account. To learn more, go to www.choc.org/choclink.

Please visit 'Coronavirus (COVID-19): A Resource for Parents' at choc.org/coronavirus for frequently asked questions and up to date information from Centers for Disease Control and Prevention, Orange County Health Care Agency, California Department of Health, and CHOC Children's including visitor information.

If you have questions about your child's health and COVID-19, call our 24/7 Nurse line at **1-844-GET-CHOC**.