## Chapter 1 Exercises: Number Systems

Kyle Swanson

February 5, 2015

1.8 What is the largest 32-bit unsigned number?

$$2^{32} - 1 = 4,294,967,295$$

1.10 What is the largest 32-bit binary number that can be represented with

- (a) unsigned numbers?
- $2^{32} 1 = 4,294,967,295$
- (b) two's complement numbers?
- $2^{31} 1 = 2,147,483,647$
- (c) sign/magnitude numbers?
- $2^{31} 1 = 2,147,483,647$

 ${\bf 1.12}$  What is the smallest (most negative) 32-bit binary number that can be represented with

(a) unsigned numbers?

If you can't have a sign, either it's impossible to have a negative value, or you assume that all values are negative. A negative-int data type I suppose.

$$-2^{32} - 1 = -4,294,967,295$$

- (b) two's complement numbers?
- $-2^{31} = -2,147,483,648$
- (c) sign/magnitude numbers?

$$-2^{31} - 1 = -2,147,483,647$$

1.14 Convert the following unsigned binary numbers to decimal.

(a) 1110<sub>2</sub>

$$2^{3} + 2^{2} + 2^{1} + 0 = 14$$

- (b)  $100100_2$
- $2^5 + 2^2 = 36$
- (c) 11010111<sub>2</sub>

$$2^7 + 2^6 + 2^4 + 2^2 + 2^1 + 20 = 215$$

(d) 011101010100100<sub>2</sub>

$$2^{13} + 2^{12} + 2^{11} + 2^9 + 2^7 + 2^5 + 2^2 = 15,012$$

1.16 Repeat 1.14, but convert to hexadecimal.

Since they are split into 4 bit sections, just match each section with it's hex digit.

- (a)  $1110_2$
- 0xE
- (b) 10 0100<sub>2</sub>
- 0x24
- (c) 1101 0111<sub>2</sub>
- 0xD7

(d)  $011\ 1010\ 1010\ 0100_2$  0x3AA4

1.18 Convert the following hexadecimal numbers to decimal.

(a) 
$$0x4E$$

$$4 * 16^1 + 14 * 16^0 = 78$$

(b) 
$$0x7C$$

$$7*16^1 + 12*160 = 124$$

(c) 
$$0xED3A$$

$$14 * 16^3 + 13 * 16^2 + 3 * 16^1 + 10 * 16^0 = 60,730$$

$$4 * 16^7 + 3 * 16^5 + 15 * 16^4 + 11 * 16^3 + 1 * 16^0 = 1,077,915,649$$

1.20 Repeat 1.18, but convert to unsigned binary.

Simply take each digit, and match it to it's binary equivalent. Then push the resulting sections together.

(a) 0x4E

0100 1110

(b) 0x7C

0111 1100

(c) 0xED3A

1110 1101 0011 1010

(d) 0x403FB001

0100 0000 0011 1111 1011 0000 0000 0001

1.22 Convert the following two's complement binary numbers to decimal.

(a) 
$$1110_2$$

The left most bit is 1, flip the bits, 1110 = 0001 then add 1. 0001 + 0001 = 0010 = 2

Since the left most bit was 1, the result is negative. -2

(b) 100011<sub>2</sub>

$$011100 + 000001 = 011101$$

$$2^4 + 2^3 + 2^2 + 2^0 = -29$$

(c) 01001110<sub>2</sub>

The left most bit is 0, so this is a positive number. Continue as normal.

$$2^6 + 2^3 + 2^2 + 2^1 = 78$$

(d) 10110101<sub>2</sub>

01001010 + 00000001 = 01001011

$$2^6 + 2^3 + 2^1 + 2^0 = -75$$

1.26 Convert the following decimal numbers to unsigned binary numbers.

- (a) 14
- 14/2 = 7r0
- 7/2 = 3r1
- 3/2 = 1r1
- 1/2 = 0r1
- 14=1110
- (b) 52
- 52/2 = 26r0
- 26/2 = 13r0
- 13/2 = 6r1
- 6/2 = 3r0
- 3/2 = 1r1
- 1/2 = 0r1
- 52 = 110100
- (c) 339
- 339/2 = 169r1
- 169/2 = 84r1
- 84/2 = 42r0
- 42/2 = 21r0
- 21/2 = 10r1
- 10/2 = 5r0
- 5/2 = 2r1
- 2/2 = 1r0
- 1/2 = 0r1
- 339 = 101010011
- (d) 711
- 711/2 = 355r1
- 355/2 = 177r1
- 177/2 = 88r1
- 88/2 = 44r0
- 44/2 = 22r0
- 22/2 = 11r0
- 11/2 = 5r1
- 5/2 = 2r1
- 2/2 = 1r0
- 1/2 = 0r1
- 711 = 1011000111

1.28 Repeat Exercise 1.26, but convert to hexadecimal.

Since it's converted to binary already, you can start with that, then go to hex.

(a) 14

14 = 1110 = 0xE

(b) 52

 $52 = 0011\ 0100 = 0x34$ 

(c) 339

 $339 = 0001\ 0101\ 0011 = 0x153$ 

(d) 711

 $711 = 0010\ 1100\ 0111 = 0x2C7$ 

1.30 Convert the following decimal numbers to 8-bit two's complement numbers or indicate that the decimal number would overflow the range.

(a) 24

24/16 = 1r8

 $24 = 0x18 = 0001\ 1000$ 

Since it's positive, the leading bit should be 0.

(b) -59

59/16 = 3r11

 $59 = 0x3B = 0011\ 1011$ 

Invert bits and add 1

 $1100\ 0100 + 0001 = 1100\ 0101$ 

(c) 128

128/16 = 8r0

 $128 = 0x80 = 1000\,0000$ 

But, we have a problem. Since 128 on it's own takes up 8 bits, this would read as -128 in two's complement. With only 8bits, there is no way to represent +128. Overflow

(d) -150

150/16 = 9r6

 $150 = 0x96 = 1001\ 0110$ 

Invert and add 1,  $0110\ 1001 + 0001 = 0110\ 1010$ 

Again, since we only have 8 bits, this causes an overflow.

(e) 127 127/16 = 7r15 $127 = 0x7F = 0111\ 1111$ 

Luckily, this does not overflow, but it is at the end of an 8bit two's complement numbers range.

- **1.34** Convert the following 4-bit two's complement numbers to 8-bit two's complement numbers.
- (a) 0111

1111 1001

Since this is a positive number nothing needs to be changed. 0000 0111

(b) 1001 Just add 1's to the bits in front, since this is a negative number.

**1.36** Repeat Exercise 1.34 if the numbers are unsigned rather than two's complement.

I'm not completely sure I understand this question. Are we converting two's complement numbers to 8bit unsigned? Or 4bit unsigned to 8bit two's complement? Or just unsigned to unsigned?

Either way, the answer should just be the original plus 0000 at the front. This is since, if they are unsigned to start with, they are positive, and therefore the two's will be positive. If they are two's moving to unsigned, b just doesn't really make sense. Anyway, here's the answer assuming they are unsigned, moving to two's complement.

- (a) 0111 0000 0111
- (b) 1001 0000 1001
- **1.38** Base 8 is referred to as octal. Convert each of the numbers from 1.26 to octal.
- (a) 14 14/8 = 1r6 1/8 = 0r114 = 016
- (b) 52

$$52/8 = 6r4$$
  
 $6/8 = 0r6 \ 52 = 064$ 

(c) 
$$339$$
  
 $339/8 = 42r3$   
 $42/8 = 5r2$   
 $5/8 = 0r5$ 

339 = 0523

(d) 
$$711$$
  
 $711/8 = 88r7$   
 $88/8 = 11r0$   
 $11/8 = 1r3$   
 $1/8 = 0r1$   
 $711 = 01307$ 

 ${f 1.40}$  Convert each of the following octal numbers to binary, hexadecimal, and decimal.

In the following questions, converting to binary, then binary to hex is easiest.

(a) 023

binary:  $010\,011$  hex: 0x13

decimal:  $2 * 8^1 + 3 * 8^0 = 19$ 

(b) 045

binary:  $100\ 101$  hex: 0x25

decimal:  $4 * 8^1 + 5 * 8^0 = 37$ 

(c) 0371

binary: 011 111 001

hex: 0.11111001 = 0xF9

decimal:  $3 * 8^2 + 7 * 8^1 + 1 * 8^0 = 249$ 

(d) 02560

binary: 010 101 110 000

hex:  $0101\ 0111\ 0000 = 0x570$ 

decimal:  $2 * 8^3 + 5 * 8^2 + 6 * 8^1 + 0 * 8^0 = 1392$ 

1.42 How many 7-bit two's complement numbers are greater than 0? How

many numbers are less than 0

The range of a 7-bit two's complement number is  $-(2^6)$  to  $2^6-1$ , which is -64 to 63

So the number **greater** than 0, is 63, and the number **less** is 64.

1.44 How many bytes are in a 64-bit word?

8-bits to one byte. Therefore, 64/8 = 8 bytes

1.50 Draw a line number analogous to Figure 1.11 for 3-bit unsigned, two's complement, and sign/magnitude numbers.

TODO! TODO!

- 1.52 Perform the following additions of unsigned binary numbers. Indicate whether or not the sum overflows a 4-bit result.
- (a) 1011 + 0100

1111

(b) 
$$1101 + 1011$$
 $111$ 
 $1101$ 
 $+1011$ 
 $11000$ 

This does overflow the 4-bit result. The 4-bit version would simply be 1000

- **1.54** Repeat exercise 1.52, assuming that the binary numbers are in two's complement form.
- (a) 1011 + 0100
- 1011 = -5
- 0100 = 4

$$1011 + 0100 = 1111 = -1$$

- (b) 1101 + 1011
- 1101 = -3
- 1011 = -5
  - 111
  - $1\,1\,0\,1$
- +1011
- $1\ 1000$

While it does overflow, in this instance, 1000 = -8, so the answer is still correct.

1.56 Convert the following decimal numbers to 6-bit two's complement binary numbers and add them. Indicate whether or not the sum overflows a 6-bit result.

```
(a) 16 + 9
16 = 010000
9 = 001001
010000 + 001001 = 011001
(b) 27 + 31
27/2 = 13r1
13/2 = 6r1
6/2 = 3r0
3/2 = 1r1
1/2 = 0r1
27 = 011011
31/2 = 15r1
15/2 = 7r1
7/2 = 3r1
3/2 = 1r1
1/2 = 0r1
31 = 011111
   11111
   0\,1\,1\,0\,1\,1
 +0111111
   111010
111010 = -6
```

Even though there are no lost bits, since they msb is set to 1, with not enough bits for a leading 0. This is represented as a -6, which is incorrect.

```
(c) -4 + 19

-4 = 000100 = 111011 + 0001 = 111100

19/2 = 9r1

9/2 = 4r1

4/2 = 2r0

2/2 = 1r0

1/2 = 0r1

19 = 010011
```

```
\begin{array}{c}
11 \\
111100 \\
+010011 \\
1001111
\end{array}
```

This overflows properly, since -4 + 19 is a positive number, the msb should be 0.

$$001111 = 15$$

(d) 
$$3 + -32$$
  
 $3 = 000011$   
 $-32 = 100000$   
 $000011 + 100000 = 100011 = -29$ 

(e) 
$$-16 + -9$$

$$(f) -27 + -31$$

- 1.58 Perform the following additions of unsigned hexadecimal numbers. Indicate whether or not the sum overflows an 8bit (two hex digit) result.
- **1.60** Convert the following decimal numbers to 5-bit two's complement binary numbers and subtract them. Indicate whether or not the difference overflows a 5-bit result.
- **1.65** Answer the following questions related to *Binary Coded Decimal* systems.