

Bursty infectiousness as a hidden driver of epidemic variability

Stephen M. Kissler^{1,3,4*} [Full authorship and ordering TBD]

*Correspondence: stephen.kissler@colorado.edu

Abstract

The population-level infectiousness profile, $A(\tau) = R_0 g(\tau)$, determines the mean-field dynamics of an epidemic but need not resemble any individual's infectiousness trajectory. We introduce a smoothness parameter, $\psi \in [0, 1]$, that governs how much of the generation interval's variability arises within *vs.* between individuals, and show that ψ creates effects invisible to standard renewal-equation models. We demonstrate that overdispersion in secondary infection counts can arise even in populations with no variation in biological infectiousness or average contact rates—a result not captured by existing frameworks for super-spreading. This overdispersion emerges from the interaction between narrow infectiousness windows and time-varying contacts, and it cannot be reduced by identifying high-risk individuals. Using transmission-tree data from eight pathogens, we find preliminary evidence that ψ varies substantially across pathogens, from near zero (measles, pneumonic plague) to near one (COVID-19, smallpox), motivating further investigation with purpose-collected data. We show that ψ governs the sensitivity of detect-and-isolate interventions to detection timing and modulates the overdispersion-reducing effects of gathering size restrictions. These findings identify an overlooked axis of variation in epidemic dynamics.

To isolate this effect, we introduce the Gamma burst model for individual infectiousness. At the population level, we assume $g(\tau)$ is (approximately) Gamma-distributed with shape α and rate β . At the individual level, secondary infections are distributed according to a $\text{Gamma}(\psi\alpha, \beta)$ distribution — the “burst” — which follows a latent period of length $l_i \sim \text{Gamma}((1 - \psi)\alpha, \beta)$, with $\psi \in [0, 1]$ (**Figure XX**). The burstiness parameter, ψ , defines the width of the individual infectiousness kernel relative to $g(\tau)$: $\psi = 0$ is maximally bursty, with all infectiousness happening at a single moment, while $\psi = 1$ is maximally smooth, with infectiousness spread across the full generation interval distribution. Since the latent period and post-latent infection time are both Gamma-distributed, their sum follows a $\text{Gamma}(\alpha, \beta) = g(\tau)$ distribution, ensuring that any value of ψ yields the same generation interval distribution. This model of infectiousness has biological grounding in within-host viral dynamics: stochastic viral replication early in an infection causes variability in when the virus can first be detected (the variable latent period), while the kinetics thereafter are largely consistent across individuals (the infectiousness burst) [Morris et al., 2025].

EMPIRICAL ESTIMATES OF ψ . The relative lack of attention to ψ may stem partly from the difficulty in empirically estimating it. Ideally, contact tracing data would capture precise infection times and perfectly link each index case to all their secondary infections; from this, estimating ψ would be straightforward. The reality is far messier: case ascertainment is imperfect, infection times are blurry, and detection lags and multiple exposures can make it hard to disentangle who infected whom. Incorporating viral kinetics can help: Goyal et al. [2021] showed that the dose-response relationship between viral load and transmission probability restricts the window of high transmission risk to less than 1 day for both SARS-CoV-2 and influenza, consistent with low ψ .

Bearing these limitations in mind, we estimated ψ from published transmission tree data for eight diseases: COVID-19, MERS, measles, Ebola, smallpox, pneumonic plague, norovirus, and hepatitis A. We

extracted serial intervals from the OutbreakTrees database [x], grouping them by index case to form clusters. For each pathogen, we fixed the generation interval distribution to published estimates and placed a flat prior on ψ , and the likelihood was computed by integrating over all possible latent periods for each cluster (**Methods**).

The resulting estimates spanned a wide range (**Figure XX**). Measles and pneumonic plague showed the strongest evidence of burstiness (95% credible interval for ψ : [0.00, 0.11] and [0.00, 0.20], respectively), while smallpox and MERS showed evidence of more diffuse infectiousness (95% credible interval for ψ : [xx,xx] and [xx,xx], respectively). These estimates should be regarded as illustrative rather than conclusive: multiple biases could be in play. Uncertainty in infection time can inflate estimates of ψ by causing infection times to appear artificially diffuse; while case ascertainment may be biased towards superspreading events and miss less explosive, more distributed-in-time transmission, driving estimates of ψ downward. Nevertheless, the breadth of the estimated range — from near 0 to near 1 — suggests that ψ may vary meaningfully across pathogens.

A simulation-based power analysis indicates that ψ is identifiable from contact-tracing data under reasonably favorable conditions: if the distribution of uncertainty in infection times is known, 50 index cases sufficed for well-powered inference of ψ when $R_0 \geq 6$ (e.g., SARS-CoV-2 or measles). Inference was more challenging for lower R_0 , where fewer secondary cases per index case limit the amount of information available on the correlation of infection times within clusters.

BURSTY INFECTIOUSNESS DRIVES COINCIDENCE SUPERSPREADING.

BURSTINESS AND THE EPIDEMIC LATENT PERIOD.

COMMON INTERVENTIONS ARE SENSITIVE TO BURSTY INFECTIOUSNESS.

BURSTINESS COMPLICATES ESTIMATION OF KEY EPIDEMIOLOGICAL PARAMETERS.

Population-level analyses often use average quantities to describe heterogeneous systems, particularly when variation does not arise from identifiable groups[Lloyd-Smith et al., 2005]. A central example is the infectiousness profile, $A(\tau) = R_0 g(\tau)$, which captures the expected amount and timing of transmission from an infected individual and is the key parameter in the renewal equation framework for epidemic modeling[Wallinga and Lipsitch, 2007, Breda et al., 2012]. But $A(\tau)$ is an expectation over individuals: person i 's infectiousness trajectory is $a_i(\tau) = v_i g_i(\tau)$, where v_i is the individual reproduction number and $g_i(\tau)$ is the individual generation interval distribution[Champredon and Dushoff, 2015, Svensson, 2007]. Substantial work has characterized how variation in v_i —the area under $a_i(\tau)$ —drives superspreading and shapes outbreak dynamics[Lloyd-Smith et al., 2005]. But the impact of variation in $g_i(\tau)$ —how $a_i(\tau)$ is distributed over time—has not been studied systematically.

Since $A(\tau) = E_i[a_i(\tau)]$, the population-level profile need not resemble any single person's trajectory. The standard SEIR model illustrates this starkly: each person has a discontinuous, stepwise infectiousness profile (zero during latency, constant during infection), yet the population average is a smooth, bell-shaped curve (Fig. ??a). Other choices of $a_i(\tau)$ can also produce the same $A(\tau)$: setting $a_i(\tau) \equiv A(\tau)$ for all individuals gives a "smooth" profile in which each person follows the population mean; at the opposite extreme, concentrating each person's infectiousness at a single random moment gives a "spike" profile. All produce identical deterministic epidemic dynamics, since R_0 , $g(\tau)$, and $A(\tau)$ are unchanged. The stochastic dynamics, however, can differ profoundly.

To isolate this effect, we introduce the Gamma time-shifted model, parameterized by a single smoothness parameter $\psi \in [0, 1]$. Each person's secondary infections occur at times $\tau_{ij} = l_i + \epsilon_j$, where $l_i \sim \text{Gamma}((1 - \psi)\alpha, \beta)$ is a shared latent period and $\epsilon_j \stackrel{iid}{\sim} \text{Gamma}(\psi\alpha, \beta)$ is an independent jitter (Fig. ??b). Since both are Gamma-distributed with common rate β , their sum follows $\text{Gamma}(\alpha, \beta) = g(\tau)$ regardless of ψ , ensuring that all population-level quantities remain invariant. When $\psi = 1$, there is no between-individual variation: every person's infections are independently drawn from $g(\tau)$. When $\psi = 0$, there is no within-individual variation: each person's infections are concentrated at a single moment drawn from $g(\tau)$. This decomposition has biological grounding in within-host viral dynamics: stochastic early-infection processes produce between-individual variation in the timing of peak infectiousness, while subsequent kinetics are more stereotyped across individuals[Morris et al., 2025], yielding exactly the time-shifted structure captured by the $l_i + \epsilon_j$ decomposition. We combine this model with a general contact-process framework (Supplementary Methods) in which the individual reproduction number is $v_i = b_i \int_0^\infty g_i(\tau) c_i(t_i + \tau) d\tau$, where b_i is biological infectiousness and $c_i(t)$ is a time-varying contact modulator.

Overdispersion without individual-level heterogeneity. Superspreading arises when secondary infection counts are overdispersed, conventionally parameterized by the negative binomial dispersion parameter k , where smaller k indicates greater overdispersion[Lloyd-Smith et al., 2005]. In the standard framework, overdispersion requires variation in the individual reproduction number v_i , arising from variation in biological infectiousness b_i ("super-shedders") or average contact rates $E_t[c_i(t)]$ ("super-contactors")—both persistent individual traits that are in principle identifiable and targetable. We show that overdispersion can also arise in populations with *no* such individual-level heterogeneity: when all individuals have identical biological infectiousness ($b_i \equiv R_0$) and identical average contact rates ($E_t[c_i(t)] = 1$), narrow infectiousness profiles still generate variation in v_i through the stochastic interaction between each person's infectiousness window and their time-varying contact pattern. A person whose brief burst of infectiousness happens to coincide with a period of elevated contacts produces many secondary infections; one whose burst falls during a contact lull produces few. Crucially, this variation is not attributable to any persistent individual char-

acteristic and therefore cannot be reduced by identifying or targeting high-risk individuals—it represents a floor on overdispersion that persists even after all “super-shedder” and “super-contactor” heterogeneity has been accounted for.

We derived a closed-form expression for the overdispersion parameter under periodic contacts, $c_i(t) = 1 - c_{\text{amp}} \cos(2\pi t/c_{\text{per}})$ (Supplementary Methods):

$$\text{OD} \equiv 1/k = \frac{c_{\text{amp}}^2}{2} \left(1 + \left(\frac{2\pi\bar{g}}{\alpha c_{\text{per}}} \right)^2 \right)^{-\psi\alpha} \quad (1)$$

When contacts are constant ($c_{\text{amp}} = 0$), $\text{OD} = 0$ regardless of ψ : no contact variation, no overdispersion. When contacts vary, narrow infectiousness profiles ($\psi \rightarrow 0$) drive OD toward $c_{\text{amp}}^2/2$, while broad profiles ($\psi \rightarrow 1$) average over the contact variation and suppress overdispersion. The sensitivity of OD to ψ is governed by the ratio of the mean generation time $\bar{g}$ to the contact period c_{per} : weekly contact variation ($c_{\text{per}} = 7$ days) produces ψ -sensitive overdispersion for most respiratory pathogens, whereas daily variation gets averaged out and monthly variation yields near-maximal overdispersion regardless of ψ . Empirical contact surveys provide direct evidence that contacts vary substantially on these relevant timescales: the POLYMOD study[Mossong et al., 2008] and subsequent surveys[?] document 20–40% fewer contacts on weekends than weekdays across European populations, corresponding to $c_{\text{amp}} \approx 0.1$ –0.2 at $c_{\text{per}} = 7$ days. Larger-amplitude variation occurs at the timescale of school terms and holidays. Even at a moderate $c_{\text{amp}} = 0.15$, Eq. 1 predicts $\text{OD} \approx 0.01$ for punctuated profiles ($\psi = 0$) with weekly contacts, which—while modest in isolation—combines multiplicatively with other sources of overdispersion and grows substantially at higher amplitudes or with stochastic contact variation (Supplementary Fig. XX). Previous transmission models have implicitly included the ingredients for this interaction—for example, Goyal et al.[Goyal et al., 2021] coupled narrow peak-infectivity windows (~ 0.5 –1 day) with overdispersed daily contact draws and showed that differing contact distributions explain SARS-CoV-2 superspreading—but attributed overdispersion to the contact process alone rather than isolating the role of infectiousness profile shape.

For three benchmark pathogens—influenza ($R_0 = 2$), SARS-CoV-2 omicron ($R_0 = 6$), and measles ($R_0 = 12$)—simulations confirmed these analytic predictions across a range of contact amplitudes and periods, and showed that the resulting overdispersion translates into systematically higher epidemic extinction probabilities (Fig. ??). Similar patterns emerged for stochastic contact processes with individual-level variation (Supplementary Fig. XX).

Empirical estimates of ψ . We estimated ψ from published transmission-tree data for eight diseases: COVID-19, MERS, measles, Ebola, smallpox, pneumonic plague, norovirus, and hepatitis A. We extracted serial intervals from the OutbreakTrees database[?], supplemented with COVID-19 data from refs. ? and ? . Serial intervals were grouped by index case to form clusters (restricted to ≥ 2 secondary cases per cluster). For each pathogen, we fixed the generation interval distribution to published estimates and placed a flat prior on ψ ; the likelihood was computed by integrating over the shared latent period within each cluster (Methods).

The resulting estimates spanned a wide range (Fig. ??). Measles and pneumonic plague yielded strongly punctuated estimates ($\hat{\psi} \approx 0$, 95% CI [0.00, 0.11] and [0.00, 0.20]), consistent with secondary infections tightly clustered in time. COVID-19, smallpox, and MERS yielded diffuse estimates ($\hat{\psi} > 0.7$), consistent with infections spread broadly across each index case’s infectious period.

These estimates should be regarded as hypothesis-generating rather than definitive. The available data are subject to ascertainment bias toward large clusters and superspreading events (which can arise from event-based transmission rather than genuinely narrow infectiousness windows), incomplete observation of infectious periods, date discretization to symptom onset day, and sensitivity to the assumed generation interval parameters (Supplementary Discussion). The most strongly punctuated estimates—measles and pneumonic plague—are also the most susceptible to these biases, since transmission of both pathogens is often event-mediated and well-traced. Conversely, local susceptible depletion in high- R_0 settings could mimic low ψ : an index case may rapidly exhaust nearby susceptibles upon reaching peak infectiousness, yielding tightly clustered secondary infections even if the biological window of infectiousness is broad. Disentangling such epidemiological effects from the intrinsic shape of $a_i(\tau)$ will require data from settings where these confounds are minimal, such as household studies with complete ascertainment or controlled exposure experiments. Nevertheless, the breadth of the estimated range—from near 0 to near 1—suggests that ψ varies meaningfully across pathogens. Independent evidence from viral kinetics modelling supports narrow infectivity windows: Goyal et al.[Goyal et al., 2021] showed that the dose-response relationship between viral load and transmission probability restricts the window of high transmission risk to ~ 0.5 –1 day for both SARS-CoV-2 and influenza, consistent with low ψ . The biological plausibility of variation in ψ across pathogens—brief shedding bursts *vs.* sustained infectiousness—motivates investment in the data needed to resolve these estimates.

A simulation-based power analysis indicates that ψ is identifiable from contact-tracing data under reasonably favourable conditions: 50 index cases sufficed for well-powered inference for SARS-CoV-2 and measles across a range of observation delays (Supplementary Fig. XX). Inference was more challenging for influenza ($R_0 = 2$), where fewer secondary cases per index case limit within-cluster information.

Intervention sensitivity. Detect-and-isolate (D&I) interventions aim to identify infected individuals and prevent onward transmission. Their efficacy depends on the timing of detection relative to infectiousness. When detection timing is anchored to the individual infectiousness profile—as is realistic for symptom-based or viral-load-based detection—the shape of $a_i(\tau)$ determines how sensitively the intervention responds to shifts in detection timing.

We defined testing effectiveness (TE) as $TE = \rho\eta P(\tau > D)$, where ρ is the participation rate, η is isolation effectiveness, and D is the detection time[Middleton and Larremore, 2024]. For symptom-based detection (symptoms arising at a $\text{Normal}(\delta_{\text{symp}}, \sigma_{\text{symp}}^2)$ offset from peak infectiousness) and test-based screening (tests at fixed intervals with a detectability window around peak infectiousness), we computed TE by numerical integration (Methods). For both detection mechanisms, TE dropped as detection shifted from before to after peak infectiousness, with the sharpest drops for narrow infectiousness profiles (Fig. ??a,b). These sharp transitions reflect the all-or-nothing character of punctuated profiles: detection either catches the entire burst of infectiousness or misses it entirely.

Beyond TE, mode-anchored D&I interventions produce two additional ψ -dependent effects. First, among individuals who escape detection, the effective generation interval $g^*(\tau)$ is truncated. For smooth profiles, detection reliably removes late transmission, shortening the mean generation interval; for punctuated profiles, detection is all-or-nothing, so survivors transmit with an untruncated $g^*(\tau) = g(\tau)$. This truncation can accelerate epidemic growth among those who escape detection (Supplementary Methods). Second, variation in detection timing across individuals creates variation in the individual reproduction number $\nu_i^* = \nu_i F_e(m_e + \iota_i)$, where F_e is the CDF of the infectiousness kernel and ι_i is the individual’s isolation time relative to peak infectiousness. This introduces overdispersion in offspring counts, the magnitude of which

depends on ψ and the detection timing distribution (Supplementary Fig. XX).

Gathering size restrictions also interact with ψ . Truncating the contact distribution at a maximum gathering size $c_{\max}$ reduces R_{eff} by a factor μ_T that is independent of ψ (Supplementary Methods). However, the truncation also reduces overdispersion, and the absolute reduction is greatest when the baseline overdispersion is highest—*i.e.*, when ψ is small. Simulations confirmed that gathering-size-restricted epidemics had systematically higher extinction probabilities than R_{eff} -matched epidemics without contact-process adjustment, with the largest discrepancies for narrow infectiousness profiles (Fig. ??c). Evaluating interventions purely in terms of their effect on R_{eff} can therefore be misleading; the full distribution of secondary infections—not just its mean—determines the epidemic consequences.

Further effects on epidemic dynamics. Narrow infectiousness profiles also inflate early-epidemic stochasticity. In the branching-process approximation to early-epidemic growth, the cumulative infection count can be written as $Z(t) \approx e^{r(t+\varsigma)}$, where ς is a random time shift [Morris et al., 2024]. We showed analytically that $\text{Var}[\varsigma]$ increases as $\psi \rightarrow 0$, with $\text{Var}[W] \propto c(R_0, \alpha)^{(1-\psi)\alpha}$ for a constant $c > 1$ (Supplementary Methods). Simulations confirmed that punctuated profiles produced wider distributions of epidemic establishment times: by day 20, only 55% of SARS-CoV-2 epidemics with $\psi = 0$ had reached 5% cumulative prevalence, compared to 82% with $\psi = 1$ (Supplementary Fig. XX). Additionally, serial intervals from the same index case are correlated when $\psi < 1$, yielding an effective sample size $n_{\text{eff}} = n / (1 + (R_0 - 1) \cdot \text{ICC})$ for generation interval estimation, where the intraclass correlation ICC increases as $\psi \rightarrow 0$. Treating serial intervals as independent, as is common practice, can produce overconfident estimates of $g(\tau)$'s parameters (Supplementary Fig. XX).

Implications. Our results demonstrate that the shape of the individual infectiousness profile—an epidemiological quantity that is invisible to deterministic, mean-field models—can meaningfully impact epidemic dynamics and intervention effectiveness. The parameter ψ captures an axis of variation complementary to the dispersion parameter k introduced by Lloyd-Smith et al. [Lloyd-Smith et al., 2005]: whereas k describes how much individuals vary in the *amount* of transmission, ψ describes how much they vary in its *timing*. The two interact: the timing-contact mechanism we identify generates overdispersion ($k < \infty$) even in a population of identical shedders with identical average contact rates, setting a *floor* on the overdispersion that remains after all individual-level risk factors have been accounted for. This has a direct implication for outbreak control: individual-specific interventions that target biological or behavioural superspreaders—which Lloyd-Smith et al. showed can dramatically outperform population-wide measures—cannot address the timing-driven component of overdispersion, because it does not arise from any identifiable individual trait. The long-noted qualitative distinction between superspreading *events* and superspreader *individuals* [Lloyd-Smith et al., 2005] maps onto our framework: individual-driven overdispersion (variation in b_i or $\bar{c}_i$) produces superspreader individuals, while timing-driven overdispersion (low ψ with variable contacts) produces superspreading events. Anecdotal observations support the latter mechanism—individuals who generate large clusters in one setting yet fail to infect household contacts are difficult to reconcile with persistent individual-level superspreading, but are expected when infectiousness is concentrated in a brief window that may or may not coincide with high-contact periods. Quantifying the relative contributions of biological variation, contact variation, and timing-contact interaction to observed overdispersion is an important avenue for future work; our framework provides the tools to do so.

Several concrete recommendations follow. First, contact-tracing studies should record the identity of the index case for each observed serial interval, enabling cluster-level analyses that correctly account for

within-cluster correlation. Second, projection and forecasting models should incorporate ψ —even as a sensitivity parameter—to account for a source of uncertainty that is currently ignored. Third, evaluations of D&I interventions should assess the sensitivity of testing effectiveness to small shifts in detection timing, which can produce disproportionate changes in epidemic outcomes when infectiousness profiles are narrow. Better data are needed: longitudinal viral kinetics studies linked to transmission outcomes, household studies capturing full sets of secondary infections, and controlled exposure experiments could all inform estimates of $a_i(\tau)$ ’s shape. We hope this framework will motivate such data collection and support more nuanced assessments of epidemic risk.

Methods

Simulation framework. We used stochastic, individual-based epidemic simulations implemented in R (version 4.5.0) with performance-critical loops in C++ via Rcpp. For each infected individual, the number of secondary infection attempts was drawn from $\text{Poisson}(R_0)$, with infection times $\tau_{ij} = l_i + \epsilon_j$ under the Gamma time-shifted model. Time-varying contacts were incorporated via Poisson thinning. We simulated both finite-population ($N = 10,000$) and infinite-population (branching process) variants. Full simulation details are in Supplementary Methods.

Benchmark pathogens. We used generation interval parameters from the literature: influenza ($\bar{g} = 3.2$ days, $\alpha = 2.32$, $R_0 = 2$; ref. 3), SARS-CoV-2 omicron ($\bar{g} = 7.05$ days, $\alpha = 2.39$, $R_0 = 6$; ref. 7), and measles ($\bar{g} = 12.2$ days, $\alpha = 11.36$, $R_0 = 12$; ref. 5).

Overdispersion. Overdispersion was quantified by fitting a negative binomial distribution to simulated offspring counts via moment matching ($k = R_0^2 / (\text{Var}[\chi] - R_0)$). The closed-form expression (Eq. 1) was derived using the power spectral density of the contact process filtered through the individual infectiousness kernel (Supplementary Methods).

Testing effectiveness. TE was computed by numerical integration over the detection time distribution and the infectiousness kernel, for both symptom-based ($D \sim \text{Normal}(\delta_{\text{symp}}, \sigma_{\text{symp}}^2)$ relative to peak infectiousness) and test-based (fixed-interval screening with detectability window) detection mechanisms (Supplementary Methods).

Empirical ψ estimation. Transmission trees were obtained from the OutbreakTrees database[?], supplemented with COVID-19 data from refs. ? and ?. Serial intervals were grouped by index case into clusters (≥ 2 secondary cases). For each pathogen, the likelihood was computed as a product over clusters of the Gamma time-shifted density, integrating over the shared latent period l_i , with generation interval parameters (α, β) fixed from published estimates. A flat prior on $\psi \in [0, 1]$ was evaluated on a grid of 101 points. Posterior mode, mean, and 95% credible intervals were reported.

Power analysis. We simulated serial interval data for 10, 25, 50, and 100 index cases at $\psi \in \{0, 0.5, 1\}$ for each benchmark pathogen, with Gamma-distributed observation delays (mean 1, 2, or 4 days). Posterior estimates for ψ were generated using a flat prior. Power was defined as $> 80\%$ of 200 replicates assigning $\geq 95\%$ posterior mass within 0.2 of the true ψ .

References

- D. Breda, O. Diekmann, W. F. De Graaf, A. Pugliese, and R. Vermiglio. On the formulation of epidemic models (an appraisal of Kermack and McKendrick). *Journal of Biological Dynamics*, 6(sup2):103–117, September 2012. ISSN 1751-3758, 1751-3766. doi: 10.1080/17513758.2012.716454. URL <http://www.tandfonline.com/doi/abs/10.1080/17513758.2012.716454>.
- David Champredon and Jonathan Dushoff. Intrinsic and realized generation intervals in infectious-disease transmission. *Proceedings of the Royal Society B: Biological Sciences*, 282(1821):20152026, December 2015. ISSN 0962-8452, 1471-2954. doi: 10.1098/rspb.2015.2026. URL <https://royalsocietypublishing.org/doi/10.1098/rspb.2015.2026>.
- Louis Yat Hin Chan, Sinead E. Morris, Melissa S. Stockwell, Natalie M. Bowman, Edwin Asturias, Suchitra Rao, Karen Lutrick, Katherine D. Ellingson, Huong Q. Nguyen, Yvonne Maldonado, Son H. McLaren, Ellen Sano, Jessica E. Biddle, Sarah E. Smith-Jeffcoat, Matthew Biggerstaff, Melissa A. Rolfes, H. Keipp Talbot, Carlos G. Grijalva, Rebecca K. Borchering, and Alexandra M. Mellis. Estimating the generation time for influenza transmission using household data in the United States. *Epidemics*, 50:100815, March 2025. ISSN 17554365. doi: 10.1016/j.epidem.2025.100815. URL <https://linkinghub.elsevier.com/retrieve/pii/S1755436525000039>.
- Ashish Goyal, Daniel B Reeves, E Fabian Cardozo-Ojeda, Joshua T Schiffer, and Bryan T Mayer. Viral load and contact heterogeneity predict SARS-CoV-2 transmission and super-spreading events. *eLife*, 10: e63537, February 2021. ISSN 2050-084X. doi: 10.7554/eLife.63537. URL <https://elifesciences.org/articles/63537>.
- Don Klinkenberg and Hiroshi Nishiura. The correlation between infectivity and incubation period of measles, estimated from households with two cases. *Journal of Theoretical Biology*, 284(1):52–60, September 2011. ISSN 00225193. doi: 10.1016/j.jtbi.2011.06.015. URL <https://linkinghub.elsevier.com/retrieve/pii/S0022519311003146>.
- J. O. Lloyd-Smith, S. J. Schreiber, P. E. Kopp, and W. M. Getz. Superspreading and the effect of individual variation on disease emergence. *Nature*, 438(7066):355–359, November 2005. ISSN 0028-0836, 1476-4687. doi: 10.1038/nature04153. URL <https://www.nature.com/articles/nature04153>.
- Mattia Manica, Alfredo De Bellis, Giorgio Guzzetta, Pamela Mancuso, Massimo Vicentini, Francesco Venturelli, Alessandro Zerbini, Eufemia Bisaccia, Maria Litvinova, Francesco Menegale, Carla Molina Grané, Piero Poletti, Valentina Marziano, Agnese Zardini, Valeria d’Andrea, Filippo Trentini, Antonino Bella, Flavia Riccardo, Patrizio Pezzotti, Marco Ajelli, Paolo Giorgi Rossi, and Stefano Merler. Intrinsic generation time of the SARS-CoV-2 Omicron variant: An observational study of household transmission. *The Lancet Regional Health - Europe*, 19:100446, August 2022. ISSN 26667762. doi: 10.1016/j.lanepe.2022.100446. URL <https://linkinghub.elsevier.com/retrieve/pii/S2666776222001405>.
- Casey Middleton and Daniel B. Larremore. Modeling the transmission mitigation impact of testing for infectious diseases. *Science Advances*, 10(24):eadk5108, June 2024. ISSN 2375-2548. doi: 10.1126/sciadv.adk5108. URL <https://www.science.org/doi/10.1126/sciadv.adk5108>.
- Dylan Morris, John Maclean, and Andrew J. Black. Computation of random time-shift distributions for stochastic population models. *Journal of Mathematical Biology*, 89(3):33, September 2024. ISSN

304 0303-6812, 1432-1416. doi: 10.1007/s00285-024-02132-6. URL <https://link.springer.com/10.1007/s00285-024-02132-6>.
305

306 Dylan J. Morris, Lauren Kennedy, and Andrew J. Black. Random time-shift approximation enables hier-
307 archical Bayesian inference of mechanistic within-host viral dynamics models on large datasets. *arXiv*
308 *preprint arXiv:2507.02884*, 2025. URL <https://arxiv.org/abs/2507.02884>.

309 Joël Mossong, Niel Hens, Mark Jit, Philippe Beutels, Kari Auranen, Rafael Mikolajczyk, Marco Massari,
310 Stefania Salmaso, Gianpaolo Scalia Tomba, Jacco Wallinga, Janneke Heijne, Malgorzata Sadkowska-
311 Todys, Magdalena Rosinska, and W. John Edmunds. Social Contacts and Mixing Patterns Relevant
312 to the Spread of Infectious Diseases. *PLoS Medicine*, 5(3):e74, March 2008. ISSN 1549-1676. doi:
313 10.1371/journal.pmed.0050074. URL <https://dx.plos.org/10.1371/journal.pmed.0050074>.

314 Åke Svensson. A note on generation times in epidemic models. *Mathematical Biosciences*, 208(1):300–311,
315 July 2007. ISSN 00255564. doi: 10.1016/j.mbs.2006.10.010. URL <https://linkinghub.elsevier.com/retrieve/pii/S0025556406002094>.
316

317 J Wallinga and M Lipsitch. How generation intervals shape the relationship between growth rates and
318 reproductive numbers. *Proceedings of the Royal Society B: Biological Sciences*, 274(1609):599–604, February
319 2007. ISSN 0962-8452, 1471-2954. doi: 10.1098/rspb.2006.3754. URL <https://royalsocietypublishing.org/doi/10.1098/rspb.2006.3754>.
320

321 Figure Legends

322 **Figure 1 | The individual infectiousness profile and the smoothness parameter ψ .** **a**, The population-
 323 level infectiousness profile $A(\tau)$ (black curve) can arise from different individual-level profiles $a_i(\tau)$. Left:
 324 stepwise profiles (SEIR model). Centre: smooth profiles matching the population average. Right: spike pro-
 325 files concentrated at single moments. All produce the same $A(\tau)$ and thus the same deterministic epidemic
 326 dynamics. **b**, The Gamma time-shifted model parameterizes this decomposition with $\psi \in [0, 1]$. Secondary
 327 infections from index case i occur at times $\tau_{ij} = l_i + \epsilon_j$, where $l_i \sim \text{Gamma}((1 - \psi)\alpha, \beta)$ is a shared latent
 328 period and $\epsilon_j \sim \text{Gamma}(\psi\alpha, \beta)$ is independent jitter. Examples shown for three values of ψ .

329 **Figure 2 | Narrow infectiousness profiles generate overdispersion through contact-timing interaction.**
 330 **a–c**, Schematic of the g_i – c_i interaction mechanism. A narrow infectiousness window (left) can coincide with
 331 high or low contacts, producing variable offspring counts; a broad window (right) averages over contact
 332 variation. **d–f**, Estimated overdispersion ($\text{OD} = 1/k$) from simulated offspring counts (10,000 index cases
 333 per scenario) as a function of ψ and contact amplitude (c_{amp}), for periodic contacts with $c_{\text{per}} = 7$ days.
 334 Rows: influenza, SARS-CoV-2, measles. **g–i**, As in **d–f**, for stochastic contacts with varying dispersion
 335 parameter σ .

336 **Figure 3 | Empirical estimates of ψ across eight pathogens.** Posterior estimates (mode and 95% credi-
 337 ble interval) of ψ from transmission-tree data. Sample sizes (number of serial intervals) shown at right.
 338 Generation interval parameters fixed from published estimates (Supplementary Table XX).

339 **Figure 4 | Sensitivity of interventions to individual infectiousness timing.** **a**, Testing effectiveness (TE)
 340 under symptom-based detection as a function of the mean symptom-to-peak offset δ_{symp} , for three values
 341 of ψ . Rows: benchmark pathogens. **b**, As in **a**, for test-based screening as a function of test interval Δ_{test} . **c**,
 342 Epidemic extinction probability under gathering size restrictions (truncated stochastic contacts, solid lines)
 343 compared with R_{eff} -matched epidemics without contact-process adjustment (dashed lines), for three values
 344 of ψ .